CATALYTIC AND CO-ORDINATION CHEMISTRY OF ENOLATE
OXAZOLINE κ^2 -N,O-LIGANDS TO GROUP 9 & 11 TRANSITION
METALS

by

Kathleen L. May
Bachelor of Science, Chemistry
Ryerson University, Toronto, Ontario, Canada 2016

A thesis presented to Ryerson University
in partial fulfillment of the
requirements for the degree of
Master of Science
in the program of
Molecular Science

Toronto, Ontario, Canada 2018
© Kathleen L. May, 2018

AUTHOR'S DECLARATION FOR ELECTRONIC SUBMISSION OF A THESIS

I hereby declare that I am the sole author of this thesis. This is a true copy of the thesis, including any required final revisions, as accepted by my examiners.

I authorize Ryerson University to lend this thesis to other institutions or individuals for the purpose of scholarly research.

I further authorize Ryerson University to reproduce this thesis by photocopying or by other means, in total or in part, at the request of other institutions or individuals for the purpose of scholarly research.

I understand that my thesis may be made electronically available to the public.

CATALYTIC AND CO-ORDINATION CHEMISTRY OF ENOLATE OXAZOLINE κ^2 - N,O-LIGANDS TO GROUP 9 & 11 TRANSITION METALS

Master of Science, 2018

Kathleen May

Molecular Science

Ryerson University

Abstract

The co-ordination chemistry and resulting catalytic potential of Group 9 and 11 transition metal centered complexes with the 2-acylmethyl-2-oxazoline skeleton are detailed. Each of the synthesized ligand species were treated with either a Copper (II) or Cobalt (II) salt to promote co-ordination, in all cases deprotonation of the ligand occurred. These complexes have, thereafter, been examined by infrared spectroscopy (IR), UV-Visible spectroscopy (UV-Vis), mass spectrometry (MS), cyclic voltammetry (CV), combustion analysis (EA) and x-ray diffraction and probed for their potential redox properties. In the case of Cu(II) chelated complexes, no desirable redox behaviour was observed. Although, with respect to Co(II) complexes, one complex displayed favourable redox potential. The redox active species have been shown to effectively catalyze the polymerization of methyl methacrylate with tosyl chloride as an initiator, through an ATRP-like mechanism. Work within also reflects preliminary co-ordination to Iridium and Rhodium metal centers. Successfully synthesized Iridium complexes have been tested with respect to their oxidative properties. Positive results have been observed in their ability to perform oxidative addition with both HSnPh_3 and MeI . Both small molecule species have been effectively added to an Ir(I) complex, formally changing its oxidation state to Ir(III).

Acknowledgements

Firstly, I'd like to thank Prof. Gossage for taking me as a graduate student. Mainly I'd like to thank him for putting up with my naïve excitement and overall over-the-top attitude and enthusiasm when it comes to our research, not only for my master's degree but also for my doctoral studies (in which I hope he can continue to put up with me). He has been an excellent mentor in both my research as well as helped me to grow as a writer.

Thank you to all my committee members (Dr. B. Koivisto, and Dr. D. Foucher) for literally grinding information into my brain. I've learned so much over the last couple years and am very lucky to have met such intelligent individuals such as yourselves.

All my lab mates, past and present; Jeff, Joe, Jas, Gloria, Kristy, Rachel, Jeanette, Tristan, and Alex. most importantly, the other two-thirds of the **OG3**; Des and Julie. Without any of these people I could have accomplished much more in my thesis since there would have been less late UNO/Anomia/Fun Farm/Killer game nights, Blue Jays games, and memorable family lunches. Although, I wouldn't have it any other way.

To my parents, for they let their oldest daughter move out of the house to live closer to the lab with some guy she literally met 2 months prior to moving out. They have supported me in more ways than I can fathom. I love you, Mom, Dad and my awesome little sister Sarah (please don't grow up).

Lastly, I'd like to thank my loving boyfriend, William Scannell. For all those countless nights watching dumb YouTube videos, playing bad LAN video games, all the fun times we've had and all the laughs we shared. I truly could not have asked for a better friend, even though you severely lack creativity when it comes to making birthday/anniversary cards.

Table of Contents:

Abstract.....	iii
Acknowledgements	iv
List of Abbreviations:.....	xiii
List of Figures:	xvii
List of Schemes:.....	xx
List of Equations:	xxiii
List of Tables:	xxiv
List of Appendices	xxv
1.0 Introduction.....	1
1.1 Oxazolines	1
1.2 Coordination Chemistry	2
1.2.1 Denticity	3
1.3. κ^2 -N,O Chelating Ligands	4
1.3.1 Classes of N,O Chelating Ligands	5
1.3.2 Synthesis of Ligands	5
1.3.2.1 Ligands via Diacylation and Hydrolysis	6
1.3.2.2 Ligands via Crossed-Claisen Reaction	7
1.3.2.3 Ligands via S,S-dithio-Ketenenes	7

1.3.2.4 Ligands via alternative methods – pyridine/lithiation	8
1.3.3 <i>Tautomers of the Target Ligand Series</i>	9
1.4 Relevant Transition Metal N,O-complexes	10
1.4.1 <i>Copper(II)/Cobalt(II/III) Complexes</i>	10
1.4.2 <i>Rhodium(I/III)/Iridium(I/III) Complexes</i>	11
1.5 Polymers	12
1.5.1 <i>Mechanisms of Polymer Growth</i>	13
1.5.2 <i>Metal Catalyzed Atom Transfer Radical Polymerization (ATRP)</i>	14
1.5.2.1 Copper(I/II) Catalysts in ATRP	15
1.5.2.2 Cobalt(II/III) Catalysts in ATRP	16
1.6 Past Work	17
1.7 Objective of This Thesis	18
2.0 Results and Discussion	19
2.1 General Considerations	19
2.2 Synthesis and Characterization of Intermediate Compounds (3a-i , 5a-d)	19
2.3 Synthesis and Characterization of a Ligand Series with 1a (4a-i and 4j-l)	26
2.3.1 <i>Synthesis and Characterization of ligands 4a-i</i>	26
2.3.2 <i>Synthesis and Characterization of ligands 4j-l</i>	29
2.4 Synthesis and Characterization of a Ligand Series with 2-methyl-2-oxazoline (6a-d)	33
2.5 Synthesis and Spectrochemical Investigation of Transition Metal Complexes	37
2.5.1 <i>Copper Complexes (7a-l)</i>	39

2.5.2 Other Copper Complexes (8a-c).....	47
2.5.3 Cobalt Complexes (9a-l).....	49
2.5.3.1 Attempted Oxidation of a Co^{2+} Complex.....	55
2.5.4 Cobalt Complexes (10a-d and 11a/11c).....	59
2.5.4.1 Usual Oxidation from Co^{2+} to Co^{3+} Complexes (11a/c).....	63
2.6 Polymerization Studies with a Co^{2+} Catalyst.....	67
2.7 Synthesis and Characterization Other Group 9 complexes.....	70
2.7.1 Attempted Synthesis and Characterization of Rh Complexes.....	70
2.7.2 Attempted Synthesis and Characterization of Vaska-type Complexes.....	71
2.7.3 Synthesis and Characterization of Ir Complexes (14 and 15).....	74
2.7.3.1 Oxidation Addition of various substrates to 15 (16-18)	79
2.7.3.2 Attempted displacement of the COD ring of complex 15 (19a-e)	88
3.0 Conclusions and Future directions:	90
4.0 Experimental	92
4.1 General Considerations	92
4.2 Synthesis and Characterization of Intermediates species (3a-h).....	94
4.2.1 Synthesis of (E)-2-(3-benzoyl-4,4-dimethyloxazolidin-2-ylidene)-1-phenylethan-1-one (3a).....	94
4.2.2 Synthesis of (Z)-2-(3-(3,4-dichlorobenzoyl)-4,4-dimethyloxazolidin-2-ylidene)-1-(3,4-dichlorophenyl)ethan-1-one (3b).....	95

4.2.3 (E)-2-(3-(4-methoxybenzoyl)-4,4-dimethyloxazolidin-2-ylidene)-1-(4-methoxyphenyl)ethan-1-one (3c)	96
4.2.4 Synthesis of (Z)-2-(3-(3,4-dimethylbenzoyl)-4,4-dimethyloxazolidin-2-ylidene)-1-(3,4-dimethylphenyl)ethan-1-one (3d)	97
4.2.5 Synthesis of (Z)-2-(3-(4-chloro-3-methylbenzoyl)-4,4-dimethyloxazolidin-2-ylidene)-1-(4-chloro-3-methylphenyl)ethan-1-one (3e).....	98
4.2.6 Synthesis of (Z)-2-(3-(3-chloro-4-methylbenzoyl)-4,4-dimethyloxazolidin-2-ylidene)-1-(3-chloro-4-methylphenyl)ethan-1-one (3f)	99
4.2.7 (Z)-(2-(2-hydroxy-2-(naphthalen-1-yl)vinyl)-4,4-dimethyl-4,5-dihydro-3H-oxazol-3-yl)(naphthalen-1-yl)methanone (3g).....	100
4.2.8 (E)-1-(4,4-dimethyl-3-pivaloyloxazolidin-2-ylidene)-3,3-dimethylbutan-2-one. (3h) ^{20,35}	101
4.3 Synthesis and Characterization for the Ligand Series made with 1a	102
4.3.1 Synthesis of (Z)-2-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)-1-phenylethen-1-ol (4a)	102
4.3.2 (E)-1-(3,4-dichlorophenyl)-2-(4,4-dimethyloxazolidin-2-ylidene)ethan-1-one (4b)	103
4.3.3 Synthesis of (E)-2-(4,4-dimethyloxazolidin-2-ylidene)-1-(4-methoxyphenyl)ethan-1-one (4c)	104
4.3.4 Synthesis of (E)-2-(4,4-dimethyloxazolidin-2-ylidene)-1-(3,4-dimethylphenyl)ethan-1-one (4d).....	105

4.3.5 Synthesis of (E)-1-(4-chloro-3-methylphenyl)-2-(4,4-dimethyloxazolidin-2-ylidene)ethan-1-one (4e)	106
4.3.6 Synthesis of (E)-1-(3-chloro-4-methylphenyl)-2-(4,4-dimethyloxazolidin-2-ylidene)ethan-1-one (4f)	107
4.3.7 Synthesis of (E)-2-(4,4-dimethyloxazolidin-2-ylidene)-1-(naphthalen-1-yl)ethan-1-one (4g).....	108
4.3.8 Synthesis of (E)-1-(4,4-dimethyloxazolidin-2-ylidene)-3,3-dimethylbutan-2-one (4h).....	109
4.3.9 Synthesis of (E)-1-(4,4-dimethyloxazolidin-2-ylidene)propan-2-one (4j).....	110
4.3.10 Synthesis of (E)-3-(4,4-dimethyloxazolidin-2-ylidene)-1,1,1-trifluoropropan-2-one (4k)	111
4.3.11 Synthesis of (E)-1,1,1-trichloro-3-(4,4-dimethyloxazolidin-2-ylidene)propan-2-one (4l)	112
4.4 Synthesis of the ligand series with 1b	113
4.4.1 Synthesis of (E)-2-(oxazolidin-2-ylidene)-1-phenylethan-1-one (5a to 6a);.....	113
4.4.2 Synthesis of (E)-1-(3,4-dichlorophenyl)-2-(oxazolidin-2-ylidene)ethan-1-one (5b to 6b).....	114
4.4.3 Synthesis of (E)-1-(4-methoxyphenyl)-2-(oxazolidin-2-ylidene)ethan-1-one (5c to 6c)	115
4.4.4 Attempted Synthesis of (E)-1-(4-nitrophenyl)-2-(oxazolidin-2-ylidene)ethan-1-one (5d/6d)	116
4.5 Synthesis and Characterization of Cu complexes made with 1a	117

4.5.1 Chelation of 4a to Copper metal to obtain 7a	117
4.5.2 Chelation of 4b to Copper metal to obtain 7b	118
4.5.3 Chelation of 4c to Copper metal to obtain 7c	119
4.5.4 Chelation of 4d to Copper metal to obtain 7d	120
4.5.5 Chelation of 4e to Copper metal to obtain 7e ;	121
4.5.6 Chelation of 4f to Copper metal to obtain 7f ;.....	122
4.5.7 Chelation of 4g to Copper metal to obtain 7g ;.....	123
4.5.8 Chelation of 4h to Copper metal to obtain 7h	124
4.5.9 Chelation of 4i to a Copper metal to obtain 7i ;	125
4.6 Synthesis and Characterization of Cu complexes made with 1b	126
4.6.1 Chelation of 6a to Copper metal to obtain 8a	126
4.6.2 Chelation of 6b to Copper metal to obtain 8b	127
4.7 Synthesis and Characterization of Co complexes made with 1a	128
4.7.1 Chelation of 4a to a Cobalt metal to obtain 9a	128
4.7.2 Chelation of 4b to a Cobalt metal to obtain 9b	129
4.7.3 Chelation of 4c to a Cobalt metal to obtain 9c	130
4.7.4 Chelation of 4d to a Cobalt metal to obtain 9d	131
4.7.5 Chelation of 3e to a Cobalt metal to obtain 9e	132
4.7.6 Chelation of 4f to a Cobalt metal to obtain 9f	133
4.7.7 Chelation of 4g to a Cobalt metal to obtain 9g	134
4.7.8 Chelation of 4h to a Cobalt metal to obtain 9h	135
4.7.9 Chelation of 4i to a Cobalt metal to obtain 9i	136

4.7.10 Chelation of 4j to a Cobalt metal to obtain 9j	137
4.7.11 Chelation of 4k to a Cobalt metal to obtain 9k	138
4.7.12 Chelation of 4l to a Cobalt metal to obtain 9l	139
4.8 Synthesis and Characterization of Co complexes made with 1a	140
4.8.1 Chelation of Cobalt metal to obtain 10a	140
4.8.1.1 Oxidation of 10a to 11a	141
4.8.2 Chelation of Cobalt metal to obtain 10b	142
4.8.3 Chelation of Cobalt metal to obtain 10c	143
4.8.3.1 Oxidation of 10c to 11c	144
4.9 Attempted Oxidation of Co complexes made with 1a	145
4.9.1 Attempted oxidation of 9a to 9aX ;	145
4.9.2 Attempted oxidation of 9h to 9h-1 and then 9h-2	146
4.10 Using 9h in ATRP	147
4.10.1 Polymerization of MMA with 9h to PMMA	147
4.10.2 Attempted polymerization of THF with 9h	148
4.11 Synthesis of Other Group 9 Complexes	149
4.11.1 Synthesis of Vaska's complex ($\text{IrCl}(\text{CO})(\text{PPh}_3)_2$) (12)	149
4.11.2 Synthesis of bis(1,5-cyclooctadiene)diiridium(I)dichloride (14)	149
4.11.3 Substitution of 14 with 4a to give 15	150
4.11.4 Oxidative addition of methyl iodide to 15	151
4.11.5 Synthesis and oxidative addition of triphenylstannane to 15	152
Appendix	153

References	512

List of Abbreviations:

Acetonitrile	ACN
Aluminum(III)chloride	AlCl ₃
Aryl	Ar
Atom transfer radical polymerization	ATRP
Benzene-d ₆	C ₆ D ₆
<i>tertiary</i> -Butyl	<i>t</i> -Bu
<i>normal</i> -Butyllithium	<i>n</i> -BuLi
2,2'-Bipyridine	Bpy
Carbon disulphide	CS ₂
Chloroform	CHCl ₃
Chloroform-d ₁	CDCl ₃
Correlation spectroscopy	COSY
Cobalt(II) bromide	CoBr ₂
Cobalt(II) hydroxide	Co(OH) ₂
Cobalt(II) nitrate hexahydrate	Co(NO ₃) ₂ • 6 H ₂ O
Copper(II) bromide	CuBr ₂
Copper(II) nitrate hemipentahydrate	Cu(NO ₃) ₂ • 2.5 H ₂ O
Cyclic voltammery	CV
<i>cis,cis</i> -1,5-Cyclooctadiene	COD
Day(s)	d
Daltons	Da
Dichloromethane	DCM

Dimethyl formamide	DMF
Dimethyl sulphoxide	DMSO
Distilled water	dH ₂ O
1,1'-bis(Diphenylphosphino)ferrocene	dppf
Diethyl ether	Et ₂ O
Elemental Analysis	EA
Electron volts	eV
Ethanol	EtOH
Ethyl acetate	EtOAc
Gel Permeation Chromatography	GPC
Heteronuclear multiple-bond correlation spectroscopy	HMBC
Heteronuclear single- quantum correlation spectroscopy	HSQC
Hertz	Hz
Infrared spectroscopy	IR
Ligand to metal charge transfer	LMCT
Magnesium sulphate	MgSO ₄
Mass spectrometry	MS
Melting point	mp
Metal to ligand charge transfer	MLCT
Methanol	MeOH
Methyl iodide	MeI
Methyl methacrylate	MMA
N-bromosuccinamide	NBS

1-Naphthyl	nap
Nuclear magnetic resonance spectroscopy	NMR
Nuclear Overhauser effect spectroscopy	nOesy
The Oak Ridge Thermal Ellipsoid Plot	ORTEP
Parts per million	ppm
Poly(aniline)	PANi
Poly(methyl methacrylate)	PMMA
Poly(styrene)	PS
Poly(vinylchloride)	PVC
Polydispersity Index	PDI
Potassium hydroxide	KOH
Pyridine	py
Reversible Addition-Fragmentation Chain Transfer	RAFT
Room temperature	rt
Sodium carbonate	Na ₂ CO ₃
Sodium chloride	NaCl
Sodium sulphate	Na ₂ SO ₄
Sulphuric acid	H ₂ SO ₄
Tetrabutylammonium hexafluorophosphate	[N(n-Bu) ₄ ⁺][PF ₆ ⁻]
Tetrahydrofuran	THF
N, N, N', N'-Tetramethylethylenediamine	TMEDA
(2,2,6,6-Tetramethylpiperidin-1-yl)oxyl	TEMPO
Thin Layer Chromatography	TLC

Thionyl chloride	SOCl_2
Trichloroacetyl chloride	TCAC
Triethylamine	NEt_3
Triethylamine hydrochloride	$\text{NEt}_3 \cdot \text{HCl}$
Triethylphosphite	$\text{P}(\text{OEt})_3$
Trichlorosilane	HSiCl_3
Trifluoroacetic anhydride	TFAA
Triphenylphosphine	PPh_3
Triphenylphosphine oxide	$\text{O}=\text{PPh}_3$
Triphenylsilane	HSiPh_3
Triphenylstannane	HSnPh_3
Tosyl chloride	TsCl
Ultraviolet-Visible spectroscopy	UV-Vis
Vaska's complex	$\text{IrCl}(\text{CO})(\text{PPh}_3)_2$
Zinc Chloride	ZnCl_2

List of Figures:

Figure 1: Crystal structure of D , seen in the enamine form.....	9
Figure 2: One of the intermediate compounds, 3e , labeled for aid in NMR discussion.	21
Figure 3: ^1H -NMR spectrum of 3e	21
Figure 4: ^{13}C -NMR spectrum for 3e	22
Figure 5: <u>A</u> - Prior to work-up, before filtering the salt ($[\text{NEt}_3][\text{HCl}]$), <u>B</u> - After work-up and attempted recrystallization (5a).	24
Figure 6: ^1H -NMR spectrum of 5d	25
Figure 7: Zoom in on the aromatic region in the ^1H -NMR spectrum of 5d . Coupling constants are displayed.	25
Figure 8: ^1H -NMR spectrum of 4e , boxed in regions depict characteristic signals for 4 (red = alkyl, green = aromatic).....	28
Figure 9: HMBC-NMR spectrum of 4e . (blue = aromatic interactions, green = carbonyl interactions)	29
Figure 10: ^{13}C - and ^{19}F -NMR spectrum of 4j	30
Figure 11: Crude ^1H - and ^{13}C -NMR spectra of the mixture of 4l and 1a ²	32
Figure 12: ^1H - and ^{13}C -NMR spectra for 4l . The enamine and keto forms are present in a 10:1 ratio.	33
Figure 13: <u>A</u> - 6a after attempted recrystallization, <u>B</u> - before work-up of 5a to 6a	34
Figure 14: An example of one of the repeated attempts to synthesize 6d by Tohda's method. (^1H -NMR spectrum)	35
Figure 15: The ^1H -NMR spectrum for 6b	37
Figure 16: ^1H -NMR spectrum of 7a (paramagnetic).	40

Figure 17: An example of a UV-Vis spectrum of copper complexes (7e).	41
Figure 18: ORTEP drawings of the unit cell of 7g and 7h (x-ray crystallographic representation)	42
Figure 19: Possible polymorphic behaviour in 7a/b	45
Figure 20: Both brown and green powders crystalline forms for 7b	46
Figure 21: Voltammogram of 7h/i displaying irreversibility and ligand reduction.	46
Figure 22: UV-Vis profiles for 8a showing three main absorption maxima.....	48
Figure 23: ¹ H-NMR spectrum of 9a (paramagnetic).	51
Figure 24: UV-Vis trace of 6e	52
Figure 25: ORTEP representations of 9j-l	52
Figure 26: Voltammograms for 9e and 9h in DCM.	55
Figure 27: A qualitative test of Trial 1 (9a + Br ₂)	57
Figure 28: ¹ H-NMR spectrum of proposed 9h-1	59
Figure 29: <u>A</u> - 10a and 10c as a powder. <u>B</u> - RIGHT: 10c (orange) during synthesis.	60
Figure 30: UV-Vis trace for 10b	61
Figure 31: Cyclic voltammetry for 10a and 10c	62
Figure 32: Solution darkening and oxidation of Co ²⁺ to Co ³⁺ (10a to 11a)	63
Figure 33: ORTEP representations of 11a and 11c	65
Figure 34: Voltammogram of 11a and 11c in DCM.	67
Figure 35: ³¹ P-NMR spectra of reaction between Wilkinson's catalyst and 4a	71
Figure 36: Target group 9 complexes for substitution.	72
Figure 37: Substitution of 4a to either 12 (left) or 13 (right).	73
Figure 38: ¹ H-NMR spectrum of 15	75

Figure 39: ^1H -NMR stacked of both R_f spots on the TLC plate of 15	76
Figure 40: ^1H -NMR stacked of both R_f spots on the TLC plate of 15	77
Figure 41: ORTEP representation of 15 (left) and cubic crystals of 15 (right)	78
Figure 42: ^1H -NMR spectrum of 16	81
Figure 43: ^1H - ^1H nOesy NMR spectrum of 16 . Marked is the CH_3 (15) interactions with the oxazoline protons (1/1a).	82
Figure 44: ORTEP representation of 16	82
Figure 45: <u>A</u> - ^{119}Sn -NMR spectrum of 18 , <u>B</u> - ^1H -NMR spectrum of 18	85
Figure 46: nOesy NMR spectrum of 18 (red box highlights NOE interactions)	87
Figure 47: ORTEP representation of 18	87

List of Schemes:

Scheme 1: 2-, 3-, and 4- oxazolines (left to right).....	1
Scheme 2: Three known routes to 2-oxazolines.....	2
Scheme 3: Examples of mono-charged and neutral ligands	3
Scheme 4: Conformational and rotation isomers of cobalt complexes	3
Scheme 5: Examples of Mono-, bi-, and tridentate oxazoline transition metal complexes ..	4
Scheme 6: Types of N,O-donor pro-chelates.	5
Scheme 7: A general look at three of Tohda's procedures to make the target oxazoline ligand class.....	6
Scheme 8: Tohda's diacylation and deacylation procedure to make the target compound, D	6
Scheme 9: Tohda's procedure for synthesis of the target compound, E , <i>via</i> AlCl ₃	7
Scheme 10: Generation of the S,S-dithio ketene and synthesis of the target compound, G . 8	
Scheme 11: Reaction to make target compound, H , with cold pyridine.....	8
Scheme 12: Lithiation of the oxazoline and reaction with EtOAc to give the target compound, E	9
Scheme 13: Tautomers of this ligand class, along the N-C-C-C-O backbone.....	9
Scheme 14: A few relevant copper and cobalt complexes (I-L)	11
Scheme 15: Wilkinson's catalyst (M) and Vaska's complex (N).	12
Scheme 16: Carlton's and Love's chelated iridium complexes.	12
Scheme 17: General mechanism for Cu-mediated ATRP.	14
Scheme 18: Matyjaszewski's varying ligand design for copper ATRP catalysts.	15
Scheme 19: Relevant cobalt(II) ATRP catalysts	16

Scheme 20: A library of ligands leading towards the formation of κ^2 -N,O-donor complexes	17
Scheme 21: Complexation of T1-4/U1-2/D to a cobalt(II) metal center	17
Scheme 22: Select compounds/complexes synthesized during this thesis.....	18
Scheme 23: General reaction conditions for the formation of 3a-i . Compound 3i was supplied by J. Adjei.	20
Scheme 24: General procedure on how to synthesize diacylated compounds 5a-d	23
Scheme 25: Formation of the ligand series with reacted initially with 1a . Compound 4i was made by J. Adjei.	26
Scheme 26: General reaction of tri-halogenated species with 1a to give the target alkyl ligands.....	30
Scheme 27: The synthetic route of 4l	31
Scheme 28: Formation of the ligand series with reacted initially with 1b	33
Scheme 29: Attempted synthesis of 6d	35
Scheme 30: A general metal complexation reaction of 4 or 6 to either Cu or Co	38
Scheme 31: An alternate route to Cu/Co complexes from 3 or 5	39
Scheme 32: Towards copper complexes with the ligand series 4	39
Scheme 33: Synthesis of Cu complexes with 6	47
Scheme 34: Towards cobalt complexes with the ligand series 4	50
Scheme 35: Attempted bromination of 9h	58
Scheme 36: Synthesis of complexes 10a-d	59
Scheme 37: General route to polymerization of MMA.....	67
Scheme 38: Formation of PMMA, based on best trial from Table 11	68

Scheme 39: Proposed mechanism for the formation of PMMA <i>via</i> 9h	69
Scheme 40: Attempted substitution of Wilkinson's Catalyst with 4a	70
Scheme 41: Synthetic route to make 15	77
Scheme 42: Oxidative addition of MeI to 15 to give 16	79
Scheme 43: Attempted oxidative addition to 15 to make 17	84
Scheme 44: Synthesis and oxidative addition of HSnPh ₃ to 15 to make 18	85
Scheme 45: Various attempted displacement reactions with P/N nucleophiles with 15	89

List of Equations:

Equation 1: Polydispersity Index.	13
Equation 2: Calculating the geometry index of 7g and 7h	43

List of Tables:

Table 1: Relevant bond angles (°) and bond lengths (Å) for 7g	44
Table 2: Select bond angles (°) and bond lengths (Å) for 7h	44
Table 3: Calculated τ^4 values for all Co complexes	53
Table 4: Selected bond lengths (Å) and angles (°) for 9j	54
Table 5: Selected bond lengths (Å) and angles (°) for 9k	54
Table 6: Selected bond lengths (Å) and angles (°) for 9l	54
Table 7: Attempted oxidation of a Co ²⁺ center.	56
Table 8: Selected bond lengths (Å) and angles (°) for 10a	62
Table 9: Select bond angles (°) and lengths (Å) for 11a	66
Table 10: Select bond angles (°) and lengths (Å) for 11c	66
Table 11: Trials and data for the polymerization of MMA	68
Table 12: Relevant bond lengths (Å) and angles (°) for 15	78
Table 13: Select bond lengths (Å) and angles (°) for 16	83
Table 14: Select bond lengths (Å) and angles (°) for 18	88
Table 15: Trials for oxidative addition of halogens to 9a	145

List of Appendices

Appendix A: Nuclear Magnetic Resonance Spectroscopy (NMR)	153
Appendix B: Infrared Spectroscopy (IR).....	250
Appendix C: Ultra-Violet-Visible Spectroscopy (UV-Vis)	285
Appendix D: Cyclic Voltammetry (CV).....	315
Appendix E: Gel-Permeation Chromatography (GPC)	343
Appendix F: Single Crystal X-Ray Diffraction (X-Ray Crystallography)	345

1.0 Introduction

1.1 Oxazolines

Oxazolines, or 4,5-dihydro-2-oxazoles, are five-membered heterocycles containing a nitrogen and oxygen between an sp^2 -hybridized carbon atom. They can differ sterically by substitution appended to any of its five vertices (except the oxygen atom) or isomerically by positioning of the π -bond (**Scheme 1**).

Scheme 1: 2-, 3-, and 4- oxazolines (left to right)

Oxazolines can be manufactured in a variety of methods including the Witte-Seeliger reaction (metal-catalyzed cross-coupling),¹ a ring-closing reaction with thionyl chloride,² from oxidation of an oxazolinidine or by simply heating a carboxylic acid or aldehyde with a 1,2-amino alcohol (**Scheme 2**).³ Oxazolines have gained synthetic interest due to not only their relatively stable and inert nature but also their close relation to acyl derivatives of α -amino acids.⁴ They are unreactive with most nucleophiles, radical species, bases and weak acids. Oxazolines were first used in the field of polymer chemistry as a monomer; the resulting poly(2-oxazolines)⁵ are like that of similar to poly(amides) making them strong candidates in biomedical applications.⁶ The aspect of this that is attractive to organic chemists is that the oxazoline protecting group can be installed during synthesis by using mild conditions and are moderately good precursors for functional later group transformation.^{4,6,7}

Scheme 2: Three known routes to 2-oxazolines.

More recently, oxazolines have been widely studied for their role in polymerization and protecting group chemistry^{8,9} as well as their ability to be a respectable monodentate ligands in asymmetric and symmetric catalysis.^{10,11} Over the years, oxazolines have had, and continue to exhibit, a vast array of applications from polymer chemistry to organic chemistry to coordination chemistry.⁴⁻⁷

1.2 Coordination Chemistry

Coordination chemistry entails the formation of covalent/coordination bonds (usually dative interactions) between, normally, a metal ion and neutral or charged ligands. In the case that the resulting complex carries a charge, an ion with opposite charge is present to balance it to a neutral state. As previously mentioned, ligands can be either neutral or charged (cationic or anionic). Neutral ligands include *N*-heterocyclic carbenes, pyridine, acetonitrile and water (**Scheme 3**).

Scheme 3: Examples of mono-charged and neutral ligands

Most charged ligands are anionic in nature as transition metal centers are generally cationic. Charged ligands can be mono-, di- or tri- anionic such as halogen species (Cl^- , Br^- , F^- , I^-) and deprotonated oxygen species (e.g., 8-hydroxyquinolinolate) (**Scheme 3**). Once coordinated, octahedral ligated metal complexes can exhibit different geometric isomers; *mer/fac*, *cis/trans*, as well as lambda/delta (Λ/Δ) rotational isomers (**Scheme 4**).

Scheme 4: Conformational and rotation isomers of cobalt complexes

1.2.1 Denticity

The number of donor groups from a specific ligand attached to a central metal atom denotes its denticity.¹² Oxazolines on their own can act as monodentate ligands^{13–16} or polydentate ligands when functionalized with other groups, meaning they can be bound to the central atom through one or more bonds.^{17–19} There are multiple examples of oxazoline-based ligands with varying

denticity in the literature including mono- (**A**)¹⁴, bi- (**B**)²⁰, and tridentate (**C**)¹⁹ architectures (**Scheme 5**).

Scheme 5: Examples of Mono-, bi-, and tridentate oxazoline transition metal complexes

The examples given in **Scheme 5** represent transition metal complexes containing chelating ligands. Strong donating interactions, such as in the case of the nitrogen in the oxazoline ring, readily form bonds with a formally cationic metal center. If the oxazoline ring is appended with another anionic functional group, the compound can form a chelate bond with the metal center effectively changing its bond mode from κ^1 to κ^2 (or higher).

1.3. κ^2 -N,O Chelating Ligands

There are several types of nitrogen-based bidentate chelating ligands such as those with NN, NS, NP, NC, or NO-donor atom functionality. N,O chelates have been targeted as a class of ligands in the fields of catalytic and coordination chemistry due to their variable bonding modes and binding affinities for cationic metal species.²¹ After being bound to a metal center the participating ligands are employed in applications ranging from small molecule activation to stoichiometric bond activation and catalysis. In an N,O donor ligand, the oxygen atom is often anionic in character, making it a hard (Lewis basic) donor to the metal center.²² The nitrogen is an intermediate donor, meaning it can exhibit properties of both a hard and soft donor. This gives the nitrogen atom freedom to bond then release itself from the metal. Overall, this hemilabile quality makes N,O donor ligands attractive for use in catalysis when coordinated to a metal center.

1.3.1 Classes of N,O Chelating Ligands

A lot of focus in the literature has been put on N,O chelates, such as those derived from amino acids.^{23,24} More common to the field of inorganic coordination chemistry are the 1,3- and 1,4-N,O chelates. These functional groups (**Scheme 6**), most noticeably quinolinolates^{25–27} and picolinates,^{28–30} have seen a span of uses catalytically. Sulfonamides³¹ and phosphoramidates^{32,33} have started to see more use in the field as well.

Scheme 6: Types of N,O-donor pro-chelates.

1.3.2 Synthesis of Ligands

A target of the Gossage groups' research has been centered around a 1,5-N,O pro-ligand, first synthesized by Meyers³⁴ and later popularized by the simple synthesis of the series from Prof. Tohda in 1984.³⁵ In his studies of these compounds, Tohda crafted clever, high yielding synthetic pathways to these compounds.^{36,37} He later went on to continue his investigation of the formation of abnormal Michael adducts that these particular oxazoline based compounds can adopt.⁸ He devised three simple synthetic pathways (**Scheme 7**); 2-step acylation and deacylation, a crossed Claisen condensation with AlCl₃ coordination (for acid chloride compounds with α -protons to the carbonyl), and a reaction that utilizes pyridine to synthesize compounds with more electron

withdrawing donor groups. Another, less popular method to make these types of compounds comes from the use of a dithio-ketene^{9,38,39} (*vide infra*).

Scheme 7: A general look at three of Tohda's procedures to make the target oxazoline ligand class

1.3.2.1 Ligands via Diacylation and Hydrolysis

The first, and most versatile, synthetic pathway developed by Tohda is a two-step reaction.³⁵ The starting oxazoline is acylated at the *N*-position first, which then activates the methyl group on the oxazoline at position 2 (**Scheme 8**). This is done in a polar aprotic solvent as the transition state of the intermediate is very polar. The isolated intermediate material, in high yields (82-99 %), is subjected to a basic mixture of excess potassium hydroxide dissolved in methanol (1.5 M KOH). This cleaves the C-N bond at the *N*-acetylated position to give the desired compound in moderate to high yields (57-95 %). This reaction has been optimized to the point where one can produce > 30 grams of **D** without sacrificing yield or materials.³⁵

Scheme 8: Tohda's diacylation and deacylation procedure to make the target compound, **D**

1.3.2.2 Ligands via Crossed-Claisen Reaction

The second pathway developed by Tohda in 1986 was a way to synthesize these oxazoline derived compounds if an alkyl-carbonyl substituent bonded to the heterocycle. The 2,4,4-trimethyl-2-oxazoline and triethylamine are added to a solution of aluminum chloride.³⁶ After which an anhydride species is added over 30 minutes at 0 °C. Once obtained, but not isolated, the intermediates species is immediately dissolved in a basic potassium hydroxide/methanol solution to cleave the C-N bond on the 3-position of the oxazoline (**Scheme 9**). The pure product (**E**) is obtained by *via* co-distillation with ethylene glycol. Although this method is effective for synthesizing these types of compounds with alkyl chains that have protons directly next to the carbonyl, yields do not exceed 75 %.³⁶

Scheme 9: Tohda's procedure for synthesis of the target compound, **E**, *via* AlCl₃

1.3.2.3 Ligands via *S,S*-dithio-Ketenes

A less popular route used in the synthesis of these oxazoline-appended compounds is the reaction of a *S,S*-dithio-ketene with an amino alcohol. The dithio ketene is generated *in situ* in a one-pot reaction between NaH, carbon disulphide (CS₂) and methyl iodide (MeI) (**Scheme 10**).^{9,38,39} In this reaction, a sample of a benzyl ketone is irreversibly deprotonated and reacted with CS₂ and MeI to give the dithio intermediate (**F**). The intermediate compound is immediately dropped into a dispersion of Na and the chosen amino alcohol. After reflux, the target ring-closed compound is isolated as an off-white powder (**G**).⁴⁰ Drawbacks of this reaction pathway is the overall technical difficulty of the synthesis and lower yields obtained of the targets (49-63 %).

Scheme 10: Generation of the S,S-dithio ketene and synthesis of the target compound, **G**

1.3.2.4 Ligands via alternative methods – pyridine/lithiation

Two last methods are more uncommon as they can make a smaller range of such modified oxazoline compounds. The first method is by using a mixture of pyridine in acetonitrile with a strongly electron withdrawing anhydride. In this route, the reaction is kept at a temperature of 0 °C to prevent unwanted by-products and is mediated by pyridine (**Scheme 11**). After completion, the target compound can be isolated as an off-white powder in excellent yields (90-93 %).⁸

Scheme 11: Reaction to make target compound, **H**, with cold pyridine.

The second method is by using *n*-BuLi in an electrophilic substitution of a ketone species to obtain the target compound (**Scheme 12**). The starting 2-oxazoline compound is subjected to *n*-butyllithium. This lithiates the 2-position of the oxazoline, forming the supposed lithio-oxazoline intermediate. Afterwards the lithio-oxazoline is reacted with the ester of choice and can be isolated as an off-white powder (**E**) after extraction and work-up (85 %).³⁴

Scheme 12: Lithiation of the oxazoline and reaction with EtOAc to give the target compound, **E**.

1.3.3 Tautomers of the Target Ligand Series

This class of compounds display at least three different tautomers (resonance contributors); the enol, keto and enamine forms (**Scheme 13**). Although, in the solid state of the compound series, the enamine form is most prominently observed (**Figure 1**) and in the solution state a mix of the enamine and keto form can be observed.³⁵

Scheme 13: Tautomers of this ligand class, along the N-C-C-C-O backbone.

Figure 1: Crystal structure of **D**, seen in the enamine form.

1.4 Relevant Transition Metal N,O-complexes

1.4.1 Copper(II)/Cobalt(II/III) Complexes

The area of copper(II) and cobalt(II) complexes bearing N,O-donor ligands has been extensively researched with literature searches returning complexes bound to all classes of ligands, including a wide variety of amino acid functionalized complexes. (**Scheme 14, I**).^{41–43} Other N,O-donor ligands that are commonly found bound to metals are those of the amidate, phosphoramidate, quinolinolate and the pyridonate classes (**Scheme 6**). Cobalt and copper have received increasing attention over the years as substitutes for heavier transition metals due to their cheaper cost and facile synthesis.^{44,45} Tendenborg in 1999 synthesized a small library of ligands (both N,O- and N,N-chelates) and coordinated them to both copper and cobalt metal salts (**Scheme 14, J**).⁴⁶ They later tested those complexes in their ability to catalyze the asymmetric aziridination of styrene. They received high yields with modest enantiomeric excesses in short reaction times (10 minutes). In 2016, preliminary studies on copper(II) coordination polymers were disclosed by the Amo-Ochoa group and their pyridonate N,O-ligands showed their exceptional binding utility by attracting other copper complexes to form a polymeric substance (**Scheme 14, K**).⁴⁷ More recently, in 2017 Shakhdofa and coworkers presented their work on copper and cobalt anti-tumor complexes. The tested complexes [NNO, NN, OO-(dinuclear Cu species) and NO ligated complexes] showed a modest level of activity against the human leukemia, human liver cancer, and human breast cancer cell lines although not as active as a current anti-tumor drug, Doxorubicin (**Scheme 14, L**).⁴⁸

Scheme 14: A few relevant copper and cobalt complexes (I-L)

1.4.2 Rhodium(I/III)/Iridium(I/III) Complexes

Further down Group 9 lies Iridium and Rhodium. In recent years (1960-present), there have been a plethora of studies surrounding their interesting properties. Rhodium, and Iridium to a degree, were initially used as hydrogenation catalysts (addition of H_2 across a π -bond)^{49–52} as is in the case of Wilkinson's catalyst (**Scheme 15, M**).⁵³

Both Rhodium and Iridium have gained significant interest in the research field as they can efficiently facilitate oxidative addition/reductive elimination-type reactions (redox). This lead to their increasingly popularity in the field of catalytic-coordination chemistry. One of the best candidates for studying hydrogen transfer, as well as other general oxidative addition reactions is Vaska's complex, $\text{IrCl}(\text{CO})(\text{PPh}_3)_2$ (**Scheme 15, N**). These reactions can be visualized physically as well they tend to lose their bright yellow colour and through infrared spectroscopy as they observed to have higher stretching frequencies ($> 2000 \text{ cm}^{-1}$ once oxidative addition occurs at the **Ir** center).

Scheme 15: Wilkinson's catalyst (**M**) and Vaska's complex (**N**).

Two coordination compounds to note are **O**⁵⁴ and **P**.^{32,33,55} Using a simple two-step method (use of an iridium-COD dimer⁵⁶ then coordination of their ligand) then by the addition of a reaction nucleophilic species, the Carlton group has been able to synthesise an array of H-Ir-R complexes (see **Scheme 16, O**). This reactions series showcases the redox ability of the iridium centered complex as it oxidizes from an Ir(I) to Ir(III).⁵⁴ In a more recent example, the Love and Schafer groups have been able to synthesize 1,3-N,O-chelated phosphoramidate complexes (**Scheme 16, P**). These complexes have shown to perform a variety of oxidative addition/reductive elimination reactions with various boron substrates. In doing so, they've demonstrated the versatility of these types of complexes by achieving a high yielding, selective hydroboration.^{32,33,55}

Scheme 16: Carlton's and Love's chelated iridium complexes.

1.5 Polymers

Polymers are macromolecular structures or more simply explained as long chains with repeating units. There are two main different types of polymers, organic (PANi, PVC, PS) and inorganic polymers (polyphosphazenes, polysilanes, poly(ferrocenylsilanes)). These

macromolecules can consist of one singular repeat unit (homopolymer) to many different repeat units (co-polymer) arranged neatly (block co-polymer) or unpredictably (random co-polymer). Polymers are characterized the degree of their uniformity (**Equation 1**). This measurement is known as the polydispersity index (PDI).⁵⁷

$$\text{PDI} = \frac{M_w}{M_n}$$

← Weight-average molar mass
← Number-average molar mass

Equation 1: Polydispersity Index.

1.5.1 Mechanisms of Polymer Growth

The two generally accepted mechanisms surrounding polymer growth are step-growth (condensation) and chain growth (addition). Step-growth polymerization occurs by the monomer units adding to each other in a more unpredictable fashion, having monomers reacting to make dimers, dimers reacting to form tetramers, or a monomer unit and a dimer reacting to form a trimer. These seemingly random chain length oligomers react with each other to increase the polymer chain length. A polymer polymerized with this mechanism generally produces low molecular weights, high percent conversion but also a more broad PDI.⁵⁸

The other main accepted mechanism is by chain growth. With a chain growth mechanism, the reactive species, usually a radial species, initiates and facilitates the lengthening of the polymer chain. In this case a new monomer is added tip-to-tail or step by step to the polymer chain. High polymer molecular weights, along with a narrow PDI but low conversion rate are consistent with this mechanism.

1.5.2 Metal Catalyzed Atom Transfer Radical Polymerization (ATRP)

Atom transfer radical polymerization (ATRP) was simultaneously discovered by Mitsuo Sawamoto,⁵⁹ Jin-Shan Wang and Krzysztof Matyjaszewski in 1995.⁶⁰ ATRP starts off with the initiation of the catalytic species (see **Scheme 17**). The catalyst chosen for metal-based ATRP must be able to have a reasonable affinity for halogens, be able to accommodate the extra halogen in its coordination sphere as well as have an easily accessible oxidation state with typically a one electron difference.⁶¹

Scheme 17: General mechanism for Cu-mediated ATRP.

Afterwards the initiated radical species ($\text{R}\cdot$) seeks out the olefinic monomer. Once found, the radical species bonds to the olefin, propagating along the backbone of the molecule ($\text{P}\cdot$).⁶¹ This step occurs continuously and stops once all of the monomer units have been bonded together. The polymer is terminated by the halogen species that was originally bonded to the initiator as the catalyst “terminates” the initiator and the polymerization is ended. Additionally, ATRP is a living polymerization, meaning that the process can be stopped and restarted. This process is ideal for co-block polymerizations as one block can be polymerized, stopped, and a second monomer can be polymerized onto the first block. This facilitates an effective and controlled synthesis.^{60,61}

Another type of radical polymerization is nitroxide mediated polymerization (NMP). This method is akin to ATRP although makes no use of metal species and is usually initiated and/or mediated by TEMPO, or other radical nitroxide functional molecules (nitroxide = $\text{N-O}\cdot$).

1.5.2.1 Copper(I/II) Catalysts in ATRP

ATRP, when it was first discovered, primarily used copper(I) catalysts as they could accommodate the halogen initiative species in their coordination sphere, lead to minimal side reactions and its higher oxidation state (Cu^{2+}) can be easily accessed. The first copper complex to be used in this manner by Matyjaszewski is the *in situ* generated catalyst, copper(I)bipyridine chloride.⁶⁰ Their catalyst worked efficiently, giving PDI's < 1.5 (narrow molecular weight distributions). They noted that without the additive (bpy) the molecular weight and PDI were ill controlled and gave poor results.⁶⁰ Matyjaszewski and co-workers have been building to expand this field with the use of their copper catalysts by varying the coordinated ligand.⁶² Different ligand scaffolds they've recently used (see, **Scheme 18**) contain four N-donor atoms, are potentially tetradentate ligands. Currently, their pre-catalyst $[\text{Cu}(\text{TPMA}^{\text{NMe}_2})\text{Br}][\text{Br}]$ exhibits the highest $k_{\text{ATRP}} (= 1)$, meaning the rate at which ATRP can be started and the rate at which it can be stopped are almost equal. A good ATRP catalyst has a k_{ATRP} value of 1, meaning the ease of which polymerization can be started and stopped and started again is high, making an effective catalyst for co-block polymerizations. This means for a true reversible, radical polymerization.⁶²

Scheme 18: Matyjaszewski's varying ligand design for copper ATRP catalysts.

1.5.2.2 Cobalt(II/III) Catalysts in ATRP

Since the discovery of ATRP, chemists have set out to find other metal containing-complexes that can perform ATRP more efficiently than its copper predecessors. Simple systems such as **Q** and **R** (Scheme 19) as well as more complex systems such as **S** have been recently explored.

Scheme 19: Relevant cobalt(II) ATRP catalysts

Matyjaszewski and co-workers have also tested a simple cobalt catalyst (**R**) analogue in the polymerization of MMA and styrene.⁶³ This catalyst was specifically tested in the copolymerization of two different acrylate species, ultimately terminated by TEMPO. Here they observed narrow PDI's (< 1.2) at very high conversion rates with high molecular weights (~ 40 000 Da).⁶³ The Zhang group have shown with their cobalt catalyst (**Q**) they can achieve very high molecular weight with high conversion rates with narrow PDI's (< 1.35). What is interesting about their synthesis is that they've used RAFT agents to neatly facilitate polymerization *via* an ATRP mechanism.⁶⁴ More recently, the Yan group have showcased the catalytic ability of the cobalt enriched polymer. With this macromolecule, they could polymerize a variety of acrylates (PMMA, 40 % conversion, PDI = 1.84) as well as styrene (PS, 42 % conversion, PDI = 1.92) in modest yields and narrow dispersities.⁶⁵

1.6 Past Work

Previous work in the Gossage group on this project pertains to synthesis of the target ligand series as well as preliminary coordination work with copper²⁰ and cobalt.⁶⁶ Pertaining to ligand synthesis either Tohda's, or Meyer's procedures (**Scheme 8**, **Scheme 9**) were used to synthesize **T1-4**, **D**, **E** (**Scheme 20**).

Scheme 20: A library of ligands leading towards the formation of κ^2 -N,O-donor complexes

Compound **D** was selected as a candidate to test the coordination abilities of the ligand series. The compound was ligated to a copper metal center and is observed to be the deprotonated enolate form of the ligand. This work was repeated with a cobalt metal salt to receive **T1-3**, **U1/2**, **D** (**Scheme 21**).⁶⁶ The reaction, although low yielding (in the case of **Co/Cu**), worked well enough to isolate single crystals of each of the synthesized species.

Scheme 21: Complexation of **T1-4/U1-2/D** to a cobalt(II) metal center

1.7 Objective of This Thesis

The purpose of this thesis is to synthesize, characterize and expand the already established library of κ^2 -N,O-bidentate oxazoline ligands. Varying the -R substituents as well as the substituents on the 4-position of the oxazoline (-R') will influence the electronic properties of the ligand therefore varying the redox properties of the metal complex. The series of κ^2 -N,O-bidentate ligands will be coordinated to either cobalt, copper or iridium, to obtain novel Group 9 and 11 complexes (**Scheme 22**). All organic compounds will have their purity tested by a series of NMR techniques and melting point. All transition metal complexes will be characterized by UV-Vis, IR, CV, X-ray diffraction, MS, melting point, and a combination of NMR spectroscopy experiments. The complex series will be trialed as polymerization catalysts with monomers such as methyl methacrylate. Oxidation of an Iridium(I) centered complex giving two novel Iridium(III) complexes will be explored and characterized by IR, a combination of NMR spectroscopic experiments and X-ray diffraction where applicable. Synthesis of Rhodium complexes will also be attempted.

Scheme 22: Selected compounds/complexes synthesized during this thesis.

2.0 Results and Discussion

2.1 General Considerations

Each of the following reactions (**Sec. 2.2-2.4**) were completed in either the exact method or an adaption of the methods first used by Tohda *et al.*,^{8,35} Pittman and coworkers,⁹ or Meyers and colleagues³⁴ and are referenced appropriately to their article or origin. The metal complexation reactions were modified from a preparation by Gossage *et al.* to further optimize their synthesis.²⁰ The ligands can be obtained in one to two steps and the metal complexes can be obtained in an additional one to two step(s). Samples of compounds **3i** and **4i** were obtained from J. Adjei (Ryerson University), synthesized by the same methods as described below.⁶⁷

2.2 Synthesis and Characterization of Intermediate Compounds (3a-i, 5a-d)

The diacylated intermediate compounds (**3a-i**) were synthesized using commercially available 2,4,4-trimethyl-2-oxazoline (**1a**) and the corresponding acid chloride. In each case, 1 equivalent of **1a** was heated to reflux with 2.2 equivalents of the acid chloride (**2a-i**) and 3 equivalents of NEt₃ in acetonitrile for three hours (**Scheme 23**). After which, the mixture is filtered to exclude the triethylammonium chloride salt formed. Once all volatile components have been removed, the mixture is subjected to two washes; the first with equal parts chloroform and distilled water, and the second wash with equal parts chloroform (previous wash) and a 5 % solution of sodium carbonate in water (Na₂CO_{3(aq)}). The washes extract excess triethylamine (and [NEt₃][HCl]) as well as by products such as benzoic acid derivatives. All impurities are assumingly removed in the washing process. The organic phases were dried (MgSO₄ or Na₂SO₄), filtered, then all remaining volatile components were removed under vacuum. Once removed, the crude product was revealed as a yellow-orange sludge. The crude mixture is purified by recrystallizing with a

1:1 toluene/*n*-hexanes combination, except for **3d** which was done with a 1:2 mixture of toluene/*n*-hexanes. Depending in on the “*R*” substituent, the pure product emerged was either a light orange or a pale yellow solid. Yields ranged accordingly to the steric bulk associated with the “*R*” group itself with seemingly little to no electronic influence of the substituents in the variable group. The obtained yields were moderate to excellent for the intermediate species (45-99 %)

Scheme 23: General reaction conditions for the formation of **3a-i**. Compound **3i** was supplied by J. Adjei.

All compounds are novel, with the exception of **3a/c/h/i**^{20,35,67} which are known in the literature. For all the synthesized intermediate species, characterization was carried out by melting point (mp), NMR, TLC and in the case of the novel compounds by elemental analysis (EA). For these intermediate compounds, the ^1H -NMR spectrum generally displays eight to eleven different proton environments for the aryl derivatives (**3a-g/i**) and five different proton environments for the alkyl derivative (**3h**).

Figure 2: One of the intermediate compounds, **3e**, labeled for aid in NMR discussion.

Figure 3: ^1H -NMR spectrum of **3e**.

Resonance signals indicative of the desired product are those of positions **1**, **3**, **5** and the overall ‘doubling’ of aryl resonance signals (**Figure 2**). These protons (**1**, **3**, **5**) resonate at approximately $\delta_{\text{H}} = 1.55$, 4.25 , 5.00 ppm depending on the “*R*” group attached to the carbonyl group at position **6**. Specifically, **3e** displays resonance signal at $\delta_{\text{H}} = 1.62$, 4.33 and 4.95 ppm (**Figure 3**).

For this compound (**3e**), there are six distinct aromatic signals which resonate between 7.0 - 8.0 ppm. Of these signals, two display doublet character due to their four-bond coupling (^4J) with either the protons on the neighboring carbon’s methyl group or from the other *ortho* carbon’s

proton. The other four signals show a characteristic doublet of doublets splitting pattern. This splitting pattern is indicative of a proton interacting with an immediate neighboring proton (^3J) and with second order effects from a further proton ($^{4/5}J$).

Figure 4: ^{13}C -NMR spectrum for **3e**.

In the carbon NMR (^{13}C -NMR) spectra, there are generally fifteen to twenty-three different carbon environments for the aryl species (**3a-g**) and eleven different carbon environments for the alkyl species (**3h**). The frequency of which the carbons at positions **1**, **3**, and **5** (**Figure 2**) resonate are the first sign of the successful synthesis of the target intermediate species. In general, these positions resonate at approximately $\delta_{\text{C}} = 25$, 80 and 100 ppm for **1**, **5**, and **3** respectively. Like the proton NMR spectrum, these signals vary greatly depending on the “*R*” group attached to the enolic carbonyl. For this compound, **3e**, the carbons **1**, **5** and **3** resonate at $\delta_{\text{C}} = 22.8$, 79.8, 85.4 ppm respectively (**Figure 4**). By observing the coupling constants and a variety of two-dimensional

NMR experiments (COSY ^1H - ^1H , HSQC ^1H - ^{13}C , HMBC ^1H - ^{13}C) assignments have been made to each of the compounds.

The second variety of diacylated compounds (**5a-d**) were made in a similar manner where the only starting oxazoline differed to 2-methyl-2-oxazoline (**1b**). In each of the four cases, 1 equivalent of **1b** was refluxed with 2.2 equivalents of the corresponding acid chloride and 3 equivalents of NEt_3 in tetrahydrofuran for 3 hours (**Scheme 24**). This process differs from the synthesis of **3a-3g** as described by Pittman and coworkers as the reaction leads to a ring-opened chlorinated intermediate, **5**, as displayed.⁹

Scheme 24: General procedure on how to synthesize diacylated compounds **5a-d**.

After completion of the reaction, the salt is filtered from the solution where then the volatile components can be removed. The sludge is afterwards subjected to two washes, once with equal part chloroform and water and the second wash with equal parts chloroform and a 5 % solution of $\text{Na}_2\text{CO}_3(\text{aq})$. All resulting impurities were removed during the washing process. The organic phases were dried (MgSO_4 or Na_2SO_4), filtered, then all remaining volatile components were removed. Once removed, the crude product was revealed as a yellow-orange sludge. The crude mixture was attempted purification by recrystallizing with a 1:1 mixture of toluene/*n*-hexanes, as in the

methylated oxazoline compounds (**3**). All attempts in this matter lead to the mixture of compounds oiling out of solution together instead of solidifying and recrystallizing separately (**Figure 5**).

Figure 5: A- Prior to work-up, before filtering the salt ($[\text{NEt}_3][\text{HCl}]$), B- After work-up and attempted recrystallization (**5a**).

Thus, the intermediate species (**5a-d**) were attempted to be purified by column chromatography, although all attempts were met with failure as there were various impurities mixed together with similar retention factors in the case of **5d**. It is known as revealed by Pittman and co-workers⁹, that 2-methyl-2-oxazoline undergoes a ring opening mechanism when reacted under the same conditions as described above.³⁵ This, most likely, gave the intermediate species as a ring open oxazoline (**5**) instead of the target ring closed oxazoline species (**3**). Although by careful column chromatography a sample of **5d** was obtained. Compound **5d** recrystallized from 45% EtOAc/*n*-hexanes and was obtained as a fully off-white powder. This sample was characterized only by ¹H-NMR spectroscopy. In the ¹H-NMR spectrum signals corresponding to the proposed compound, **5d**, have been observed (**Figure 6**).

Figure 6: ^1H -NMR spectrum of **5d**.

Figure 7: Zoom in on the aromatic region in the ^1H -NMR spectrum of **5d**. Coupling constants are displayed.

When looking at the coupling constants of the aromatic region, there are four resonance signals which coupling to other one other aromatic resonance signal. Their corresponding ^3J values correlate to one other signal, which is indicative of a *para* substituted phenyl ring (**Figure 7**).

The use of these intermediate compounds will be detailed in [Section 2.3](#) (compounds **3a-i**) and [Section 2.4](#) (compounds **5a-d**).

2.3 Synthesis and Characterization of a Ligand Series with **1a** (**4a-i** and **4j-l**)

2.3.1 Synthesis and Characterization of ligands **4a-i**

Once the intermediate compounds have been obtained, they are subjected to a 1.5 M methanolic potassium hydroxide solution (KOH dissolved in MeOH).

Scheme 25: Formation of the ligand series with reacted initially with **1a**. Compound **4i** was made by J. Adjei.

These basic conditions cleave the C-N bonded acyl group. Typically, the reagents are stirred at room temperature for six hours. Once the reaction time elapsed, all volatiles are removed from the flask and are subjected to two washes with distilled water. During the second distilled water wash, sometimes a scoopula tip of brine is added to help separate the layers. The organic phase is dried

(MgSO₄), filtered and again volatile components are removed. The resulting pale yellow or light orange sludge is recrystallized with a 1:4 mixture of toluene/*n*-hexanes. This reveals the target ligand as either pale yellow needles (**4d-i**) or white needles (**4a-c**) in moderate to high yields (23-99 %).

All compounds are novel, with the exception of **4a/c/h**^{20,35} which are known in the literature and **4b/g/i**⁶⁷ which have been previously synthesized by our group.⁶⁶ For all the synthesized ligand species, characterization was carried out using their melting point (mp), NMR spectroscopy (a combination of various 1D/2D experiments), TLC and in the case of the novel compounds by elemental analysis (EA). The ligand series exhibits characteristic resonances in both the proton and carbon NMR spectra. Such as the case of **4e** series (**Figure 8**), the proton NMR spectrum displays all the characteristic resonance signals for the target ligand series.

Resonance signals for the two-methyl groups (~1.43 ppm) and the CH₂ (~4.15 ppm) on the heterocycle as well as the enolic resonance (~5.50 ppm) support the formation of the ligand. For more evidence the aromatic region can be examined. In this case in the aromatic region there are three signals being split to observe a doublet (~7.23 ppm), a doublet of doublets (~7.62 ppm), and a singlet with distinguishable ⁴*J* coupling to another proton environment to give a doublet (~7.82 ppm).

9 and **13** positions. Meaning those carbons interacting with position **12** must be situated opposite to the oxazoline. With this information one can deduce the correct assignments of positions **8**, **10** and **11**. Position **8** lies next to the carbonyl as it interacts with protons **9** and **13**, position **11** is the chlorinated carbon in the *para*-position (this is supported by its very downfield shift), and position **10** (the only remaining assignment) is the methyl-carbon *meta* to the carbonyl.

Figure 9: HMBC-NMR spectrum of **4e**. (blue = aromatic interactions, green = carbonyl interactions)

2.3.2 Synthesis and Characterization of ligands **4j-l**

To make shorter alkyl variants of the target ligands series different reaction pathways were explored. The first is the reaction of **1a** with either trichloroacetyl chloride (TCAC) or with trifluoroacetic anhydride (TFAA). Once activated by the pyridine, the tri-halogenated species become very reactive and thus the reaction mixture is kept cold. This synthetic methodology has been described by Tohda *et al.*⁸ The reaction was done by cooling all reagents down in separate flasks then by adding the oxazoline and pyridine together in ACN. Once the tri-halogenated species was cooled it was added dropwise to the stirring solution of **1a**/py. The reaction was stirred at 0 °C for 6 hours. After the reaction was completed the volatile components were removed *in vacuo*. The resulting sludge was dissolved in distilled water and washed with DCM twice. The organic

phased was dried (MgSO₄), filtered and again the volatile components were removed. The sludge was dissolved in hot *n*-hexanes and left to recrystallize. This revealed the product (both **4j** and **4k**) as white powder. This reaction is done in a 1:2.2:2 ratio of **1a**:TCAC/TFAA:py (**Scheme 26**).

Scheme 26: General reaction of tri-halogenated species with **1a** to give the target alkyl ligands.

The alkyl ligands, **4j/k**, were analyzed by both ¹H- and ¹³C-NMR spectroscopy where all characteristic resonance signals were observed as in the literature.⁸ In addition, compound **4j** gave one resonance in the ¹⁹F-NMR spectrum at δ_F = -79.8 ppm as expected (**Figure 10**).⁸

Figure 10: ¹³C- and ¹⁹F-NMR spectrum of **4j**.

The second reaction pathway known to produce alkyl derivative of **4** is the lithiation of **1a** followed by nucleophilic substitution of ethyl acetate to form **4l** (**Scheme 27**).³⁴

Scheme 27: The synthetic route of **4l**.

This reaction was carried out by first chilling **1a** in ACN to $-84\text{ }^{\circ}\text{C}$, this is done by creating a cold bath by freezing ethyl acetate with liquid nitrogen. Once chilled, $n\text{-BuLi}$ is added dropwise with rapid stirring of the oxazoline solution. This would lithiate the methyl on the 2 position of the oxazoline (**Scheme 27**). This then was stirred, initially at $-84\text{ }^{\circ}\text{C}$ but was left to gradually warm to room temperature, overnight. Once completed, the lithio oxazoline species was cooled back down to $-84\text{ }^{\circ}\text{C}$ where cold, dried EtOAc ($4\text{\AA}$ sieves and MgSO_4 for 3d, $-26\text{ }^{\circ}\text{C}$) was added dropwise. This solution was left to stir for 2 hours, where the solution was warmed to room temperature, to let it fully react. Upon completion, distilled degassed water was added slowly to the solution to quench excess $n\text{-BuLi}$. The mixture was washed with diethyl ether twice, the organic phased dried (MgSO_4), filtered and had all the volatile components removed. This left behind a white sludge which was attempted to be recrystallized by washing with cold $n\text{-heptanes}$. The isolated material was analyzed by ^1H - and ^{13}C -NMR spectroscopy.

Figure 11: Crude ^1H - and ^{13}C -NMR spectra of the mixture of **4l** and **1a²**.

When analyzed by NMR spectroscopy, a mess of resonance signals sprouted in the alkyl region of the proton NMR spectrum and in the $\text{sp}^2\text{-C}$ /alkyl regions of the carbon NMR spectrum (**Figure 11**). This impurity, previously reported by Meyers and coworkers, is said to be a dimer (**Scheme 27**).³⁴ This oxazoline dimer, **1a²**, is most likely synthesized in the lithiation step; presumably if a lithio-oxazoline species finds another molecule of **1a** before it finds its target (EtOAc) and hence a dimer is formed. Compound **1a²** was not isolated from this reaction but was instead treated with warm distilled water (60 °C) for four hours while stirring.³⁴ The solution is washed again, with diethyl ether twice. The organic phase is dried (MgSO_4), filtered and had all the volatile components removed. This process cleaves the oxazolidine moiety and gives **4l**. When pure **4l** is analyzed by ^1H - and ^{13}C -NMR spectroscopy, both the enamine and the keto form of the ligand are observed in a 10:1 ratio (E:K) as noted by Meyers *et al.* (**Figure 12**).³⁴

Figure 12: ^1H - and ^{13}C -NMR spectra for **4l**. The enamine and keto forms are present in a 10:1 ratio.

2.4 Synthesis and Characterization of a Ligand Series with 2-methyl-2-oxazoline (**6a-d**)

All these compounds (**Scheme 28**) are known in the literature except **6b**. Compounds **6c/d** are known but have been synthesized by reacting a dithio ketene (made by reacting CS_2 , NaH with acetophenone [**Sec 1.3.2.3**]) with 2-amino-2-methyl-1-propanol. This way the target compounds are made in moderate yields (49-63 %).^{7,8} In this study, Pittman and co-workers' procedure⁹ was more closely followed for the synthesis of **6a**, whereas Tohda's procedure was followed in an attempted to synthesize **6b-d**.³⁵

Scheme 28: Formation of the ligand series with reacted initially with **1b**.

Just like prior routes to make **4a-i**, **6a-d** were synthesized by dissolving the crude mixture of **5**, previously noted in **Section 2.4**, in a 1.5 M solution of potassium hydroxide (KOH) dissolved

in methanol (MeOH) (**Scheme 28**). This basic mixture facilitates the cleavage of the C-O and then the C-N acyl groups as well as reforming the heterocyclic ring. Once the 18-hour reaction period is over, the solution is filtered so that any precipitated salts can be excluded. A 1.0 M solution of sulphuric acid in water ($\text{H}_2\text{SO}_{4(\text{aq})}$) is added dropwise with stirring to MeOH solution to obtain a pH neutral state. Once neutral, the solution is washed with distilled water twice. The organic phased is dried (MgSO_4), filtered and all volatile components are removed.

Figure 13: A- **6a** after attempted recrystallization, B- before work-up of **5a** to **6a**

The resulting dark red/brown sludge was attempted to be recrystallized with varying ratios of toluene/*n*-hexanes although each time the ligand ‘oiled’ out of solution instead of solidifying (**Figure 13**). Column chromatography eluting with 30 % acetone/*n*-hexanes as in the Pittman procedure, was then employed for purification. The target, **6a/c** (33-64 %), were isolated as pale-pink powders whereas **6b** was isolated as a pale-yellow powder (83 %). Unfortunately, all attempts to synthesize compound **6d** resulted in failure. Regardless of purity of **5d**, when subjected to the basic KOH/MeOH mixture the compound was not able to be isolated. The reaction was tried several ways, including varying reaction time, KOH concentration, and purifying the intermediate species.

Scheme 29: Attempted synthesis of **6d**.

Once the reaction was completed and volatiles removed, the sludge was washed twice with water, although each wash became progressively harder and harder to separate. The addition of several milliliters of brine (and eventually replacing water with brine) aided in separation. Regardless of good separation of the phases, when it came to preforming the column (eluting with a 30 % acetone/*n*-hexanes gradually raised to 100 % acetone in some cases), the well separated fractions did not contain the desired product (**Figure 14**). Due to time constraints, further attempts to yield **6d** were abandoned.

Figure 14: An example of one of the repeated attempts to synthesize **6d** by Tohda's method. (^1H -NMR spectrum)

These compounds (**6a-c**) were analyzed by their characteristic resonance signals in their ^1H - and ^{13}C -NMR spectra. In the ^1H -NMR spectrum, the two methylene groups (both of those on the heterocycle [**1,2**]) resonate in the alkyl region at approximately $\delta_{\text{H}} = 3.86$ and 4.54 ppm for positions **1** and **2**, respectively. The absolute assignment of these two protons can be affirmed by their coupling constants. Where the signal at $\delta_{\text{H}} = 3.86$ ppm has a $^3J = 8.54$ Hz and the $\delta_{\text{H}} = 4.54$ ppm signal has also a $^3J = 8.54$ Hz. Since these two values are the same (as well as taking their splitting patterns into consideration) it can be assumed that they are indeed coupling to one another, making them neighbours and suggesting their assignments are correct.

For this compound, **6b**, another characteristic feature of its isolation lies within the aromatic region. In this region, there are three resonance signals with splitting patterns of a doublet, a doublet of doublets, and a doublet (due to long-range splitting with a close proton) (**Figure 15**). As before, these signals can be assigned by calculating their coupling constants. The coupling constants of those in **Figure 15** labeled as **8**, **12** and **11** are $J = 1.99$ Hz, $J = 8.54$, 1.99 Hz, $J = 8.54$ Hz. Using the same path of thinking as before the protons can be properly assigned. Displaying a smaller coupling constant is indicative of coupling with a proton neighbor more than 3 bonds away. Here, protons **8** and **12** are interacting and because of this give the ^4J coupling constant of 1.99 Hz. The same can be said for proton **11**, as it only interacts with proton **12**. Thus, giving the ^3J coupling constant of 8.54 Hz. With the calculated coupling constants as well as analyzing their splitting pattern protons **8**, **12** and **11** can be and have been correctly assigned. This method was taken into consideration when assigning the protons and carbons (above, **Sec. 2.3.1**) for both **6a** and **6c**.

Figure 15: The ^1H -NMR spectrum for **6b**

2.5 Synthesis and Spectrochemical Investigation of Transition Metal Complexes

Using the fifteen derivatives produced of the 2-acylmethyl-2-oxazoline scaffold, the synthesis of several copper and cobalt complexes were undertaken. These complexes were initially made using the preparation described by Gossage *et al.* but was later altered to further optimize reaction conditions.²⁰ Two equivalents of **4** or **6** are dissolved in EtOH and 2.3 equivalents of NEt_3 are added. Separately, 1 equivalent of a metal nitrate salt ($\text{Co}(\text{NO}_3)_2 \cdot 6 \text{H}_2\text{O}$ or $\text{Cu}(\text{NO}_3)_2 \cdot 2.5 \text{H}_2\text{O}$) is dissolved in EtOH. The metal solution is syringed into the ligand solution while stirring. A noticeable colour change occurs and, in most cases, immediate formation of a coloured precipitate is noted (**Scheme 30**).

Scheme 30: A general metal complexation reaction of **4** or **6** to either **Cu** or **Co**.

Another explored route was that shown in **Scheme 31**. This involves *in situ* generation of the ligand where it immediately coordinates to the metal. Potassium hydroxide is dissolved in MeOH to make a 1.5 molar solution. Once dissolved the intermediate species is added, at room temperature, to the basic mixture. After fully dissolved, the metal salt is added to the stirring solution as a solid. After approximately 10 minutes, a colour change can be observed from the starting dark red colour to a lighter orange (cobalt) or dark green (copper) with a notable amount of precipitate. Once the reaction is done, all volatile components were removed. The blackish sludge is diluted with DCM and washed with a 1.0 M solution of NaCl in water. In the cobalt complexation reaction, 3 layers are obtained in the separatory funnel; the clear aqueous layer, the bright red organic layer, and a third black layer (not observed in the copper complexation reaction). This black layer likely contained some Co(OH)_2 , insoluble in most organic solvents (and water to an extent). The organic phase is dried (MgSO_4), filtered and all volatile components are removed. In the case of both reaction pathways, the precipitates are recrystallized by slow evaporation of a solution of 1:1 DCM/MeOH. This method is effective for all ligands tested as detailed below.

Scheme 31: An alternate route to Cu/Co complexes from **3** or **5**.

2.5.1 Copper Complexes (**7a-l**)

One alkyl and eight aryl derivatives of the copper complexes were made in moderate to excellent yields (**Scheme 32**) and synthesized in one of two methods previously described. The colour of the reaction solutions was initially clear or pale yellow (depending on the colour of the pure ligand) and turned a dark green-brown when the sky-blue metal salt was added. All copper complexes were characterized by cyclic voltammetry (CV), elemental analysis (EA), their melting point (mp), IR spectroscopy, Ultraviolet-visible spectroscopy (UV-Vis), and in some cases, single crystal x-ray diffraction.

Scheme 32: Towards copper complexes with the ligand series **4**.

Since these complexes have a presumed metal electron configuration of d^9 , they are paramagnetic in nature, meaning they have one unpaired electron. This unpaired electron makes analyzing by NMR spectroscopy less conclusive. In the ^1H -NMR spectrum, the closer a proton is to the paramagnetic center the broader the resonance signal will be for that proton (**Figure 16**).⁶⁸ Due to this behaviour, other methods of characterization had to be used such as IR, X-ray, and UV-Vis.

Infrared spectroscopy shows the interaction of the ligand with the metal by observing a shift in the absorbance frequency when comparing to the free ligand. The stretching frequency assigned to either the C=C or C=N bond(s) resonate within a range of $1514\text{--}1548\text{ cm}^{-1}$, which is far lower than the reported values of the bare ligands ($1615\text{--}1635\text{ cm}^{-1}$).³⁵ The Cu center is presumably drawing electron density away from the bound ligand.

Figure 16: ^1H -NMR spectrum of **7a** (paramagnetic).

Figure 17: An example of a UV-Vis spectrum of copper complexes (**7e**).

The UV-Vis spectra bore assigned d-d transitions in the visible region and metal-to-ligand charge transfer (MLCT) bands in the ultraviolet region (**Figure 17**). The lower energy absorptions, in the ranges of 423-457 nm, are representative of a d-d transition. These transitions are seen due to the dark green-orange colour of the copper complexes (in solution). This higher energy absorptions in the range of 243-276 nm, have been assigned to the π - π^* transition localized about either the aromatic ring (C=C) or the imine (C=N) within the oxazoline ring. This is likely the case as MLCT transition bands are primarily observed in low oxidation state metals whereas LMCT bands are more often observed in higher oxidation state transition metal complexes. The spectroscopic data obtained had confirmed the electronic properties of the complexes, such as aiding in differing the electron withdrawing complexes, **7b**, from the electron donating complexes, **7d**. The complexes bearing electron withdrawing ligands, **7b/e-f**, experience a red shift (to lower frequencies).

To elucidate the exact structure of the copper complexes, we turned to single-crystal x-ray diffraction (x-ray crystallography). The structures were collected by Dr. Alan J. Lough at the University of Toronto at 150 K. Single crystals were obtained after slow evaporation of a 1:1 solution of DCM/MeOH. The crystals were washed with cold MeOH and dried over vacuum. This revealed the crystals as dark green-brown needles for **7a/c-i**, as well as a mixture of green plates and brown needles for **7b**. X-ray crystallography was only performed on **7g-h** to help confirm the relative structure of the other derivatives. Structures of **7a** and **7i** have already been solved by our group.^{20,66} **Figure 18** shows the ORTEP drawings of **7g** and **7h** obtain *via* x-ray crystallography.

Figure 18: ORTEP drawings of the unit cell of **7g** and **7h** (x-ray crystallographic representation)

Of these two compounds, **7g** exhibits a square planar geometry around the metal center while **7h** (along with other **Cu** complexes analyzed by our group) exhibit a distorted square planar geometry. The geometry can be properly determined by calculating the τ^4 value (geometry index).^{69,70} When the calculated geometry index is 0 the complex is said to be square planar, whereas then the index is calculated to be 1 the complex holds a tetrahedral geometry. Pictured below in **Equation 2** are the calculations for their τ^4 values. Where θ is the ideal bond angle for a tetrahedral compound (109.5°), and where α and β are the two largest angles amongst the coordination center. The calculated values for **7g** and **7h** as well as other similar copper complexes by our group are in agreeance with those in the literature as d^9 copper complexes normally adopt

a more square planar geometry. A reason as to why **7g** adopts the perfect square planar geometry, where **7h** does not, may be due to the steric effects of the ligand. The ligand (**4g**) when bonded to the copper center presumably imposes an axis of chirality, there the naphthyl rings prohibit free rotation about the Cu center. The methyl groups on the oxazoline as well as the naphthyl group appended on the opposite end of the ligand make it so that it can lock the orientation of the complex, making it unable to twist in such a way that **7h** has. This phenomenon has been observed previously by our group where the oxazoline has been modified to a bulky chiral derivative. The chiral ligand locks the rotation of itself, against the other ligand, pinning it in place.^{20,66} As per their unit cell parameters, both compounds crystallized in the triclinic crystal system and both adopt the P-1 space group. Relevant bond lengths and angles are displayed in

Table 1 and **Table 2**. The angles are used to calculate the τ^4 values, which give the corresponding value concluding their distorted square planar geometry. **7h**, giving a τ^4 value of 0.49 confirms the distorted geometry, adopting a more seesaw geometry about the Cu center, whereas **7g** adopts a perfect square planar geometry. The average length for a Cu-O bond is 1.982 Å and for a Cu-N bond is 2.036 Å. The experimental bond lengths and angles for **7g** and **7h** are consistent with other N,O-donor ligand complexes in the literature.^{20,47}

Equation 2: Calculating the geometry index of **7g** and **7h**.

7g	$\tau^4 = \frac{360^\circ - (\alpha + \beta)}{360^\circ - 2\theta}$ $\tau^4 = \frac{360^\circ - (180 + 180)}{360^\circ - 2(109.5)}$ $\tau^4 = 0.00$	7h	$\tau^4 = \frac{360^\circ - (\alpha + \beta)}{360^\circ - 2\theta}$ $\tau^4 = \frac{360^\circ - (144.37 + 146.69)}{360^\circ - 2(109.5)}$ $\tau^4 = 0.489$

Table 1: Relevant bond angles (°) and bond lengths (Å) for **7g**

Bond Lengths (Å)		Bond Angles (°)	
Cu(1)-O(1)#1	1.9122(10)	O(1)#1-Cu(1)-O(1)	180

Cu(1)-O(1)	1.9122(10)	O(1)#1-Cu(1)-N(1)	88.54(5)
Cu(1)-N(1)	2.0007(11)	O(1)-Cu(1)-N(1)	91.46(5)
Cu(1)-N(1)#1	2.0007(11)	O(1)#1-Cu(1)-N(1)#1	91.46(5)
		O(1)#1-Cu(1)-N(1)#1	88.54(5)
		N(1)-Cu(1)-N(1)#1	180

Table 2: Select bond angles (°) and bond lengths (Å) for **7h**.

Bond Lengths (Å)		Bond Angles (°)	
Cu(1B)-O(1B)	1.923(2)	O(1B)-Cu(1B)-O(3B)	144.32(9)
Cu(1B)-O(3B)	1.928(2)	O(1B)-Cu(1B)-N(2B)	97.60(10)
Cu(1B)-N(2B)	1.933(2)	O(3B)-Cu(1B)-N(2B)	92.58(9)
Cu(1B)-N(1B)	1.942(2)	O(1B)-Cu(1B)-N(1B)	92.50(9)
		O(3B)-Cu(1B)-N(1B)	98.69(9)
		N(2B)-Cu(1B)-N(1B)	144.66(10)

For compound **7b**, two different forms of crystals were obtained. From a reaction where the **Cu** metal salt was CuBr₂ the crystals took the form of brown needles. Whereas when the metal salt was Cu(NO₃)₂ · 2.5 H₂O, the crystals took the form of green plates (**Figure 19**). This strange type of behaviour was first noted when the metal salt was first added to the stirring ligand solution where the copper(II)bromide solution was notably dark brown in colour, and the copper(II)nitrate solution was a lighter brown-green colour. Once the precipitates were collected the colour difference was more noticeable, where the nitrate derived complex appeared as a dark green powder and the bromine derived complex appeared as a dark brown powder. Upon crystallization and analysis by IR spectroscopy, an obvious variance between them was revealed. The brown needles gave the C=N/C=C bond stretch at 1455 cm⁻¹, whereas the green plates gave, presumably, the same IR bond stretch at the much higher value of 1531 cm⁻¹.

Figure 19: Possible polymorphic behaviour in **7a/b**.

When sent for crystallographic analysis the green plates were determined to be the same material that was previously published by Gossage *et al.*²⁰ The brown needles have yet to be analyzed by x-ray crystallography but are assumed to be a polymorph of **7a**, perhaps differing by packing within the unit cell. This behaviour has been assumed for complex **7b** as it also displayed similar characteristics. **7b** precipitates out of solution as a brown or green powder (copper(II)bromine and copper(II)nitrate respectively) but when crystallized, both take the same crystal form (**Figure 20**).

Figure 20: Both brown and green powders crystalline forms for **7b**.

To work catalytically, many Cu and Co systems function by reversible redox events taking place within the metal complex, whether it's a reversible oxidation or reduction or both. This can be explained by cyclic voltammetry (CV). Thus, CV was performed by taking a blank sample of only the electrolyte, dissolving ~193 mg of $[N(n\text{-Bu})_4]^+[\text{PF}_6]^-$ in sparged DCM (0.1 M) then dissolving ~20 mg of **7**. After blank and sample analysis, ferrocene (0.1 mg) was added to the solution to serve as an internal standard. All scans, unless otherwise stated, started with the oxidative sweep then switched to the reductive sweep at the indicated potential. The scan rate was held constant at 100 mV/s for all scans unless otherwise stated. When tested for their redox capabilities, none of the synthesized Cu complexes displayed favourable redox reversibility.

Figure 21: Voltammogram of **7h/i** displaying irreversibility and ligand reduction.

Instead, they either exhibited irreversible oxidation presumably from Cu(II) to Cu(III), poor reversibility (**7h**) or in the case of **7i**, likely reduction of the *p*-NO₂ group within the ligand. This was the case regardless of the scan rate used, step in potential or direction of initial potential (starting with the reductive sweep instead of an oxidative sweep). In each case (alkyl and aryl), these undesirable properties were observed (**Figure 21**). In the interest of making new catalysts, these copper complexes would likely not be active in redox mediated chemistry and therefore we have halted their study in this area at this time.

2.5.2 Other Copper Complexes (**8a-c**)

One aryl derivative was successfully synthesized (**8a**, 70 % yield) and two other aryl derivatives were attempted (**8b** and **8c**). Their synthetic scheme followed one of two methods previously described (**Scheme 33**). All copper complexes were characterized by cyclic voltammetry (CV), their melting point (mp), IR spectroscopy, ultraviolet-visible spectroscopy (UV-Vis), and mass spectrometry (MS).

Scheme 33: Synthesis of **Cu** complexes with **6**.

As in the preceding section, these complexes have a formal metal based electron configuration of d^9 meaning they are paramagnetic in nature. This coupled with their limited solubility in any organic solvent made them very difficult to characterize. Unfortunately, due to this solubility issue (insoluble even in DMSO and DMF), the compounds were unable to be recrystallized for x-ray crystallography purposes.

Unfortunately, **8a** did not give a molecular ion (M^+) when analyzed by mass spectrometry. Due to the poor results of MS, **8a** and **8b** were sent for elemental analysis (EA) [done by Atlantic Microlabs]. The analysis for **8a** came back positive, giving elemental percentages within experimental error. In contrast, **8b** negative results came back, stating that correct compound was

not present or highly contaminated. **8a** could be dissolved enough to perform UV-Vis spectroscopy (as a suspension in solution). Using the EA data, the molar extinction coefficient (ϵ) was estimated (**Figure 22**). **8a** exhibited a similar UV-vis profile to those copper complexes known in the literature where there are three distinct absorption maxima present in the ultraviolet region (~324, ~285, ~240 nm) [MLCT].⁷¹ In the IR spectra, **8a** showcases similar profiles to complexes **7a-i** with two main strong signals at frequencies of 1540 and 750 cm^{-1} .

Figure 22: UV-Vis profiles for **8a** showing three main absorption maxima.

The attempted synthesis of compound **8c** was carried out by adding $\text{Cu}(\text{NO}_3)_2$ to a 1.5 M methanolic potassium hydroxide solution after **5d** had dissolved. The reaction was stirred at room temperature for 22 hours in open air. After the reaction was completed, the volatile components were removed and the sludge was diluted with DCM. The organic phase was washed twice, once with a 1.0 M solution of NaCl and a second time with a 0.1 M solution of Na_2CO_3 . The organic phase was dried (MgSO_4), filtered and volatile components were removed. The resulting brown sludge was attempted to be recrystallized by two methods; the first was a layering of a mixture of

1:1 DCM/MeOH and the second being a vapor diffusion of diethyl ether into a solution of DCM and the complex. Unfortunately, none of these two pathways yielded any pure materials.

2.5.3 Cobalt Complexes (**9a-l**)

Four alkyl and eight aryl derivatives of the cobalt complexes were synthesized in moderate to excellent yields (50-99 %) in one of two methods previously described (**Scheme 34**). The colour of the reaction solutions was initially clear or pale yellow (depending on the colour of the pure ligand) and turned a dark orange-red when the magenta-coloured metal salt was added. All cobalt complexes were characterized by cyclic voltammetry (CV), elemental analysis (EA), their melting point (mp), IR spectroscopy, ultraviolet-visible spectroscopy (UV-Vis), and in some cases, single crystal x-ray diffraction.

Scheme 34: Towards cobalt complexes with the ligand series **4**.

This series of cobalt complexes adopt the presumed d^7 electron configuration, making the paramagnetic analogous to the aforementioned copper complexes. This nature makes characterizing

by NMR spectroscopy difficult as most resonance signals appear as broad singlets (**Figure 23**). The signals that are more deeply broadened are closer to the **Co** center whereas the sharper signals are further away. Using this, we can approximate the structure although this approximation would be a rough guess, at best. Similarly to the **Cu** complexes, as the **Co** complexes display this paramagnetic behaviour they were analyzed by IR, UV-Vis, EA, their mp, MS, and in most cases by x-ray crystallography.

Figure 23: ^1H -NMR spectrum of **9a** (paramagnetic).

Infrared spectroscopy showed the interaction of the ligand with the metal by displaying a shift in the absorbance frequency when compared to the ligand. The stretching frequency assigned to either C=C or C=N stretching resonated within a range of $1498\text{--}1525\text{ cm}^{-1}$, which is far lower than the reported values of the bare ligands within the range of $1615\text{--}1635\text{ cm}^{-1}$.³⁵ The C-O stretching frequencies were seen at frequencies of $753\text{--}767\text{ cm}^{-1}$, indicative of a change from the

free ligand.³⁵ Due to the drastic shift in stretching frequencies in the IR spectra, we hypothesize that the Co center is drawing electron density away from the bound ligand.

The UV-Vis spectra displayed weak d-d transitions in the visible region and metal-to-ligand charge transfer (MLCT) bands in the ultraviolet region (**Figure 24**). Lower energy absorptions, in the ranges of 534-550 nm, are representative of d-d transitions. These transitions are observed due to dark red/pink colour of the solution. This higher energy absorptions in the range of 296-321 nm, have been assigned to the $\pi-\pi^*$ transition localized to either the aromatic ring (C=C) or the imine group (C=N). As is the case for the copper complexes, MLCT transition bands are primarily observed in low oxidation state metals, such as Co^{2+} . Complexes bearing electron withdrawing ligands, **9b/e-f**, as well as conjugation extending functional groups ($-\text{NO}_2$ etc.) will experience a bathochromic shift (to a higher wavelength).

Figure 24: UV-Vis trace of **6e**.

Figure 25: ORTEP representations of **9j-l**.

X-ray crystallography was employed to determine the exact structure for several of the **Co** complexes. These were sent to the University of Toronto, where Dr. A. J Lough analyzed and solved each of their structures. Only three structures and their respective data sets were collected (**9j-l**), **Figure 25**, as our group has previously structurally characterized complexes **9a-c** and **9g-i**.⁶⁶ X-ray quality crystals are grown with slow evaporation of a 1:1 solution of DCM and MeOH. Compounds **9a-f/i** crystallized as orange needles, **9g** crystallizes as dark red cubic shapes, and complexes **9h/j-l** crystallize as pink rectangular plates.

The calculated τ^4 values for these compounds are displayed in **Table 3** and suggest the complexes adopt a more tetrahedral geometry. This approximation is supported by the relevant bond angles in

Table 4, **Table 5**, and **Table 6**. Based on the unit cell parameters all three complexes crystallized in the orthorhombic crystal system with space groups $Pca2_1$, $Pbca$, and $Pna2_1$, for **9j-l** respectively. It is expected for the Co-N bond to be longer than the Co-O bond as the former is a coordination bond, meaning it is not a true bond but a mutual sharing of electrons between the two atoms. The average length for a **Co-O** bond is 1.932 Å and for a **Co-N** bond is 2.035 Å. The

experimental bond lengths and angles for **9j-l** are consistent with other N,O-donor ligand complexes in the literature.^{66,72}

Table 3: Calculated τ^4 values for all **Co** complexes

Complex	τ^4	Complex	τ^4
9i w/ DCM	0.82	9b w/ DCM	0.82
9a	0.83	9g	0.82
9h	0.81	9j	0.84
9i w/ CHCl₃	0.82	9k	0.83
9c	0.78	9l	0.80

Table 4: Selected bond lengths (Å) and angles (°) for **9j**

Bond Lengths (Å)		Bond Angles (°)	
Co(1)-O(3)	1.935(3)	O(3)-Co(1)-O(1)	118.64(13)
Co(1)-O(1)	1.944(3)	O(3)-Co(1)-N(2)	95.42(13)
Co(1)-N(2)	1.957(3)	O(1)-Co(1)-N(2)	112.90(11)
Co(1)-N(1)	1.958(3)	O(3)-Co(1)-N(1)	113.84(12)
		O(1)-Co(1)-N(1)	95.03(13)
		N(2)-Co(1)-N(1)	122.89(16)

Table 5: Selected bond lengths (Å) and angles (°) for **9k**.

Bond Lengths (Å)		Bond Angles (°)	
Co(1)-O(3)	1.9315(19)	O(1)-Co(1)-O(3)	116.69(8)
Co(1)-O(1)	1.9449(19)	O(1)-Co(1)-N(1)	94.87(9)
Co(1)-N(2)	1.964(2)	O(3)-Co(1)-N(1)	113.29(9)
Co(1)-N(1)	1.964(2)	O(1)-Co(1)-N(2)	112.19(9)
		O(3)-Co(1)-N(2)	94.78(9)
		N(1)-Co(1)-N(2)	126.71(9)

Table 6: Selected bond lengths (Å) and angles (°) for **9l**.

Bond Lengths (Å)		Bond Angles (°)	
Co(1)-O(3)	1.936(3)	O(3)-Co(1)-O(1)	117.13(15)
Co(1)-O(1)	1.949(3)	O(3)-Co(1)-N(2)	111.12(15)
Co(1)-N(2)	1.963(4)	O(1)-Co(1)-N(2)	94.21(14)
Co(1)-N(1)	1.968(4)	O(3)-Co(1)-N(1)	94.42(15)
		O(1)-Co(1)-N(1)	111.09(15)
		N(2)-Co(1)-N(1)	130.67(17)

As was for the copper complexes, cyclic voltammetry was employed to probe the redox properties of the cobalt complexes. All scans started with an oxidative sweep following by the reductive sweep with a forward scan rate of 100 mV/s. The **Co** complexes were prepared in the same manner as the **Cu** complexes by first running a blank scan of a 0.1 M solution of [N(*n*-Bu)₄]⁺[PF₆]⁻ followed by the sample (~20 mg). Ferrocene (0.1 mg) was added afterwards to serve as an internal standard. All derivatives, **9a-l**, were analyzed in this way. All complexes excluding **9h**, showed irreversible oxidation. Previous students in the Gossage group attempted polymerization by ATRP with **9a** and other derivatives but obtained no polymer material at the terminus of their reaction.⁶⁶ Complex **9h** was the only complex that displayed quasi-reversible oxidation from Co²⁺ to Co³⁺ at approximately +1.75 eV, making it a potential target for catalytic studies (*vide infra*).

Figure 26: Voltammograms for **9e** and **9h** in DCM.

2.5.3.1 Attempted Oxidation of a Co^{2+} Complex

Upon observing promising redox behaviour in the CV, we decided to attempt various reactions to purposely oxidize the Co center to a permanent Co(III) resting state. This type of reaction was tried four different ways with **9a** and once with **9h** (Table 7).

Table 7: Attempted oxidation of a Co^{2+} center.

Trial	Complex	X_2	Time (hr)	$9:\text{X}_2$	Notes	Product
1	9a	Br	0.5	1:1	light	* CoBr_2
2	9a	Br	0.25	2:1	dark	* CoBr_2
3	9a	I	1	1:1	-	9a
4	9a	I	1	1:1	-	9a
5	9h	Br	1	1:1.4	with ACN	^green oil/orange powder

In trial 1, **9a** was dissolved in toluene and bromine (Br_2) was added dropwise in a 1:1 molar ratio. During addition of Br_2 , it was noted that the solution had started to phase separate leaving a dark red top layer and a blue oil on the bottom of the flask. The reaction was done uncovered in open air for 0.5 hours. Upon removal of the excess toluene and bromine, a fragile blue-green solid. The solids were insoluble in most organic solvents (CDCl_3 , C_6D_6 , toluene, CHCl_3) but was soluble in acetone, MeOH and water. An NMR spectrum was taken in acetone- d_6 and unfortunately from

this there were no distinct signals in either the proton or carbon spectra. A qualitative test was performed by adding both DCM and water to the flask and rapidly stirring for approximately 10 minutes. This resulted in the loss of the original green colour and appearance of a pink aqueous layer and cloudy organic layer (**Figure 27, B-C**). After NMR analysis, the sample was poured into a beaker where it was diluted further with acetone. Commercial grade CoBr_2 was dissolved in both acetone and MeOH in two other beakers and the three compared (**Figure 27, A**). It seemed likely that CoBr_2 was indeed the formed product and hence decomposition of **9a** is likely the result of the above treatments.

Figure 27: A qualitative test of Trial 1 (**9a** + Br_2)

For the second trial, the reaction was carried out in a similar fashion, this time stirring in the dark. After the reaction was done and excess components removed *in vacuo* a blue powder was left behind. The blue powder was recrystallized with a 1:1 mixture of DCM/MeOH as was the case for **7** and **9**. Upon accidental contact with water the blue powder was instantly dissolved. There was a notable colour change from light blue to a pinky-purple colour. Commercial grade CoBr_2

was likewise dissolved in EtOH and revealed similar behaviour. After more research, the blue powder was concluded to be anhydrous CoBr_2 .

Trials 3 and 4 gave the same result of recovery of starting material. The iodine, in both polar aprotic and nonpolar solvents, remained unreacted in each case. This was later confirmed by NMR spectroscopy where the same broad signals associated with **9a** (**Figure 23**) were observed.

Scheme 35: Attempted bromination of **9h**.

Trial 5 was done with **9h**, the catalytic candidate (**Scheme 35**). **9h** was initially dissolved in ACN where the solution instantly changed from a light pink to a dark purple. This colour change is assumed to be coordination of either one of two ACN molecules to the cobalt center, although this has yet to be confirmed. This complex would still be paramagnetic and therefore would not be observable by NMR spectroscopy. Continuing with the reaction once fully dissolved, bromine was added dropwise and the solution took a noticeable colour change to a dark green. Once all volatile components were removed a green resin was left behind. This resin was soluble in CHCl_3 and acetone although did not turn the bright teal-blue colour once dissolved in acetone, possibly being the target complex.

Performing NMR spectroscopy on the green resin in CDCl_3 gave a paramagnetic-like spectrum with broad signals spanning many ppm units (**Figure 28**). The green oil was dissolved in MeOH, where the solution was a bright pink colour. Once the MeOH is removed, the compound becomes a resin, again green in colour. We hypothesize this may be signaling the release and re-

coordination of a ligand (Br^- or ACN) giving back **9h**. These mixtures did not yield a pure oxidized product of **9h**. More studies in this area needs to be conducted for an exact conclusion to be drawn.

Figure 28: ^1H -NMR spectrum of proposed **9h-1**.

2.5.4 Cobalt Complexes (**10a-d** and **11a/11c**)

Cobalt complexes with ligand series **6** were also synthesized. In this series, three aryl derivatives were successfully synthesized (**Scheme 36**).

Scheme 36: Synthesis of complexes **10a-d**.

The synthesis was carried out in a method like those described above where to a stirring solution of the ligand with 2.5 equivalents of NEt_3 dissolved in EtOH, 1 equivalent of a metal salt is added (also dissolved in EtOH). Upon addition of the **Co** metal salt to the ligand solution, an evident colour change from dark red/clear to a pastel orange is observed (**Figure 29, B**).

Figure 29: A- **10a** and **10c** as a powder. B- RIGHT: **10c** (orange) during synthesis.

After which, the orange/tan precipitates are collected *via* vacuum filtration and are recrystallized with a layering of 1:1 DCM/MeOH (**Figure 29, A**). These compounds, where applicable, were characterized by IR, UV-Vis, their mp, EA and in some cases, x-ray crystallography. **Co** complex **10d** was unable to be synthesized regardless of the route taken.

Infrared spectroscopy of **10a-c** indicated an interaction between the ligand and the metal by displaying a shift in the absorbance frequency when contrasted against the bare ligand. The stretching frequency assigned to either C=C or C=N stretching resonated within a range of 1527-1531 cm^{-1} . The C-O stretching frequencies were seen at frequencies of 746-770 cm^{-1} , indicative of a change from the free ligand which bears no C-O bond but a carbonyl stretch.^{9,38,39} Due to the drastic shift in stretching frequencies in the IR spectra we conclude that the **Co** center is drawing electron density away from the bound ligand.

The UV-Vis spectra displayed weak d-d transitions in the visible region and MLCT bands in the ultraviolet region (**Figure 30**). Lower energy absorptions, in the ranges of 528-548 nm, are characteristic of d-d transitions. The orange/tan colour of the solutions are dependent on these d-d transitions as they transmit the complementary colour. The higher energy absorptions in the range of 235-242 nm and 310-321 nm, have been assigned to the $\pi-\pi^*$ transition localized to either the aromatic ring (C=C) or the imine group (C=N).

Figure 30: UV-Vis trace for **10b**.

Crystals of compound **10a** were grown from vapor diffusion of diethyl ether into a solution of warm DCM. These crystals were sent to the University of Toronto for structural analysis. Complex **10a** crystallizes as dark red cubes. The calculated τ^4 value for this complex is 0.83, giving it a more tetrahedral-like geometry. **10a** crystallizes in the monoclinic crystal system with an assigned space group of $P2_1/n$, supporting its geometry. Relevant bond lengths and angles are shown in **Table 8**, which are comparable to other Co N,O bidentate complexes.⁶⁶

Table 8: Selected bond lengths (Å) and angles (°) for **10a**.

Bond Lengths (Å)		Bond Angles (°)	
Co(1)-O(3)	1.9348(12)	O(1)-Co(1)-O(3)	108.80(5)
Co(1)-O(1)	1.9360(12)	O(1)-Co(1)-N(2)	116.97(6)
Co(1)-N(2)	1.9588(14)	O(3)-Co(1)-N(2)	94.75(5)
Co(1)-N(1)	1.9621(14)	O(1)-Co(1)-N(1)	95.14(6)
		O(3)-Co(1)-N(1)	115.29(6)
		N(2)-Co(1)-N(1)	126.20(6)

Cyclic voltammetry was carried out to probe the redox properties of the **10a-c** complexes. All scans were conducted first the oxidative sweep following by the reductive sweep, scanning with a forward scan rate of 1.0 mV/s. The **Co** complexes were prepared in the same manner as the previous complexes by first running a blank scan of a 0.1 M solution of $[N(n\text{-Bu})_4]^+[\text{PF}_6]^-$ followed by the sample (~20 mg). Afterwards, ferrocene (0.1 mg) was added to serve as an internal standard. All derivatives, **10a-c**, were analyzed in this way. All complexes showed irreversible oxidation (Figure 31).

Figure 31: Cyclic voltammetry for **10a** and **10c**

2.5.4.1 Usual Oxidation from Co^{2+} to Co^{3+} Complexes (**11a/c**)

Interestingly, darkening of the crystallization solution of both **10a** and **10c** were observed. When **10a** was initially recrystallized with a 1:1 mixture of DCM/MeOH after approximately 24 hours, black crystals were apparent at the bottom of the vial (**Figure 32, E**). When sent for crystallographic determination, a novel complex, compound **11a** was identified (**Figure 32, F**). This phenomenon was tracked over the course of three days by dissolving **10a** in only DCM and observing changes each day. In each instance, **11a** was obtained.

Figure 32: Solution darkening and oxidation of Co^{2+} to Co^{3+} (**10a** to **11a**)

Experiments were done to try to synthesize **11a** from just the ligand and cobalt metal salt. This was done by dissolving 3 equivalents of **6a** in MeOH and adding 1 equivalent of the **Co** metal salt also dissolved in MeOH. Upon addition, the ligand solution turned a dark brown-red. The reaction was stirred at room temperature for 24 hours. The same reaction was tried a second time, although refluxing (with identical conditions) for 5 hours. In both cases, **10a** was the only product isolated. In another reaction, 1 equivalent of **10a** was added to 1 equivalent of **6a**, attempting to coordinate another ligand, in MeOH. This was left to stir at room temperature. Upon completion,

again the only collected product was **10a**. In another test, 34 mg of **10a** was dissolved in 2 ml of DCM. This mixture could sit till all the solvents had evaporated. The resulting complex, **11a**, was separated and weighted out. **11a** was present, an approximate 50 % conversion from **10a** to **11a**. No other pure materials could be isolated from this mixture.

Similarly, **10c** also exhibits this behaviour, formally oxidizing the metal from a Co^{2+} to Co^{3+} . The reaction is carried out in a similar way to **10a** where 2 equivalents of **6c** and 3 equivalents of NEt_3 are dissolved in EtOH. To this solution, 1 equivalent of $\text{Co}(\text{NO}_3)_2 \cdot 6 \text{H}_2\text{O}$ dissolved in EtOH was added dropwise. After 1.5 hrs, the precipitates were collected (85 %) and recrystallized. Two recrystallization vials were set up; one with layering of DCM and MeOH (1:1) and the second a vapour diffusion of diethyl ether into warm DCM. These two methods revealed red cubic crystals and short black needles (respectively, **10c** and **11c**)

On both compounds (**11a/c**) IR, CV, UV-Vis, EA, their mp, and where applicable x-ray crystallography were carried out. In the IR spectra, a noticeable shift to lower frequencies is observed. In comparison to **10a** ($1537, 746 \text{ cm}^{-1}$) a shift to 1526 and 746 cm^{-1} is observed in the C=N/C=C and C-O stretching. The same is observed to **11c**, a shift from $1498, 770 \text{ cm}^{-1}$ to 1495 and 756 cm^{-1} for the C=N/C=C and C-O stretching respectively.

In UV-Vis spectra slight red-shifting of the absorption maxima in both the d-d transitions and MLCT bands are observed. Lower energy absorptions, in the ranges of 528-549 nm for **10a/c** are shifted to approximately 567 nm with the appearance of a new weak absorption of around 445 nm. The black/green colour of the solutions is dependent on these d-d transitions. The $\pi\text{-}\pi^*$ transitions, localized to either the aromatic ring (C=C) or the imine group (C=N), in the range of 235-240 nm and 312-318 nm, have also been red-shifted slightly to now 244-245 nm and 346-347 nm respectively. Both complexes (**11a** and **11c**) were analyzed by EA. The results came back

within experimental error of their calculated elemental compositions, with the expectation of **11a** which held also half a molecule of DCM within the elemental lattice. With DCM considered, both analyses were considered accurate (within 3 % error).

Both **11a** and **11c** were sent for crystallographic analysis and returned with positive results (**Figure 33**). As per the unit cell parameters, **11a** crystallizes in a triclinic crystal system within the P-1 space group and **11c** crystallizes in a monoclinic crystal system within the P2₁/c space group. Both complexes have octahedral geometry and are both *mer* isomers (all nitrogen/oxygen atoms found along the same meridian). Of the three ligands attached to the Co metal center, in both the case of **11a** and **11c**, there is one ligand with elongated bonds (both Co-O and Co-N) to the center (**Table 9**, **Table 10**).

Figure 33: ORTEP representations of **11a** and **11c**

Table 9: Select bond angles (°) and lengths (Å) for **11a**

Bond Lengths (Å)		Bond Angles (°)			
Co(1)-O(5)	1.8983(16)	O(5)-Co(1)-O(1)	88.61(7)	O(3)-Co(1)-N(2)	92.96(8)
Co(1)-O(1)	1.8993(16)	O(5)-Co(1)-O(3)	174.68(7)	N(3)-Co(1)-N(2)	91.15(8)
Co(1)-O(3)	1.9072(16)	O(1)-Co(1)-O(3)	89.04(7)	O(5)-Co(1)-N(1)	88.15(7)
Co(1)-N(3)	1.9144(19)	O(5)-Co(1)-N(3)	94.33(7)	O(1)-Co(1)-N(1)	92.11(7)
Co(1)-N(2)	1.915(2)	O(1)-Co(1)-N(3)	84.09(8)	O(3)-Co(1)-N(1)	87.16(7)
Co(1)-N(1)	1.9219(19)	O(3)-Co(1)-N(3)	90.18(8)	N(3)-Co(1)-N(1)	175.40(8)
		O(5)-Co(1)-N(2)	89.78(8)	N(2)-Co(1)-N(1)	92.73(8)
		O(1)-Co(1)-N(2)	174.84(7)		

Table 10: Select bond angles (°) and lengths (Å) for **11c**

Bond Lengths (Å)		Bond Angles (°)			
Co(1)-N(3)	1.892(4)	N(3)-Co(1)-O(4)	91.27(16)	N(2)-Co(1)-O(1)	87.30(16)
Co(1)-O(4)	1.896(3)	N(3)-Co(1)-O(7)	93.73(16)	N(3)-Co(1)-N(1)	90.92(17)
Co(1)-O(7)	1.902(3)	O(4)-Co(1)-O(7)	173.77(14)	O(4)-Co(1)-N(1)	87.06(16)
Co(1)-N(2)	1.906(4)	N(3)-Co(1)-N(2)	89.22(17)	O(7)-Co(1)-N(1)	89.14(16)
Co(1)-O(1)	1.912(3)	O(4)-Co(1)-N(2)	93.58(16)	N(2)-Co(1)-N(1)	179.35(18)
Co(1)-N(1)	1.913(4)	O(7)-Co(1)-N(2)	90.21(16)	O(1)-Co(1)-N(1)	92.88(16)
		N(3)-Co(1)-O(1)	176.07(16)	O(7)-Co(1)-O(1)	88.11(14)
		O(4)-Co(1)-O(1)	87.13(15)		

Complexes **11a** and **11c** have been also tested for their redox behaviour using previously described methods. Unfortunately, only irreversible behaviour was observed, regardless of scan rate and starting potential, for both complexes (**Figure 34**).

Figure 34: Voltammogram of **11a** and **11c** in DCM.

2.6 Polymerization Studies with a Co^{2+} Catalyst

The successful observation of the redox properties of **9h** lead to testing of its abilities as a catalyst in polymerization. Thus, this reversible oxidation metal catalyst was tested in atom transfer radical polymerization (ATRP).

Scheme 37: General route to polymerization of MMA.

Each reaction trial was set up in the same way where all glassware was flame-dried and purged of air. The catalyst, monomer and initiator were dissolved in 3 ml of the dry solvent specified in **Table 11**. For all polymerizations, tosyl chloride (TsCl) was selected as the initiator due to it being readily available.^{60,73} This mixture was left to heat at the temperature specified in a sealed round-bottom flask for an allotted amount of time (**Table 11**). After completion, the solvent was concentrated to about 1 ml. This mixture was slowly dropped into cold, rapidly stirring MeOH ($-30\text{ }^{\circ}\text{C}$) to reveal the polymer as a white threadlike substance. The complex, **9h**, is soluble in

MeOH. During the precipitation process the MeOH, is filtered and the solvent removed *via* rotary evaporator. In each case, complete recovery of the catalyst can be made. This gives way to be reused in another trial (**Table 11**, entry 8) modest conversion of the monomer to the polymer was observed, giving percent conversions between 14-42, but mostly around 40 %. Polymerization was tried twice without an initiator, as proof of principal. Without the initiator present, no polymer was formed. This further supports the ATRP-like catalytic nature of **9h**.

Table 11: Trials and data for the polymerization of MMA

Trial	9h Mol%	Solvent	Time	Temperature (°C)	Initiator : mol %	Mw (Da) : PDI	Conversion (%)
1	2	THF	24	60	N/A	N/A	0
2	2	Tol.	24	90	N/A	N/A	0
3	2	THF	72	60	TsCl : 0.3	60K : 1.38	38
4	2	THF	24	60	TsCl : 0.3	50K : 1.19	42
5	2	THF	4	60	TsCl : 0.3	53K : 1.17	23
6	5	THF	4	60	TsCl : 0.3	39K : 1.21	14
7	10	THF	4	60	TsCl : 0.3	32K : 1.20	40
8	2	THF	12	60	TsCl : 0.3	45K : 1.18	38

Scheme 38: Formation of PMMA, based on best trial from **Table 11**.

Gel permeation chromatography (GPC) data revealed a generally well controlled polymerization of MMA showing respectable PDI's ranging from 1.17-1.38 with high molecular weights (32-60,000 Da).

Based on the data in **Table 11** and other literature protocols^{60,74}, we propose a mechanism for the polymerization of MMA with **9h** (**Scheme 39**). Upon heating the solution, the initiator (TsCl) radicalizes. A radical finds the catalyst and bonds to the cobalt metal which can accommodate the extra bonded halogen. In this step, the cobalt becomes oxidized to an oxidation state of 3+. The tosyl radical anion interacts with the monomer species by attacking the terminal olefin and creating a bond. This change in electronic structure sends the electron propagating along the rest of the monomer till another monomer near is hit by the radical monomer. Propagation along the main chain continues till there are no monomer units left or till all the radical species have been terminated. During this process, the tosyl group can unbind from the growing polymer chain to stop the polymerization process as the **9h**³⁺-Cl will always be close by the growing chain. When all monomer units have been used up, the chlorine bonded to the metal catalyst will be released and bound to the end of the polymer chain (**Scheme 38**).

Scheme 39: Proposed mechanism for the formation of PMMA via **9h**.

Ongoing experiments are being performed to further optimize not only the reaction yield but also the PDI's of the resulting polymer. Once optimized, **9h** will be tested in the polymerization of other monomers such as styrene.

2.7 Synthesis and Characterization Other Group 9 complexes

Below Cobalt in Group 9 lies both Rhodium and Iridium. In recent years, there have been various studies surrounding their redox properties as well as their candidacy as catalysts. This lead to their increasingly popularity in the field of catalytic-coordination chemistry. Two of the best complexes for studying hydrogen transfer, as well as other general oxidative addition reactions, are Vaska's complex, $\text{IrCl}(\text{CO})(\text{PPh}_3)_2$, and Wilkinson's catalyst. For these reasons, we propose a modification to these complexes to hopefully enrich their redox potential.

2.7.1 Attempted Synthesis and Characterization of Rh Complexes

Initial investigations of other group 9 complexes involved reactions of **4a** with Wilkinson's catalyst. Some groups have been able to oxidatively add neutral and other hydride species to Wilkinson's catalyst. These types of Rh-compounds tend to be used as hydrogenation catalysts.^{75,76} There have been other groups interested in the catalytic properties of nanoparticles, coating and grafting a tether to a particle. Once grafted, they can coordinate Wilkinson's catalyst to it. This makes for a highly effective and selective hydrogenation agent.⁷⁷

Our exploration of this was first done by dissolving Wilkinson's catalyst and **4a** in C_6D_6 while in the glove box. Once out of the glove box, the solution was heated at 50 °C and monitored by its ^{31}P -NMR spectrum (**Figure 35**).

Scheme 40: Attempted substitution of Wilkinson's Catalyst with **4a**.

By analyzing the NMR spectra obtained over the 24-hour period, it was concluded that Wilkinson's catalyst was decomposing in solution, releasing PPh_3 which was oxidized almost entirely to O=PPh_3 . This reaction was tried again, this time refluxing in CDCl_3 for only 6 hours (Scheme 40). Analysis by NMR spectroscopy revealed one signal at +29 ppm (O=PPh_3). Precipitation of Rh metal was also recovered from the reaction, further supporting decomposition of the catalyst. Further reactions with this material were not continued.

Figure 35: ^{31}P -NMR spectra of reaction between Wilkinson's catalyst and 4a.

2.7.2 Attempted Synthesis and Characterization of Vaska-type Complexes

Another typical candidate to study oxidative addition is Vaska's complex and its Rh analogue. Other groups have been able to take advantage of the oxidative properties of Ir^+ . In this regard, various silane species as well as small carbon-tin based spheres have been shown to oxidatively add to the Ir center.^{78,79}

Figure 36: Target group 9 complexes for substitution.

Various attempts were made to substitute ligands on **12** and **13** (**Figure 36**) with **4a**. An NMR tube was charged with 1 equivalent of **12/13** and 1 equivalent of **4a** dissolved in CDCl_3 , heated at $50\text{ }^\circ\text{C}$ and monitored by its ^{31}P -NMR spectrum over 2 days (**Figure 37**). In the reaction with **12**, the disappearance of the main signal in the NMR spectrum at 28.8 ppm as well as the appearance of two new signals, at 29.2 and 20.6 ppm, are observed. The signal at 29.2 ppm corresponds to the formation of triphenylphosphine oxide (O=PPh_3). For it to have been oxidized, PPh_3 must have been released from the complex. The appearance of a new signal at 20.6 ppm could be indicative of a new iridium complex being formed between **12** and **4a** although in all attempts to isolate the new signal from the mixture only **12** was obtained.

In the reaction with **13**, the disappearance of the starting material was observed (29.0 ppm) and the appearance of a new doublet was observed at 43.4 ppm, as well as O=PPh_3 at 29.2 ppm. This new signal at 43 ppm confirmed the formation of a new rhodium complex. Rhodium has a spin of $\frac{1}{2}$, the same as phosphorus, therefore they will couple together to appear as a doublet instead of a singlet. Unfortunately, attempted isolation of this new complex only gave the starting material (**13**) back as crystalline yellow needles.

Figure 37: Substitution of **4a** to either **12** (left) or **13** (right).

Substitution of **12/13** with **4a** was tried again, this time on a slightly larger scale with using toluene as a solvent. These mixtures were refluxed 24 hours before being cooled to room temperature where (20 ml) *n*-hexanes was added dropwise. The **Rh** and **Ir** containing flasks were set aside in a -30 °C freezer for recrystallization. After cooled, the two were subjected to vacuum filtration where in the case of **Rh**, a brown powder was revealed and washed with cold *n*-hexanes and diethyl ether. After dry, a ^{31}P -NMR had shown only one signal at +29 ppm ($\text{O}=\text{PPh}_3$). In the case of **Ir**, a yellow powder was revealed and washed with cold *n*-hexanes and diethyl ether. After dry, a ^{31}P -NMR disclosed 2 signals at +25 and +5 ppm (starting material). In each case, none of the complexes had either reacted (**12-Ir**) with **4a** or only decomposed (**13-Rh**).

The reaction was attempted again, although this time with a synthetic approach more like that of the formation of **Cu/Co** complexes [see **Sec 2.5**]. This was done by adding 1 equivalent of **12** and **4a**, dissolving them in toluene and then added 3.4 equivalents of NEt_3 . It was hoped that the base would make the ligand more nucleophilic towards the **Ir** center. The solution was refluxed for 14 hours. Upon completion, the reaction was cooled to room temperature where *n*-hexanes was layered on top and put into the -30 °C fridge. Once filtered, washed and dried the precipitates were

checked by NMR spectroscopy. Checking by ^{31}P -NMR spectroscopy, however, revealed only starting material (**12**).

2.7.3 Synthesis and Characterization of Ir Complexes (**14** and **15**)

Since group 9 carbonyl/phosphine complexes could not be readily substituted with **4a** another route was taken to explore the reactivity of **4a**. Instead, a structurally strained Ir dimer was made ($[\text{Ir}(\text{COD})\text{Cl}]_2$ (**14**)). The dimer is known to react with N,O-donor, and other pro-chelate ligands, as it readily splits to form a new chelated Ir⁺ complex.^{80–82}

Examples of this reactivity can be found in preparations by Carlton.^{54,80} They've showcased the oxidative properties of their iridium-COD complex by adding HSiPh₃, HSnPh₃ and SC₅F₆ to the iridium center, changing its formal oxidation number to +1 to +3. Using a simple two-step method (formation of an iridium-COD dimer then coordination of their ligand) then by the addition of the nucleophilic species, the Carlton group has been able to synthesise an array of H-Ir-R complexes. This reactions series showcases the redox ability of the iridium centered complex as it oxidizes from an Ir(I) to Ir(III).

Thus, 1 equivalent of **14** was added together with 2.1 equivalents of **4a** in a flask. This flask was purged and backfilled with nitrogen. The two solids were dissolved in dry DCM and 3.1 equivalents of NEt₃ was also added. This solution was left to stir at room temperature for 3 hours under a constant stream of nitrogen gas. At the terminus of the reaction, excess volatile components were removed *in vacuo* and TLC was preformed to observed conversion, if any. After development in 30 % EtOAc/*n*-hexanes, two spots will distinguishable R_f values were obtained (one at 0.26 and the other at 0.56). Column chromatography was performed the material the following day eluting with 25 % EtOAc/*n*-hexanes. Although, after developing TLC plates containing fractions from the

column only revealed a compound with an R_f of 0.26. Re-running the crude sample from the previous days TLC showed also only the one spot. Regardless after analyzing by ^1H and ^{13}C -NMR spectra, the compound which eluted from the column was determined to be **15** (**Figure 38**).

Figure 38: ^1H -NMR spectrum of **15**

The structural identity of **15** was probed by a variety of NMR spectroscopic techniques (both 1D and 2D) and *via* x-ray crystallography. In the ^1H -NMR spectrum (**Figure 38**), each of the signals belonging to the parent ligand experience a slight upfield shift of about 0.2 ppm and a notable disappearance of the NH signal (resonates at ~9.95 ppm in **4a**). This, along with the appearance of the six sets of complex multiplets indicating asymmetry in the ‘COD’ ring, give evidence that a new complex with ‘COD’ incorporated was formed. The ^{13}C -NMR shows a significant downfield shift of the enolic proton from ~58 ppm in **4a** to ~81 ppm in **15**. Assignments to **15** were made with a combination of the assignments of **4a** (made by COSY/HSQC NMR) as well as crude

assignments of the 'COD' ring were done with HSQC and COSY NMR. The olefinic protons/carbons are shifted more downfield, making them distinguishable from the alkyl signals.

The reaction was done a second time, although this time performing preparative TLC (eluting with 25 % EtOAc/*n*-hexanes) of the product mixture the same day to isolate both R_f spots on the TLC plate. When both spots were analyzed by NMR what looked like the same compound being scanned twice appeared. The only different between the two compounds were a fairly large downfield shift of the olefin protons/carbons of the 'COD' ring (**Figure 39**, **Figure 40**). We propose that less polar of the two spots may be an intermediate complex such as an Ir-COD-4a dimer which over time breaks apart into the thermodynamic product, **15**.

Figure 39: ¹H-NMR stacked of both R_f spots on the TLC plate of **15**.

Figure 40: ^1H -NMR stacked of both R_f spots on the TLC plate of **15**.

The synthetic route that gave the best yield of **15** is as described above except instead of 3 hours, the reaction is only stirred for an hour and instead of column chromatography, a rinse with cold MeOH and then decanting off the liquor is used to purify the complex (**Scheme 41**). This method gave the target compound, **15**, in high yields (90 %). **15** exhibits C-O stretching at a frequency of 776 cm^{-1} and the C=N/C=C stretches at 1534 cm^{-1} , in contrast to the free ligand, **4a**, which shows a C=N/C=C stretching frequency at 1620 cm^{-1} .³⁵

Scheme 41: Synthetic route to make **15**

The identity of complex **15** was unequivocally assigned once after it was crystallized and examined by x-ray crystallography. **15** crystallized as yellow cubic structures, grown by slow evaporation of neat CDCl_3 over a period of 30 minutes. As per the unit cell parameters, **15** crystallized in the monoclinic crystal system with the crystal space group $\text{P2}_1/\text{n}$. The average length for a Ir-O bond is 2.106 Å and for a Ir-N bond is 2.081 Å. The Ir-O bond lengths in **15** are slightly shorter than those reported in the literature (**Table 12**) although, other experimental bond lengths and angles for **15** are consistent with other Ir^+ -COD complexes in the literature.^{55,80,83,84}

Table 12: Relevant bond lengths (Å) and angles (°) for **15**.

Bond Lengths (Å)		Bond Angles (°)			
Ir(1)-O(1)	2.039(2)	O(1)-Ir(1)-N(1)	89.87(10)	C(5)-Ir(1)-C(1)	97.48(15)
Ir(1)-N(1)	2.087(3)	O(1)-Ir(1)-C(5)	158.12(13)	C(2)-Ir(1)-C(1)	39.24(13)
Ir(1)-C(5)	2.101(4)	N(1)-Ir(1)-C(5)	96.61(13)	O(1)-Ir(1)-C(6)	159.02(12)
Ir(1)-C(2)	2.108(3)	O(1)-Ir(1)-C(2)	85.69(12)	N(1)-Ir(1)-C(6)	100.44(13)
Ir(1)-C(1)	2.117(3)	N(1)-Ir(1)-C(2)	161.21(13)	C(5)-Ir(1)-C(6)	39.06(13)
Ir(1)-C(6)	2.126(4)	C(5)-Ir(1)-C(2)	81.46(14)	C(2)-Ir(1)-C(6)	89.85(14)
		O(1)-Ir(1)-C(1)	83.31(12)	C(1)-Ir(1)-C(6)	80.32(14)
		N(1)-Ir(1)-C(1)	157.88(14)		

Figure 41: ORTEP representation of **15** (left) and cubic crystals of **15** (right)

2.7.3.1 Oxidation Addition of various substrates to **15** (**16-18**)

Iridium (I) complexes are known in the literature to perform oxidation addition reactions with relative ease with a variety of substrates.^{54,80} Complex **15** was tested for oxidation addition reactions with MeI, HSiCl₃, and HSnPh₃. The first addition to **15** was with MeI. This was done by purging a two-necked flask containing **15** and back-filling with nitrogen gas three times. After assurance of an air free atmosphere, toluene was added to dissolve **15**. Once completely dissolved, a 5-fold excess of MeI was added to the flask. This mixture was left to stir under nitrogen at room temperature for 24 hours (**Scheme 42**). As the reaction proceeded a notable colour change occurred from bright yellow to dark red. A TLC was performed to monitor reaction progress, which showed completion conversion to a less polar substrate and disappearance of starting material. Upon completion of the reaction, the solvent and excess volatile components were removed by a rotary evaporator. When taken off the rotary evaporator, the resulting oil solidified. The brown solids were dissolved in a minimal amount of toluene and precipitated into rapidly stirring cold *n*-hexanes. The tan coloured precipitate was filtered and dried over vacuum, which was ultimately recrystallized with neat C₆D₆.

Scheme 42: Oxidative addition of MeI to **15** to give **16**.

Complex **16** gives a slightly shifted IR spectrum so that the C-O stretch is at 697 cm^{-1} and the C=N/C=C stretching band is now at 1434 cm^{-1} . This is expected seeing as now **16** has a lot more electron withdrawing character about the Ir center when compared to **15**.

This behaviour is also observed in the ^1H and ^{13}C spectra. Upon oxidative addition of MeI, the complex becomes asymmetrical, giving each proton and carbon their own unique chemical environment. This is observed in the ^1H -NMR spectrum where all protons on the oxazoline are diastereotopic, now splitting protons on the same carbon. In **Figure 42** there is a clear splitting of the protons labeled **1** giving rise to two singlets (an apparent doublet) and at the protons labeled **2** giving rise to an ABq splitting pattern. This signal is found at $\delta_{\text{H}} = 3.89\text{ ppm}$. This methodology can be ascertained by viewing the ^{13}C -NMR, where all carbons assigned to the 'COD' ring are dissimilar giving rise to a unique resonance signal for each. All other assignments to **16** were made by analysis by 2D NMR spectroscopy (HSQC, COSY, nOesy). One assignment of note is the one of the proton labeled **15** in **Figure 42**. This assignment was not only important from a completion stand-point but also important in detailing the exact geometry of the complex.

In an oxidative addition reaction, there are three main mechanisms of how the small molecule adds to the complex; by a concerted mechanism (where bond breaking and bond forming happen simultaneously), by a non-concerted mechanism (where bond breaking and bond forming happen one after another) or by radical insertion (where the small molecule radicalizes and adds in a non-concerted manner). In the case of **16**, we proposed a non-concerted reaction where either the iodide anion or the methyl ion first add to **15** in which the other follows. This hypothesis gives the *trans* addition product. This being the case as first either the iodide or methyl group will attack on face of the complex, leaving only the opposite face with room for attack, coming in separately. This is viewed also in the nOesy NMR spectrum where there is a subtle interaction between a few

of the ‘COD’ protons as well as the methyl’s on the oxazoline (**1/1a**) with the methyl group (**15**) directly bonded to the iridium center, proving **15** lays in between both ligands (**Figure 43**, marked).

Figure 42: ^1H -NMR spectrum of **16**.

The absolute geometry was confirmed by growing x-ray crystallography quality crystals by slow evaporation in neat C_6D_6 (**Figure 44**). **16** crystallized as brown needles in the monoclinic crystal system within in the $\text{P}2_1/\text{n}$ space group. This is ascertained by the unit cell parameters. Bond angles and Ir-O, Ir-N and Ir-C bond lengths are comparable to other Ir^{+3} -COD complexes (

Table 13).^{55,80,83,84}

Figure 43: ^1H - ^1H nOesy NMR spectrum of **16**. Marked is the CH_3 (**15**) interactions with the oxazoline protons (**1/1a**).

Figure 44: ORTEP representation of **16**.

Table 13: Select bond lengths (Å) and angles (°) for **16**.

Bond Lengths (Å)		Bond Angles (°)		Bond Angles (°)	
Ir(1)-O(1)	2.0426(19)	O(1)-Ir(1)-N(1)	91.57(8)	C(14)-Ir(1)-C(18)	94.65(12)
Ir(1)-N(1)	2.108(2)	O(1)-Ir(1)-C(22)	82.21(10)	O(1)-Ir(1)-C(21)	157.73(9)
Ir(1)-C(22)	2.184(3)	N(1)-Ir(1)-C(22)	79.87(10)	N(1)-Ir(1)-C(21)	103.83(10)
Ir(1)-C(17)	2.187(3)	O(1)-Ir(1)-C(17)	83.47(10)	C(22)-Ir(1)-C(21)	115.93(11)
Ir(1)-C(14)	2.191(3)	N(1)-Ir(1)-C(17)	158.96(11)	C(17)-Ir(1)-C(21)	87.35(11)
Ir(1)-C(18)	2.204(3)	C(22)-Ir(1)-C(17)	79.20(12)	C(14)-Ir(1)-C(21)	37.10(11)
Ir(1)-C(21)	2.211(3)	O(1)-Ir(1)-C(14)	156.93(10)	C(18)-Ir(1)-C(21)	78.67(11)
Ir(1)-I(1)	2.8598(2)	N(1)-Ir(1)-C(14)	97.87(10)	O(1)-Ir(1)-I(1)	85.50(6)
		C(22)-Ir(1)-C(14)	78.83(11)	N(1)-Ir(1)-I(1)	81.65(6)
		C(17)-Ir(1)-C(14)	80.19(11)	C(22)-Ir(1)-I(1)	157.45(8)
		O(1)-Ir(1)-C(18)	81.78(10)	C(17)-Ir(1)-I(1)	118.12(9)
		N(1)-Ir(1)-C(18)	161.82(11)	C(14)-Ir(1)-I(1)	116.60(8)
		C(22)-Ir(1)-C(18)	115.62(11)	C(18)-Ir(1)-I(1)	80.98(8)
		C(17)-Ir(1)-C(18)	37.21(12)	C(21)-Ir(1)-I(1)	81.03(8)
		O(1)-Ir(1)-N(1)	91.57(8)	C(14)-Ir(1)-C(18)	94.65(12)

The second oxidation addition reaction tried upon **15** was that with HSiCl₃. This reaction was carried out in a similar manner to that of **16**. In a small round bottom flask, **15** was purged and back filled with nitrogen gas. To the flask, toluene was added to dissolve **15**. Once fully dissolved, a 10-fold excess of HSiCl₃ was added to the reaction mixture. Upon addition of the silane species, the solution changed colour from a bright yellow to a dull beige colour as well as the appearance of an oily substance at the bottom of the reaction flask. This was stirred at room temperature for

24 hours under a constant stream of nitrogen gas (**Scheme 43**). It was noted that after the reaction was finished the mixture had a distinct odour, like that of COD. After completion, excess volatile components were removed *in vacuo*. Without further purification, the complex was dissolved in a deuterated solvent to check its progress. The beige powder which was left behind had extreme limited solubility in both CDCl_3 and C_6D_6 making analysis by NMR spectroscopy tough. An overnight ^{29}Si -NMR experiment was performed (14 hours) which revealed no silicon signals in the sample. Due to time constraints, further reactions of **15** with H-Si reagents were abandoned.

Scheme 43: Attempted oxidative addition to **15** to make **17**.

The final oxidative addition reaction attempted was the addition of HSnPh_3 to **15**. Triphenylstannane, a white material (^{119}Sn $\delta = -163$ ppm) was mixed with **15** and 1.0 ml C_6D_6 in the glovebox in a small flask. The flask, brought out of the glovebox, was filled with nitrogen gas and left undisturbed (stirring) for 24 hours under the $\text{N}_{2(\text{g})}$ atmosphere. Upon completion of the reaction the solution was concentrated further under vacuum. To this, dry *n*-hexanes was added which lead to immediate precipitation of a bright yellow powder. The powder was filtered, washed with cold Et_2O and dried (**Scheme 44**). The compound was analyzed by ^{119}Sn -NMR spectroscopy to confirm purity. The ^{119}Sn -NMR spectrum displayed one tin resonance signal at $\delta_{\text{Sn}} = -178$ ppm (**Figure 45**). This value is comparable to other Sn-Ir-H complexes.^{54,80}

Scheme 44: Synthesis and oxidative addition of $\text{H}(\text{Ph})_3\text{Sn}$ to **15** to make **18**.

Figure 45: A- ^{119}Sn -NMR spectrum of **18**, B- ^1H -NMR spectrum of **18**

In the IR spectrum, there is a notable shift in the frequency of C-O (693 cm^{-1}) and less drastic shift in the C=N/C=C (1537 cm^{-1}) stretching frequency. This is expected as **18** now has more electron withdrawing character around the center, making the resting state of the overall compound more neutral.

Assignments of **18** were made by a combination of 1D and 2D NMR experiments (^1H , ^{13}C , ^{119}Sn , COSY, HSQC, nOesy). By looking at the 1D NMR spectra information like coupling can be deduced, not only ^1H - ^1H coupling but also ^1H - ^{119}Sn and ^{13}C - ^{119}Sn . As was in the case of **16**, **18**

also experiences desymmetrization upon addition of HSnPh_3 , seeing the protons of the oxazoline ring being split in the same fashion (doublet for methyl groups, and an ABq for the methylene protons) as well as each 'COD' proton and carbon now giving their own resonance signal in each 1D NMR spectrum. Perhaps more interesting is the appearance of the lone hydride signal, resonating at ^1H δ -17.40 ppm. To this signal are distinct ^1H - ^{119}Sn coupling satellites giving a coupling constant of $^2J_{\text{cis}} = 23.0$ Hz, indicating they are no longer bonded together but are instead being separated by another atom (presumably **Ir**).⁸⁰ This behaviour is also observed in the ^{13}C NMR spectrum, in which all the *ortho* carbons in the $-\text{SnPh}_3$ group can be seen coupling with a coupling constant of $^2J_{\text{Sn-C}} = 17.0$ Hz. The interactions of the hydride can be monitored by nOesy NMR. The nOesy of **18** shows interactions of the hydride with both the methyl/methylene groups on the oxazoline and the rest of $-\text{SnPh}_3$. Geometry proposed for **18** is where the hydride and Sn atoms are *cis* to one another, supported by nOesy-NMR spectroscopy (**Figure 46**). This notion was later confirmed by x-ray crystallography, indefinitely proving a *cis*-addition of triphenylstannane to **15** (**Figure 47**). X-ray crystallography quality crystals were grown by slow diffusion of dry n-hexanes into THF at -30 °C for 8 days. **18** crystallized as yellow cubes in the monoclinic crystal system within in the $\text{P2}_1/\text{n}$ space group. This is ascertained by the unit cell parameters. Bond angles and **Ir**-O and **Ir**-N bond lengths as well as **Ir**-H and **Ir**-Sn bond lengths are comparable to other Ir^{+3} -COD complexes (**Table 14**).^{55,79,80,83–86}

Figure 46: nOesy NMR spectrum of **18** (red box highlights NOE interactions)

Figure 47: ORTEP representation of **18**.

Table 14: Select bond lengths (Å) and angles (°) for **18**.

Bond Lengths (Å)		Bond Angles (°)		Bond Angles (°)	
Ir(1)-N(1)	2.095(2)	N(1)-Ir(1)-O(1)	87.95(8)	N(1)-Ir(1)-Sn(1)	93.55(6)
Ir(1)-O(1)	2.1592(19)	N(1)-Ir(1)-C(18)	165.61(10)	O(1)-Ir(1)-Sn(1)	86.42(5)
Ir(1)-C(18)	2.166(2)	O(1)-Ir(1)-C(18)	78.79(9)	C(18)-Ir(1)-Sn(1)	91.19(7)
Ir(1)-C(19)	2.180(3)	N(1)-Ir(1)-C(19)	155.50(10)	C(19)-Ir(1)-Sn(1)	92.31(7)
Ir(1)-C(14)	2.294(3)	O(1)-Ir(1)-C(19)	116.16(9)	C(14)-Ir(1)-Sn(1)	159.20(8)
Ir(1)-C(15)	2.310(3)	C(18)-Ir(1)-C(19)	37.39(10)	C(15)-Ir(1)-Sn(1)	165.09(8)
Ir(1)-Sn(1)	2.6137(2)	N(1)-Ir(1)-C(14)	86.95(9)	N(1)-Ir(1)-H(1IR)	84.6(13)
Ir(1)-H(1IR)	1.37(3)	O(1)-Ir(1)-C(14)	114.38(9)	O(1)-Ir(1)-H(1IR)	165.3(13)
		C(18)-Ir(1)-C(14)	93.41(10)	C(18)-Ir(1)-H(1IR)	109.6(13)
		C(19)-Ir(1)-C(14)	79.33(10)	C(19)-Ir(1)-H(1IR)	72.8(13)
		N(1)-Ir(1)-C(15)	92.83(9)	C(14)-Ir(1)-H(1IR)	78.0(13)
		O(1)-Ir(1)-C(15)	80.36(10)	C(15)-Ir(1)-H(1IR)	112.6(13)
		C(18)-Ir(1)-C(15)	79.56(10)	Sn(1)-Ir(1)-H(1IR)	81.4(13)
		C(19)-Ir(1)-C(15)	87.41(11)	O(1)-Ir(1)-N(1)	91.57(8)

2.7.3.2 Attempted displacement of the COD ring of complex **15** (**19a-e**)

Attempts were also made to displace the ‘COD’ ligand of **15** with another either mono- or di-chelating nitrogen or phosphorus based nucleophiles. A series of displacement reactions were carried out on **15** in attempt to yield a series of novel Ir catalysts. Displacement of the ‘COD’ ring with TMEDA, pyridine, dppf, PPh₃, and P(OEt)₃ were attempted (**Scheme 45**). Unfortunately, each case gave either quantitative recovery of starting materials or an inseparable mixture of by-products.

Scheme 45: Various attempted displacement reactions with P/N nucleophiles with **15**.

3.0 Conclusions and Future directions:

A series of 1,5-N,O-bidentate ligands (**4a-l/6a-c**) and their respective copper (11 compounds, **7/8**) and cobalt (19 compounds, **9-11**) complexes have been successfully synthesized. With regards to heavier group 9 complexes, the synthesis of three novel iridium compounds were synthesized with ligand **4a**. The explored synthetic pathways showcased the ease of tuning both the electronic and structural properties of the oxazoline-based ligand. Recrystallization from simple organic solvents gave relatively modest yields ranging from 23-99 % for the ligands (**4/6**), 32-99 % for the copper and cobalt series (**7-11**), and yields of 42-75 % for the iridium containing compounds (**15-18**). All organic compounds were characterized by ^1H -, ^{13}C -NMR, their melting point and EA (where applicable). All transition metal containing complexes were characterized by IR, UV-Vis, EA, MS, CV, their melting point, and x-ray crystallography where appropriate. With the characterization obtained, mainly the reversible nature of its oxidation displayed in the cyclic voltammetry, only **9h** was deemed a viable candidate for catalysis and thus was trial in the ATRP of MMA. **9h** gave PMMA in 42 % conversion with a PDI of 1.19 and molecular weight of ~ 50,000 Da. Preliminary coordination studies have been carried out with **Ir** and **Rh** centers giving compound **15** in a decent yield (75 %). **15** underwent oxidation addition reactions with MeI and HSnPh_3 . Displacement reactions were undertaken of **15** in attempt to replace the 'COD' ring with another ligand although all attempts were unsuccessful.

Future directions for this project would be to perfect the ATRP of MMA by changing aspects such as initiator, time, temperature and solvent. Once optimized, different monomers (such as styrene) could be polymerized. With regards to heavy group 9 complexes, the continued attempted synthesis of 'COD' displaced compounds should be tried with other nucleophiles

ligands. To complex **15**, a good continuation of its oxidative properties could be the addition of HSiPh_3 , which has been proved to work more effectively over HSiCl_3 in these types of reactions.

4.0 Experimental

4.1 General Considerations

All reagents, solvents and deuterated solvents were obtained from commercially available sources. Dry solvents were dispensed from an Mbraun solvent purification system immediately before use. Thin layer chromatography (TLC) was performed on aluminum backed sheets pre-coated with silica F254 (0.25 nm thick; Silicycle, Quebec, Canada) adsorbent eluting with the specified solvent system used. All preparative thin layer chromatography was performed on silica plated glass-backed plates. All TLC results were observed by either UV irradiation or iodine vapour staining. Column chromatography was carried out on 250-400 mesh size silica gel, eluting with the specified solvent system for each listed compound. Nuclear magnetic resonance spectroscopy (^1H NMR = 400 MHz, $^{13}\text{C}\{^1\text{H}\}$ NMR = 101 MHz, $^{31}\text{P}\{^1\text{H}\}$ -NMR = 161 MHz, $^{29}\text{Si}\{^1\text{H}\}$ -NMR = 79 MHz, $^{119}\text{Sn}\{^1\text{H}\}$ -NMR = 149 MHz) was completed on Bruker Avance II AC-400 MHz spectrometer at 300K in the specified deuterated solvent for each compound. Signals were picked and processed based on solvent residual ^1H and ^{13}C peaks. Proton and carbon assignments were made using a variety of 2D-NMR experiments (COSY, HSQC, HMBC, nOesy). Abbreviations used to indicate multiplicity are as followed: s = singlet, d = doublet, t = triplet, m = multiplet, dd = doublet of doublets, td = triplet of doublets, ABq = AB quartet. Infrared spectroscopy was performed on an Agilent Cary 630 -FTIR Spectrometer in open air and have been left uncorrected. Cyclic voltammetry data was collected using a MetrOhm μ -Autolab Type III potentiostat/galvanostat. All solvents (DCM) were degassed prior to analysis by cyclic voltammetry. UV-Vis absorption profiles were collected using an Agilent Cary 5000 UV-vis-NIR spectrophotometer with a cuvette with a path length of 1 cm was used. Molar extinction coefficients were calculated using the Beer-Lambert law. Melting points were determined in open air and in triplicate and have been left

uncorrected. The purity of all the products was confirmed by a combination of melting point, X-ray and NMR spectroscopies. The growth of all metal complex single crystals was carried out in either a 1:1 mixture of DCM/MeOH or by slow diffusion of Et₂O into CHCl₃, whereas crystal growth of all organic compounds was carried out by slow evaporation of neat acetone. A Bruker-Nonius Kappa-CCD diffractometer was used to collect the X-ray information crystal. This machine is equipped with monochromated Mo-K α radiation source. All x-ray crystallographic data sets were obtained and solved by Dr. Alan J. Lough and the University of Toronto. Molecular weights of polymers were determined by triple detection gel permeation chromatography (TDGPC) equipped with a Refractive Index Detector (RI), a four-capillary differential viscometer (VISC), a right-angle laser light scattering detector and a low angle laser light scattering detector. GPC columns were calibrated against polystyrene standards. HPLC grade THF was used as the mobile phase with a constant flow rate of 1 mL/min. GPC samples were made at a concentration of 5 mg/mL and diluted as per the GPC experimental method. Elemental Analysis was performed by Atlantic Microlabs in Norcross, Georgia. Toluene was used *in lieu* of benzene for the purposes of re-crystallization for all organic compounds (intermediates [3/5] and ligands [4/6]). The analyses of synthesized compounds which have been previously reported in the literature are comparable to their referenced literature sources. For these compounds only new data is reported (IR/NMR spectral data not included), or have information reported for comparison with data gathered in this thesis.

4.2 Synthesis and Characterization of Intermediates species (**3a-h**)

4.2.1 Synthesis of (*E*)-2-(3-benzoyl-4,4-dimethyloxazolidin-2-ylidene)-1-phenylethan-1-one (**3a**)

At room temperature **1a** (5.0 mL, 39.2 mmol), NEt₃ (14.2 mL, 101.9 mmol) and ACN (50 mL) was added to a 250-mL round-bottom flask. After letting the solution cool to 0°C, benzoyl chloride (10.0 mL, 86.2 mmol) was added dropwise *via* an addition funnel. The suspension was refluxed for 3 hours. Upon completion of the reaction rotary evaporator was employed to remove the solvent. The solids were diluted with dH₂O (50 mL) and washed with DCM (2 × 50 mL). The organic layer was subjected to a second wash with a 10% solution of Na₂CO_{3(aq)} (3 × 100 mL). Excess solvents were removed by rotary evaporator leaving behind a sludgy orange mixture. This mixture was then recrystallized with a 2:1 hexane-toluene blend to reveal the product as a pale yellow solid (10.61 g, 84 % yield). Analysis agrees with the literature data for this compound.^{20,35}

Melting Point Range: 115-118 °C (lit: 119-120 °C)

TLC: 45% EtOAc/*n*-hexanes, R_f = 0.13

4.2.2 Synthesis of (Z)-2-(3-(3,4-dichlorobenzoyl)-4,4-dimethyloxazolidin-2-ylidene)-1-(3,4-dichlorophenyl)ethan-1-one (**3b**)

A flask was charged with **1a** (3.00 mL, 23.5 mmol), THF (50 mL), and NEt₃ (10.0 mL, 71.7 mmol). After allowing the solution to cool to 0°C, 3,4-dichlorobenzoyl chloride (10.9 g, 52.1 mmol) in THF (25 mL) was added *via* addition funnel. The mixture was set to reflux for 3 hours. The solution was cooled to room temperature and filtered to remove the integrated salt. Following filtration, the volatile components were removed *via* rotary evaporator. To the resulting pale yellow solid, dH₂O (75 mL) was added then the solution was washed with CHCl₃ (75 mL). The organic components were washed one last time with a 10 % solution of Na₂CO_{3(aq)} (1 × 100 mL). Sodium sulphate was added, filtered and the volatile constituents were removed *via* rotary evaporator. The resulting white powder was recrystallized with a 1:2 mixture of toluene-hexane. Filtration revealed a pale yellow fluffy solid (6.7 g, 63 % yield).

Melting Point Range: 79-82°C

TLC: 20% EtOAc/*n*-hexanes, R_f = 0.16

¹H-NMR: (400 MHz, CDCl₃) δ **1.66** (s, 6H, **1**), **4.38** (s, 2H, **3**), **4.93** (s, 1H, **5**), **7.21-7.23** (m, 1H, Ar-**H**), **7.36-7.38** (m, 1H, Ar-**H**), **7.54-7.65** (m, 3H, Ar-**H**), **7.85** (m, 1H, Ar-**H**)

¹³C-NMR: (101 MHz, CDCl₃) δ 22.8 (**1**); 63.4 (**2**); 79.9 (**5**); 85.2 (**3**); 126.2 (**Ar-H**); 128.1 (**Ar-H**); 129.2 (**Ar-H**); 130.3 (**Ar-H**); 130.9 (**Ar-H**); 131.1 (**Ar-H**); 132.6 (**Ar-X**); 133.8 (**Ar-X**); 133.9 (**Ar-X**); 135.8 (**Ar-X**); 137.8 (**Ar-X**); 139.6 (**Ar-X**); 161.5 (**6'**); 166.2 (**4**); 184.1 (**6**)

4.2.3 (*E*)-2-(3-(4-methoxybenzoyl)-4,4-dimethyloxazolidin-2-ylidene)-1-(4-methoxyphenyl)ethan-1-one (**3c**)

A 150-mL flask was charged with *p*-methoxybenzoic acid (3.10 g, 20.4 mmol) and SOCl₂ (50.0 mL, 0.689 mol) and was refluxed (90 °C) for 2 hours. Volatile components were removed *in vacuo*. The acid chloride (3.48 g, 20.4 mmol) was not isolated and immediately used. After cooling to 0 °C, **1a** (1.18 mL, 9.27 mmol), NEt₃ (3.88 mL, 27.81 mmol) and THF (50 mL) were added to the flask. The mixture was refluxed for 3 hours. Upon completion, the resultant salt was filtered off and the volatiles were removed *via* rotary evaporator. DCM (50 mL) was added and the solution was washed with dH₂O (50 mL × 1). The organics were washed one final time with a solution of 10% Na₂CO_{3(aq)} (30 mL) and brine (50 mL). Sodium sulphate was added, filtered out, and the volatile components removed *via* rotary evaporator. The resulting orange sludge was obtained, although not isolated. Recrystallization was carried out with a mixture of 1:2 toluene-hexane, giving a yellow powder (1.258 g, 35 % yield). Analyses are consistent with those in the literature.^{20,87}

Melting Point Range: 174-176 °C (lit: 175-176 °C)

TLC: 70 % EtOAc/Hexanes R_f = 0.09

4.2.4 Synthesis of (Z)-2-(3-(3,4-dimethylbenzoyl)-4,4-dimethyloxazolidin-2-ylidene)-1-(3,4-dimethylphenyl)ethan-1-one (**3d**)

A 150-mL flask was charged with 3,4-dimethylbenzoic acid (3.10 g, 20.4 mmol), DMF (0.10 mL, 99 mg, 1.4 mmol), SOCl₂ (50.0 mL, 82.0 g, 0.689 mol) and was refluxed for 2 hours. Volatile components were removed *in vacuo*. The acid chloride (3.48 g, 20.4 mmol) was not isolated and immediately used. After cooling to 0 °C, **1a** (1.18 mL, 1.05 g, 9.27 mmol), NEt₃ (3.88 mL, 2.82 g, 27.8 mmol) and THF (50 mL) were added to the flask. The mixture was refluxed for 3 hours. Upon completion, the resultant salt was filtered off and the volatiles were removed *via* a rotary evaporator. The solids were diluted with DCM (50 mL) and washed with dH₂O (50 mL × 1). Organics were washed one final time with a solution of 10% Na₂CO_{3(aq)} (30 mL) and brine (50 mL). The organic phase was dried (Na₂SO₄), filtered, and volatile components removed *via* a rotary evaporator. The orange oil was recrystallized with a mixture of 1:2 toluene/*n*-hexanes, giving a yellow powder (1.3 g, 35 % yield).

Melting Point Range: 122-124°C

TLC: 20% EtOAc/*n*-hexanes, R_f = 0.11

¹H-NMR: (400 MHz, CDCl₃) δ **1.68** (s, 6H, **1**), **2.17** (s, 3H, Ar-CH₃), **2.24** (s, 3H, Ar-CH₃), **2.34** (s, 3H, Ar-CH₃), **2.38** (s, 3H, Ar-CH₃), **4.37** (s, 2H, **3**), **5.12** (s, 1H, **5**), **7.00-7.02** (m, 2H, **Ar**), **7.11** (dd, *J* = 1.5, 4.8 Hz, 1H, **Ar**), **7.29** (d, *J* = 8.0 Hz, 1H, **Ar**), **7.49-7.53** (m, 2H, **Ar**).

¹³C-NMR: (101 MHz, CDCl₃) δ 19.7 (**Ar-Me**); 19.7 (**Ar-Me**); 19.8 (**Ar-Me**); 20.1 (**Ar-Me**); 22.8 (**1**); 62.6 (**2**); 79.7 (**5**); 85.7 (**3**); 124.9 (**Ar**); 126.8 (**Ar**); 128.5 (**Ar**); 129.3 (**Ar**); 130.1 (**Ar**); 130.1 (**Ar**); 132.4 (**Ar**); 136.0 (**Ar**); 137.4 (**Ar**); 138.1 (**Ar**); 140.3 (**Ar**); 142.1 (**Ar**); 161.2 (**6'**); 169.1 (**4**); 186.8 (**6**).

EA: (%): calc'd. (found) for C₂₄H₂₇NO₃: C 73.73 (73.93); H 7.33 (7.33); N 3.58 (3.61).

4.2.5 Synthesis of (Z)-2-(3-(4-chloro-3-methylbenzoyl)-4,4-dimethyloxazolidin-2-ylidene)-1-(4-chloro-3-methylphenyl)ethan-1-one (**3e**)

A flask was charged with 4-chloro-3-methylbenzoic acid (5.01 g, 29.4 mmol), DMF (0.10 mL, 99 mg, 1.4 mmol) and SOCl₂ (10.0 mL, 16.4 g, 0.138 mol) and was refluxed for 1 hour. Volatile components were removed in *vacuo*. **1a** (1.70 mL, 1.51 g, 13.3 mmol) and THF (20 mL) were added and cooled to 0 °C. The acid chloride (in THF, 20 mL) was added dropwise to **1a**-containing flask. NEt₃ (5.90 mL, 4.38 g, 42.3 mmol) was added to the flask. The mixture was refluxed for 3 hours. The resultant salt was filtered off and the volatiles were removed *via* rotary evaporator. The solids were diluted with DCM (50 mL) and washed with dH₂O (50 mL × 2). The organics were washed with a solution of 10% Na₂CO_{3(aq)} (1 × 70 mL). Organics were dried (MgSO₄), filtered and the volatile constituents were removed *via* a rotary evaporator. The target product was recrystallized with a 1:1 mixture of toluene and *n*-hexanes and was isolated as an off-yellow-coloured powder (4.8 g, 86% yield).

Melting Point Range: 154-156°C

TLC: 45% EtOAc/*n*-hexanes, R_f = 0.19

¹H-NMR: (400 MHz, CDCl₃) δ **1.62** (s, 6H, **1**), **2.30** (s, 3H, **13**), **2.46** (s, 3H, **13'**), **4.33** (s, 2H, **3**), **4.95** (s, 1H, **5**), **7.04** (d, *J* = 1.1 Hz, 1H, **8'**), **7.09** (d, *J* = 7.9 Hz, 1H, **12'**), **7.23** (dd, *J* = 8.0, 1.1 Hz, 1H, **11'**), **7.36** (d, *J* = 7.8 Hz, 1H, **8**), **7.49** (dd, *J* = 7.8, 1.1 Hz, 1H, **12**), **7.70** (d, *J* = 1.1 Hz, 1H, **11**)

¹³C-NMR: (101 MHz, CDCl₃) δ 20.1 (**13**); 20.5 (**13'**); 22.8 (**1**); 63.0 (**2**); 79.8 (**5**); 85.4 (**3**); 125.5 (**11'**); 127.2 (**11**); 127.8 (**8'**); 129.6 (**8**); 130.7 (**12'**); 131.4 (**12**); 133.5 (**Ar**); 134.0 (**Ar**); 135.2 (**Ar**); 139.4 (**Ar**); 139.6 (**Ar**); 141.8 (**Ar**); 161.4 (**6'**); 167.3 (**4**); 185.1 (**6**)

EA: (%): calc'd. (found) for C₂₂H₂₁Cl₂NO₃: C 61.19 (61.47); H 5.25 (5.34); N 3.24 (3.42).

4.2.6 Synthesis of (Z)-2-(3-(3-chloro-4-methylbenzoyl)-4,4-dimethyloxazolidin-2-ylidene)-1-(3-chloro-4-methylphenyl)ethan-1-one (**3f**)

A flask was charged with 3-chloro-4-methylbenzoic acid (5.078 g, 29.8 mmol), DMF (0.10 mL, 99 mg, 1.4 mmol) and SOCl_2 (10.0 mL, 16.4 g, 0.138 mol) and was refluxed for 1 hour. Volatile components were removed *in vacuo*. **1a** (1.70 mL, 1.51 g, 13.3 mmol) and THF (20 mL) were added and was cooled to 0 °C. The acid chloride (5.63 g, 29.8 mmol) was dissolved in THF (20 mL) and added dropwise **1a**-containing flask. NEt_3 (5.90 mL, 4.50 g, 42.3 mmol) was added to the flask. The mixture was refluxed at for 3 hours. The resultant salt was filtered off and the volatiles were removed *via* a rotary evaporator. The solids were diluted with DCM (50 mL) and washed with dH_2O (50 mL $\times$ 2). The organics were washed with a solution of 10% $\text{Na}_2\text{CO}_{3(\text{aq})}$ (1 $\times$ 70 mL). The solution was dried (MgSO_4), filtered and the volatile constituents were removed *via* a rotary evaporator. The target product was recrystallized with a 1:1 mixture of toluene and *n*-hexanes then was isolated as a pale-yellow-coloured powder (3.8 g, 68% yield).

Melting Point Range: 142-145°C

TLC: 20% EtOAc/*n*-hexanes, R_f = 0.53

$^1\text{H-NMR}$: (400 MHz, CDCl_3) δ **1.65** (s, 6H, **1**), **2.28** (s, 3H, **13**), **2.43** (s, 3H, **13'**), **4.35** (s, 2H, **3**), **4.97** (s, 1H, **5**), **7.03** (d, J = 1.5 Hz, **8'**), **7.09** (dd, J = 8.2, 1.5 Hz, **12'**), **7.22** (d, J = 8.2 Hz, **11'**), **7.47-7.52** (m, 2H, **8,12**), **7.60** (s, 1H, **11**)

$^{13}\text{C-NMR}$: (101 MHz, CDCl_3) δ 20.0 (**13**); 20.0 (**13'**); 22.8 (**1**); 79.8 (**2**); 85.6 (**5**); 126.1 (**Ar-H**); 127.2 (**Ar-H**); 127.9 (**Ar-H**); 129.0 (**Ar-H**); 129.7 (**Ar-H**); 129.8 (**Ar-H**); 131.4 (**Ar**); 135.9 (**Ar**); 137.4 (**Ar**); 138.7 (**Ar**); 139.4 (**Ar**); 141.6 (**Ar**); 161.3 (**6'**); 167.8 (**4**); 185.7 (**6'**)

EA: (%): calc'd. (found) for $\text{C}_{22}\text{H}_{21}\text{Cl}_2\text{NO}_3$: C 61.19 (61.24); H 5.25 (5.46); N 3.24 (3.32).

4.2.7 (Z)-2-(2-(2-hydroxy-2-(naphthalen-1-yl)vinyl)-4,4-dimethyl-4,5-dihydro-3H-oxazol-3-yl)(naphthalen-1-yl)methanone (**3g**)

A flask was charged with **1a** (5.0 mL, 39.19 mmol) and THF (25 mL). After allowing the solution to cool to 0°C, 1-naphthoyl chloride (13.00 g, 86.22 mmol) was added to the mixture. Once added, the solution was warmed to room temperature where NEt₃ (16.39 mL, 117.58 mmol) was then added. The mixture was set to reflux for 3 hours. Upon completion of the allotted time, the solution was cooled to room temperature and filtered to remove the integrated salt. Following filtration, the volatile components were removed *via* rotary evaporator. To the resulting pale yellow solid, dH₂O (50 mL) was added then the solution was washed with DCM (50 mL). The organic components were washed one last time with a 10 % solution of Na₂CO_{3(aq)} (1 × 60 mL). Sodium sulphate was added, filtered and the volatile constituents were removed *via* rotary evaporator. The resulting yellow powder was recrystallized with a 1:1 mixture of toluene-hexane. Filtration revealed a pale yellow fluffy solid (12.611 g, 76 % yield). EA of this compound is pending.

Melting Point Range: 140-143°C

TLC: 20% EtOAc/*n*-hexanes, R_f = 0.34

4.2.8 (E)-1-(4,4-dimethyl-3-pivaloyloxazolidin-2-ylidene)-3,3-dimethylbutan-2-one. (**3h**)^{20,35}

A flask was charged with **1a** (2.00 mL, 15.7 mmol), NEt₃ (7.00 mL, 50.2 mmol) and ACN (15 mL). Trimethylacetyl chloride (4.20 mL, 34.1 mmol) as added to the mixture. The mixture was set to reflux for 3 hours. Upon completed of the allotted time, the solution was cooled to room temperature and the volatile components were removed *via* rotary evaporator. To the resulting pale yellow solids, dH₂O (50 mL) was added then the solution was washed with CHCl₃ (50 mL). The organic components were washed one last time with a 10 % solution of Na₂CO_{3(aq)} (1 × 70 mL). Magnesium sulphate was added, filtered and the volatile constituents were removed *via* rotary evaporator. The resulting yellow oil was used crude and determined pure *via* NMR (3.849g, 87 % yield). Analyses of this compound are consistent with those in the literature.^{20,35}

Melting Point Range: 75-77 °C (lit: 75-75.5 °C)

TLC: 35 % EtOAc/Hexanes R_f = 0.28

¹H-NMR: (400 MHz, CDCl₃) δ **1.09** (s, 9H, *t*-Bu-**H** **8**), **1.23** (s, 6H, **1**), **1.27** (s, 9H, **8'**), **3.83** (s, 2H, **3**), **5.76** (s, 1H, **5**)

¹³C-NMR: (101 MHz, CDCl₃) δ 27.2 (**8'**); 27.6 (**8**); 28.2 (**1**); 37.4 (**7'**); 39.2 (**7**); 66.3 (**2**); 78.5 (**5**); 102.1 (**3**); 159.7 (**6'**); 164. 1 (**4**); 174.8 (**6**)

4.3 Synthesis and Characterization for the Ligand Series made with **1a**

4.3.1 Synthesis of (Z)-2-(4,4-dimethyl-4,5-dihydrooxazol-2-yl)-1-phenylethen-1-ol (**4a**)

In a 250-mL round-bottom flask, KOH (4.233g, 75.5 mmol, 1.5 M) was dissolved in MeOH (50 mL) at room temperature. After fully dissolved, compound **3a** (5.04g, 15.6 mmol) was added to the mixture and was stirred for 6 hours at room temperature. Any solids were filtered off the solution. The filtrate was washed twice with DCM (2×25 mL). The organic layer was washed again with dH₂O (3×25 mL) and once with a solution of brine (20 mL). Removal of solvents was done by rotary evaporator after the addition of MgSO₄ and filtration. Evaporation of excess solvents yielded a pale-yellow powder (4.22 g, 64 % yield). Analyses of this compound are consistent with those in the literature.^{20,35}

Melting Point Range: 99-101 °C (lit: 100.5-101 °C)

TLC: 45 % EtOAc/*n*-hexanes $R_f = 0.47$

4.3.2 (*E*)-1-(3,4-dichlorophenyl)-2-(4,4-dimethyloxazolidin-2-ylidene)ethan-1-one (**4b**)

In a 50-mL round-bottom flask, KOH (2.14g, 37.5 mmol, 1.5 M) was dissolved in MeOH (25 mL) at room temperature. After fully dissolved, compound **3b** (1.04g, 2.19 mmol) was added to the mixture and was stirred for 19 hours at room temperature. Volatile components were removed by rotary evaporator. The resulting white solid was diluted with dH₂O (25 mL) and washed with DCM (1 × 25 mL). Organic constituents were washed once more with dH₂O (50 mL). Sodium sulphate was added, filtered and any volatiles removed *via* rotary evaporator. The target product was collected as a white, needle-like solid (532 mg, 84% yield).

Melting Point Range: 119-121°C

TLC: 20% EtOAc/*n*-hexanes, *R_f* = 0.14

¹H-NMR: (400 MHz, CDCl₃) δ **1.43** (s, 6H, **1**), **4.16** (s, 2H, **3**), **5.46** (s, 1H, **6**), **7.44** (d, *J* = 8.25 Hz, 1H, **12**), **7.65** (dd, *J* = 1.75, 8.25 Hz, 1H, **13**), **7.92** (d, *J* = 1.75 Hz, 1H, **9**), **9.91** (s, 1H, **4**)

¹³C-NMR: (101MHz, CDCl₃) δ 27.1 (**1**); 58.7 (**2**); 73.9 (**6**); 79.1 (**3**); 126.0 (**13**); 128.9 (**9**); 130.1 (**12**); 132.4 (**10**); 134.4 (**8**); 139.9 (**11**); 169.8 (**5**); 184.5 (**7**)

EA: (%): calc'd. (found) for C₁₃H₁₃Cl₂NO₂: C 54.57 (54.50); H 4.58 (4.51); N 4.89 (4.85).

MS: (m/z): calc'd (found): [M⁺] = 286.12 (286.1)

4.3.3 Synthesis of (*E*)-2-(4,4-dimethyloxazolidin-2-ylidene)-1-(4-methoxyphenyl)ethan-1-one (**4c**)

In a 100-mL round-bottom flask, KOH (4.25g, 75.7 mmol, 1.51 M) was dissolved in MeOH (50 mL) at room temperature. After fully dissolved, compound **3c** (1.26g, 3.30 mmol) was added to the mixture and was stirred for 9 hours at room temperature. Any solids were filtered off the solution. Volatile components were removed by rotary evaporator. The resulting white solid was diluted with dH₂O (50 mL) and washed with DCM (1 × 50 mL). Organic constituents were washed once more with brine (50 mL). Magnesium sulphate was added, filtered and any volatiles removed *via* rotary evaporator. The target product was collected as an orange-yellow solid (401 mg, 49 % yield). Analyses of this compound are consistent with those in the literature.^{20,88}

Melting Point Range: 104-106°C (lit: 105-106 °C)

TLC: 45% EtOAc/*n*-hexanes, R_f = 0.19

4.3.4 Synthesis of (*E*)-2-(4,4-dimethyloxazolidin-2-ylidene)-1-(3,4-dimethylphenyl)ethan-1-one (**4d**)

In a 100-mL round-bottom flask, KOH (2.15g, 38.3 mmol, 1.53 M) was dissolved in MeOH (25.0 mL) at room temperature. After fully dissolved, compound **3d** (0.843g, 2.23 mmol) was added to the mixture and was stirred for 20 hours at room temperature. Any solids were filtered off the solution. Volatile components were removed by a rotary evaporator. The resulting solids were diluted with dH₂O (50 mL) and washed with CHCl₃ (1 × 50 mL). Organic constituents were washed once more with dH₂O (1 × 50 mL). The organic phase was dried (MgSO₄) filtered and volatiles removed *via* a rotary evaporator. The target product was collected as a pale yellow solid (0.300 g, 56 % yield).

Melting Point Range: 149-151°C

TLC: 35% EtOAc/Hexanes R_f = 0.18

¹H-NMR: (400 MHz, CDCl₃) δ **1.42** (s, 6H, **1**), **2.28** (d, *J* = 5.8 Hz, 6H, **14-15**), **4.12** (s, 2H, **3**), **5.56** (s, 1H, **6**), **7.14** (d, *J* = 7.8 Hz, 1H, **12**), **7.59** (d, *J* = 7.8 Hz, 1H, **13**), **7.64** (s, 1H, **9**), **9.91** (s, 1H, **4**)

¹³C-NMR: (101MHz, CDCl₃) δ 19.8 (**14**); 19.9 (**15**); 27.1 (**1**); 58.5 (**2**); 73.8 (**6**); 73.9 (**3**); 124.4 (**13**); 128.1 (**9**); 129.4 (**12**); 136.2; 137.6; 139.5; 169.5 (**5**); 187.7 (**7**)

EA: (%): calc'd. (found) for C₁₅H₁₉NO₂: C 73.44 (71.99); H 7.81 (7.64); N 5.71 (5.39).

- 0.5 Acetone was found in the lattice in elemental analysis

MS: (m/z): calc'd (found): [M⁺+H] = 245.3 (246.4)

4.3.5 Synthesis of (*E*)-1-(4-chloro-3-methylphenyl)-2-(4,4-dimethyloxazolidin-2-ylidene)ethan-1-one (**4e**)

In a 150-mL beaker, KOH (2.20g, 39.2 mmol, 1.57 M) was dissolved in MeOH (25.0 mL) at room temperature. The KOH/MeOH mixture was added to a 250-mL flask containing **3e** (1.51g, 3.60 mmol) was added to the mixture and was stirred for 6 hours at room temperature. Volatile components were removed by a rotary evaporator. The resulting solids were diluted with dH₂O (50 mL) and washed with DCM (1 × 50 mL). Organic constituents were washed once more with dH₂O (1 × 100 mL). Magnesium sulphate was added to dry the solution. The mixture was then filtered and the volatile constituents were removed *via* a rotary evaporator. The orange oil was recrystallized with a mixture containing 1:4 toluene/*n*-hexanes. The target product was collected as a pale yellow-coloured solid (0.588 g, 69 % yield).

Melting Point Range: 127-129°C

TLC: 45% EtOAc/*n*-hexanes, *R_f* = 0.30

¹H-NMR: (400 MHz, CDCl₃) δ **1.42** (s, 6H, **1**), **2.37** (s, 3H, **14**), **4.14** (s, 2H, **3**), **5.49** (s, 1H, **6**), **7.22** (d, *J* = 7.9 Hz, 1H, **12**), **7.61** (dd, *J* = 7.9, 1.7 Hz, 1H, **13**), **7.81** (d, *J* = 1.7 Hz, 1H, **9**), **9.79** (s, 1H, **4**)

¹³C-NMR: 20.2 (**14**); 27.2 (**1**); 58.7 (**2**); 73.9 (**6**); 79.1 (**3**); 125.1 (**9**); 127.7 (**13**); 130.8 (**12**); 134.4 (**10**); 138.6 (**8**), 139.4 (**11**); 169.8 (**5**); 185.9 (**7**)

EA: (%): calc'd. (found) for C₁₄H₁₆ClNO₂: C 63.28 (63.47); H 6.07 (5.97); N 5.27 (5.17).

MS: (*m/z*): calc'd (found): [M⁺+H] = 265.7 (266.1)

4.3.6 Synthesis of (*E*)-1-(3-chloro-4-methylphenyl)-2-(4,4-dimethyloxazolidin-2-ylidene)ethan-1-one (**4f**)

In a 100-mL round-bottom flask, KOH (4.53 g, 80.5 mmol, 1.61 M) was dissolved in MeOH (50.0 mL) at room temperature. **3f** (2.01 g, 4.80 mmol) was added to the mixture and was stirred for 16 hours at room temperature. Volatile components were removed by a rotary evaporator. The solid was diluted with dH₂O (50 mL) and washed with DCM (2 × 50 mL). Organic constituents were washed once more with dH₂O (1 × 100 mL). Magnesium sulphate was added to dry the solution. The mixture was then filtered and the volatile constituents were removed *via* a rotary evaporator. The target product was collected as a pinkish-coloured solid (0.62 g, 49 % yield).

Melting Point Range: 143-145°C

TLC: 45% EtOAc/*n*-hexanes, *R_f* = 0.28

¹H-NMR: (400 MHz, CDCl₃) δ **1.41** (s, 6H, **1**), **2.39** (s, 3H, **14**), **4.13** (s, 2H, **3**), **5.50** (s, 1H, **6**), **7.33** (d, *J* = 8.3 Hz, 1H, **9**), **7.59** (dd, *J* = 8.3, 1.8 Hz, 1H, **13**) **7.71** (d, *J* = 1.8 Hz, 1H, **12**), **9.93** (s, 1H, **4**)

¹³C-NMR: (101MHz, CDCl₃) δ 20.2 (**14**); 27.1 (**1**); 58.6 (**2**); 73.9 (**6**); 79.0 (**3**); 125.6 (**13**); 128.8 (**9**); 129.4 (**12**); 135.7 (**10**); 136.7 (**8**); 138.5 (**11**); 169.6 (**5**); 186.4 (**7**)

EA: (%): calc'd. (found) for C₁₄H₁₆ClNO₂: C 63.28 (63.17); H 6.07 (5.97); N 5.27 (5.19).

MS: (m/z): calc'd (found): [M⁺] = 265.7 (265.8)

4.3.7 Synthesis of (*E*)-2-(4,4-dimethyloxazolidin-2-ylidene)-1-(naphthalen-1-yl)ethan-1-one (**4g**)

In a 150-mL beaker, KOH (4.52 g, 80.7 mmol, 1.46 M) was dissolved in MeOH (55 mL) at room temperature. The KOH/MeOH mixture was added to a 100-mL flask containing **3g** (10.18 g, 24.08 mmol) was added to the mixture and was stirred for 16 hours at room temperature. Volatile components were removed by rotary evaporator. The resulting solids were diluted with dH₂O (50 mL) and washed with DCM (2 × 50 mL). Organic constituents were washed once more with dH₂O (1 × 100 mL). Magnesium sulphate was added, filtered and any volatiles removed *via* rotary evaporator. The orange gel was purified via flash column chromatography with a solvent mixture containing 1:1 ethyl acetate/hexanes. The target product was collected as either a pale yellow solid or white solid (1.5 g, 23 % yield).

Melting Point Range: 129-132°C

TLC: 50% EtOAc/*n*-Hexanes, *R_f* = 0.29

¹H-NMR: (400 MHz, CDCl₃) δ **1.46** (s, 6H, **1**), **4.14** (s, 2H, **3**), **5.36** (s, 1H, **6**), **7.43-7.57** (m, 3H, **11,13,16**), **7.63** (d, *J* = 7.0 Hz, 1H, **12**) **7.85** (d, *J* = 7.7 Hz, 2H, **15,17**), **8.48** (d, *J* = 8.20, 1H, **10**) **9.93** (s, 1H, **4**)

¹³C-NMR: 27.2 (**1**); 58.6 (**2**); 78.8 (**6**); 79.0 (**3**); 124.9 (**Ar**); 125.1 (**12**); 125.8 (**Ar**); 126.3 (**Ar**) 126.3 (**10**); 128.1 (**15/17**); 129.5 (**15/17**); 130.3 (**14**); 133.8 (**9**); 140.6 (**8**); 169.1 (**5**); 191.6 (**7**)

EA: (%): calc'd. (found) for C₁₇H₁₇NO₂: C 76.38 (76.29); H 6.41 (6.35); N 5.24 (5.11).

MS: (m/z): calc'd (found): [M⁺] = 258.1 (258.0)

4.3.8 Synthesis of (*E*)-1-(4,4-dimethyloxazolidin-2-ylidene)-3,3-dimethylbutan-2-one (**4h**)

In a 150-mL beaker, KOH (8.537 g, 152.0 mmol, 1.20 M) was dissolved in MeOH (100 mL) at room temperature. The KOH/MeOH mixture was added to a 100-mL flask containing **3h** (10.99 g, 39.05 mmol) was added to the mixture and stirred for 18 hours at room temperature. Volatile components were removed *via* rotary evaporator. The resulting solid mass was diluted with dH₂O (60 mL) and washed with CHCl₃ (2 × 60 mL). Organic constituents were washed once more with dH₂O (1 × 120 mL). Magnesium sulphate was added, filtered and any volatiles removed *via* rotary evaporator. The crude off-white powder was recrystallized with a mixture of 1:4 toluene/hexanes to afford white needle-like crystals. (7.3 g, 94 % yield). Analyses of this compound are consistent with those in the literature.^{20,35}

Melting Point Range: 94-95 °C (lit: 94.5-95 °C)

TLC: 45 % EtOAc/Hexanes R_f = 0.49

4.3.9 Synthesis of (*E*)-1-(4,4-dimethyloxazolidin-2-ylidene)propan-2-one (**4j**)

A Schlenk flask was charged with dry THF (50.0 mL) and **1a** (10.0 mL, 8.87 g, 78.4 mmol) and cooled to -84 °C. Once cooled, *n*-BuLi (1.0 M in *n*-hexanes, 60.0 mL, 90.0 mmol) was added dropwise over 45 minutes. As the solution stirred overnight (~18 hours), the colour turned from a pale yellow to a vibrant, deep yellow. The solution was cooled back down to -84 °C, after which EtOAc (15.0 mL, 13.5 g, 153 mmol) in THF (10 mL) was added dropwise to the stirring solution. The mixture was stirred for 2 hours where it gradually warmed to room temperature, where it turned a dark orange colour, before quenching with degassed dH₂O. It was washed with Et₂O (100 mL × 2). Magnesium sulphate was added to dry the solution. The mixture was then filtered and the volatile constituents were removed *via* a rotary evaporator. This left behind crude mixture of **4j** and the **dimer** (above). The mixture was stirred at 60 °C in dH₂O for 4 hours. After cooling to room temperature, the mixture was washed with Et₂O (60 mL × 2). Magnesium sulphate was added to dry the solution. The mixture was then filtered and the volatile constituents were removed *via* a rotary evaporator. The solids were rinsed with cold *n*-heptanes and separated to reveal **4j** (2.5 g, 20 % yield). Analyses of this compound are consistent with those in the literature.^{34,36}

Melting Point Range: 123-126°C (lit: 125-127 °C)

¹H-NMR: (400 MHz, CDCl₃) δ **1.25** (s, 6H, **1'**), **1.39** (s, 6H, **1**), **2.02** (s, 3H, **8**), **2.24** (s, 3H, **8'**), **3.39** (s, 2H, **3'**), **3.98** (s, 2H, **6'**), **4.07** (s, 2H, **3**), **4.87** (s, 1H, **6**), **9.47** (s, 1H, **4**)

¹³C-NMR: (101MHz, CDCl₃) δ 27.0 (**1**); 28.2 (**1'**); 28.7 (**8**); 29.5 (**8'**); 43.6 (**6'**); 58.3 (**2**); 67.4 (**2'**); 76.9 (**6**); 78.7 (**3'**); 79.5 (**3**); 159.9 (**5'**); 168.3 (**5**) 194.2 (**7**); 201.5 (**7'**)

4.3.10 Synthesis of (*E*)-3-(4,4-dimethyloxazolidin-2-ylidene)-1,1,1-trifluoropropan-2-one (**4k**)

A round-bottom flask was charged with **1a** (2.00 mL, 1.77 g, 15.7 mmol), py (7.60 mL, 7.46 g, 94.4 mmol) and ACN (10.0 mL) and cooled to 0 °C. In a separate round-bottom flask, TFAA (4.80 mL, 7.15 g, 34.1 mmol) and ACN (10.0 mL) were cooled to 0 °C. Once cooled, the anhydride was transferred to the oxazoline containing flask *via* dropwise addition over a period of 10 minutes. The reaction was stirred at 0 °C for a period of 6 hours, periodically replacing the cold bath. Upon completion, excess volatile components were removed *via* a rotary evaporator. The resulting sludge was dissolved in brine (30 mL) and washed with DCM (2 × 30 mL) and additionally once with brine (60 mL). Magnesium sulphate was added to dry the solution. The mixture was then filtered and the volatile constituents were removed *via* a rotary evaporator. The mixture was dissolved in a minimal amount of hot *n*-hexanes. Once cooled, the target product was revealed as a fluffy white powder (2.1 g, 64 % yield). Analyses of this compound are consistent with those in the literature.⁸

Melting Point Range: 140-142°C (lit: 141-142 °C)

¹H-NMR: (400 MHz, CDCl₃) δ **1.43** (s, 6H, **1**), **4.19** (s, 2H, **3**), **5.16** (s, 1H, **6**), **9.86** (s, 1H, **4**)

¹⁹F-NMR: (376 MHz, CDCl₃) δ -76.4 (**9**, 3F)

¹³C-NMR: (101 MHz, CDCl₃) δ 26.6 (**1**); 59.7 (**2**); 72.6 (**6**); 79.8 (**3**); 117.9 (**q**, *J*_{F-C} = 288.5 Hz, **8**); 170.3 (**5**); 175.3 (**q**, *J*_{F-C} = 33.0 Hz, **7**)

4.3.11 Synthesis of (*E*)-1,1,1-trichloro-3-(4,4-dimethyloxazolidin-2-ylidene)propan-2-one (**4I**)

A round-bottom flask was charged with **1a** (2.00 mL, 1.77 g, 15.7 mmol), pyridine (2.80 mL, 2.75 g, 34.8 mmol) and ACN (10.0 mL) and cooled to 0 °C. In a separate round-bottom flask, TCAC (3.80 mL, 6.16 g, 33.9 mmol) and ACN (10.0 mL) were cooled to 0 °C. Once cooled, the acid chloride was transferred to the oxazoline containing flask *via* dropwise addition over a period of 10 minutes. The reaction was stirred at 0 °C for a period of 6 hours, periodically replacing the cold bath. Upon completion, excess volatile components were removed *in vacuo*. The resulting sludge was dissolved in dH₂O (40 mL) and washed with DCM (2 × 40 mL). Magnesium sulphate was added to dry the solution. The mixture was then filtered and the volatile constituents were removed *via* a rotary evaporator. The mixture was washed with hot *n*-hexanes. Once cooled, the target product was revealed as a fluffy white powder (1.1 g, 25 % yield). Analyses of this compound are consistent with those in the literature.⁸

Melting Point Range: 140-142°C (lit: 142-144 °C)

¹H-NMR: (400 MHz, CDCl₃) δ **1.45** (s, 6H, **1**), **4.23** (s, 2H, **3**), **5.55** (s, 1H, **6**), **9.35** (s, 1H, **4**)

¹³C-NMR: (101 MHz, CDCl₃) δ 27.2 (**1**); 59.1 (**2**); 69.8 (**6**); 79.8 (**3**); 97.2 (**8**); 170.9 (**5**); 181.1 (**7**)

4.4 Synthesis of the ligand series with **1b**

4.4.1 Synthesis of (*E*)-2-(oxazolidin-2-ylidene)-1-phenylethan-1-one (**5a** to **6a**);

A flask was charged with **1b** (5.06 mL, 5.03 g, 59.1 mmol), THF (20 mL) and NEt₃ (20.6 mL, 14.9 g, 148 mmol). The solution was cooled to 0 °C, benzoyl chloride (15.1 mL, 18.3 g, 130 mmol) was added. The mixture was refluxed for 3 hours. The solution was cooled to room temperature, the salt was filtered and the volatile components were removed *via* a rotary evaporator. dH₂O (50 mL) was added, and the solution was washed with CHCl₃ (2 × 50 mL). The organics were washed with a 10 % solution of Na₂CO_{3(aq)} (1 × 100 mL). The organic phase was dried (Na₂SO₄), filtered and the volatile constituents removed *via* a rotary evaporator. KOH (7.31 g, 130 mmol, 2.17 M) was dissolved in MeOH (60 mL) and added to **5a** (0.20 g, 0.44 mmol). The contents were stirred for 16 hours at room temperature. Volatile components were removed *via* a rotary evaporator. dH₂O (50 mL) was added and washed with DCM (2 × 50 mL) and washed with dH₂O (1 × 100 mL). The solution was dried (Na₂SO₄), filtered and the volatile constituents removed *via* a rotary evaporator. The gel was purified with column chromatography eluting with 2:1 acetone/*n*-hexanes. The target product was collected as pinkish-white solids (3.66 g, 33.0 % yield). Analyses of this compound are consistent with those in the literature.^{9,40}

Melting Point Range: 105-107 °C (lit: 105-106 °C)

TLC: 40% acetone/*n*-hexanes, R_f = 0.30

¹H-NMR: (400MHz, CDCl₃) δ **3.82** (t, *J* = 8.5 Hz, 2H, **1**), **4.49** (t, *J* = 8.5 Hz, 2H, **2**), **5.64** (s, 1H, **5**), **7.39-7.41** (m, 3H, **9-10**), **7.87** (dd, *J* = 7.5, 1.0 Hz, 2H, **8**), **10.00** (s, 1H, **3**)

¹³C-NMR: (101MHz, CDCl₃) δ 43.2 (**1**); 67.5 (**2**), 74.3 (**5**); 127.0 (**8**); 128.3 (**9**); 130.7 (**10**); 140.1 (**7**); 170.9 (**4**); 187.8 (**6**)

4.4.2 Synthesis of (*E*)-1-(3,4-dichlorophenyl)-2-(oxazolidin-2-ylidene)ethan-1-one (**5b** to **6b**)

A flask was charged with **1b** (2.00 mL, 2.01 g, 23.4 mmol) and THF (45 mL). The solution was cooled to 0 °C, 3,4-dichlorobenzoyl chloride (10.63 g, 50.73 mmol) was added. NEt₃ (10.00 mL, 7.260 g, 71.75 mmol) was added and was refluxed for 3 hours. The solution was cooled to room temperature, the salt filtered and the volatile components were removed *via* a rotary evaporator. dH₂O (50 mL) was added, and the solution was washed with CHCl₃ (2 × 50 mL). The organics were washed with a 10 % solution of Na₂CO_{3(aq)} (1 × 100 mL). The solution was dried (Na₂SO₄), filtered and volatile constituents were removed *via* a rotary evaporator (10.13 g). KOH (2.30 g, 40.9 mmol, 1.64 M) was dissolved in MeOH (25 mL) and added to **5b** (0.20 g, 0.44 mmol). The contents were stirred for 3 hours at room temperature. The solids were diluted with dH₂O (50 mL) and washed with DCM (2 × 50 mL). Organics were washed with dH₂O (1 × 100 mL). The solution was dried (Na₂SO₄), filtered and volatile constituents were removed *via* a rotary evaporator. The gel was recrystallized with a mixture of 1:4 toluene/*n*-hexanes to collect a pale yellow solid (110 mg, 92 % yield).

TLC: 45 % EtOAc/*n*-hexanes R_f = 0.07

¹H-NMR: (400 MHz, CDCl₃) δ **3.83** (t, *J* = 8.5 Hz, 2H, **1**), **4.52** (t, *J* = 8.5 Hz, 2H, **2**), **5.53** (s, 1H, **5**), **7.44** (d, *J* = 8.5 Hz, 1H, **11**), **7.65** (dd, *J* = 2.0, 8.5 Hz, 1H, **12**), **7.92** (d, *J* = 2.0 Hz, H, **8**), **9.94** (s, 1H, **4**)

¹³C-NMR: 43.2 (**1**); 67.8 (**2**); 74.2 (**5**); 126.2 (**12**); 129.1 (**8**); 130.3 (**11**); 132.66 (**Ar**); 134.7 (**Ar**); 140.0 (**7**); 171.2 (**3**); 184.9 (**6**)

EA: (%): calc'd. (found) for C₁₁H₉Cl₂NO₂: C 51.19 (49.92); H 3.51 (3.39); N 5.43 (5.25).
0.5 water molecules found in the lattice in elemental analysis

4.4.3 Synthesis of (*E*)-1-(4-methoxyphenyl)-2-(oxazolidin-2-ylidene)ethan-1-one (**5c** to **6c**)

A flask was charged with **1b** (2.00 mL, 2.01 g, 23.4 mmol) and THF (40 mL). The solution was cooled to 0 °C, *p*-methoxybenzoyl chloride (7.01 g, 51.9 mmol) was added. NEt₃ (10.00 mL, 7.260 g, 71.75 mmol) was added and was refluxed for 3 hours. The solution was cooled to room temperature, the salt filtered and volatile components were removed *via* a rotary evaporator. dH₂O (50 mL) was added and was washed with CHCl₃ (2 × 50 mL). The organics were washed with a 10 % solution of Na₂CO_{3(aq)} (1 × 100 mL). The solution was dried (Na₂SO₄), filtered and volatile constituents were removed *via* a rotary evaporator (8.34 g). KOH (2.26 g, 40.2 mmol, 1.61 M) was dissolved in MeOH (25 mL) at room temperature and was added to a 100-mL flask containing **5c** (8.34 g, 23.4 mmol). The contents were stirred for 3 hours at room temperature. The solids were diluted with dH₂O (50 mL) and washed with DCM (2 × 50 mL). Organic constituents were washed once more with dH₂O (1 × 100 mL). The organic phase was dried (Na₂SO₄), filtered and the volatile constituents were removed *via* a rotary evaporator. The oil was purified *via* column chromatography eluting with 30% EtOAc/*n*-hexanes. The target was collected a pale yellow solid (0.63 mg, 12 % yield). Analyses of this compound are consistent with those in the literature.^{38,39}

TLC: 45 % EtOAc/*n*-hexanes R_f = 0.12

¹H-NMR: (400 MHz, CDCl₃) δ **3.80-3.88** (m, 5H, **1,13**), **4.50** (t, *J* = 8.0 Hz, 2H, **2**), **5.60** (s, 1H, **5**), **6.89** (dt, *J* = 3.0, 9.0 Hz, 2H, **9,11**), **7.83** (dt, *J* = 3.0, 9.0 Hz, 1H, **8,12**), **9.94** (s, 1H, **4**)

¹³C-NMR: (101 MHz, CDCl₃) δ 43.2 (**1**); 55.5 (**13**); 67.5 (**2**); 73.7 (**5**); 113.5 (**9,11**); 128.8 (**8,12**); 132.8 (**10**); 161.9 (**7**); 170.7 (**3**); 187.2 (**6**)

4.4.4 Attempted Synthesis of (*E*)-1-(4-nitrophenyl)-2-(oxazolidin-2-ylidene)ethan-1-one (**5d/6d**)

A flask was charged with **1b** (1.00 mL, 1.01 g, 11.8 mmol), NEt₃ (5.00 mL, 3.63 g, 35.9 mmol) and dry THF (20 mL). *p*-Nitrobenzoyl chloride (4.891 g, 26.36 mmol) was dissolved in dry THF (30 mL) and added dropwise to the stirring **1b**-mixture. The reaction was set to reflux for 3 hours. The solution was cooled to room temperature where the incorporated salt was removed by filtration and the volatile components were removed *via* a rotary evaporator. To the resulting brown sludge, dH₂O (50 mL) was added then the solution was washed with DCM (2 × 50 mL). The organic components were washed one last time with a 5 % solution of Na₂CO_{3(aq)} (1 × 100 mL). Magnesium sulphate was added to dry the solution. The mixture was then filtered and the volatile constituents were removed *via* a rotary evaporator. The resulting orange sludge was purified via column chromatography eluting with 2:1 acetone/*n*-hexanes. The resulting white powder (434 mg, 1.13 mmol) was dissolved in MeOH (55 mL) and KOH (4.53 g, 80.6 mmol, 1.46 M) was added. The contents were stirred at room temperature for 6 hours. Volatile components were removed *via* a rotary evaporator. The solids were diluted with dH₂O (60 mL) and washed with DCM (2 × 60 mL). Organic constituents were washed once more with dH₂O (1 × 120 mL). The solution was dried (MgSO₄) filtered and the volatile constituents were removed *via* a rotary evaporator. The crude wine-red powder was purified by column chromatography, eluting with 45% EtOAc/*n*-hexanes. All attempts of this reaction were met with no success. All NMR spectra showed resonance signals indicative of the starting material (*p*-nitrobenzoyl chloride).

4.5 Synthesis and Characterization of **Cu** complexes made with **1a**

4.5.1 Chelation of **4a** to Copper metal to obtain **7a**

A round-bottom flask was charged with **4a** (0.502 g, 2.39 mmol), NEt₃ (1.00 mL, 0.726 g, 7.17 mmol), and EtOH (5 mL). After the yellow-coloured solids were completely dissolved, Cu(NO₃)₂•2.5 H₂O (0.269 g, 1.16 mmol), a bright blue compound, was dissolved in EtOH (5 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a deep green. The solution was stirred at room temperature for 1 hour. Following completion of the reaction, green precipitates were collected which was recrystallized from a mixture of 1:1 DCM/MeOH (474 mg, 81 % yield). Analyses of this compound are consistent with those in the literature.²⁰

MP: Green plates: 164-156 °C (lit: 165 °C decomp.)

IR: Green plates (cm⁻¹) 2892, 1592, 1572, 1531, 1487, 1412, 1351, 1309, 1173, 1106, 1032, 994, 944, 861, 837, 790, 751, 732, 659

MP: Brown needles: 158-160 °C

IR: Brown needles (cm⁻¹) 2882, 1561, 1527, 1496, 1455, 1429, 1349, 1312, 1250, 1194, 1140, 1105, 1048, 1025, 993, 949, 873, 836, 832, 756, 735, 695

4.5.2 Chelation of **4b** to Copper metal to obtain **7b**

A round-bottom flask was charged with **4b** (1.01 g, 3.49 mmol) and EtOH (10 mL). After the yellow-coloured solids were completely dissolved, Cu(NO₃)₂•2.5 H₂O (0.402 g, 1.75 mmol), a soft black compound, was dissolved in EtOH (10 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark green. The solution was stirred at room temperature for 16 hours. Following completion of the reaction, dark green precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (0.628, 57% yield).

Melting Point Range: 208-210°C

IR: (cm⁻¹) 2964, 1750, 1593, 1527, 1458, 1353, 1304, 1179, 1139, 1101, 1027, 997, 943, 889, 823, 764, 727, 671

UV-Vis: (DCM: 1.89 × 10⁻⁵M) λ_{max1} = 434 nm (ε = 1.62 × 10³ L·mol⁻¹·cm⁻¹), λ_{max2} = 335 nm (ε = 3.39 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max3} = 296 nm (ε = 3.42 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max4} = 246 nm (ε = 4.21 × 10⁴ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₂₆H₂₄N₂O₄Cl₄Cu: C 49.27 (49.13); H 3.82 (3.63); N 4.42 (4.45).

MS: (m/z) found (calc'd) [M+H]⁺ = 634.1 (633.8)

4.5.3 Chelation of **4c** to Copper metal to obtain **7c**

A round-bottom flask was charged with **4c** (0.10 g, 0.40 mmol) and EtOH (15 mL). NEt₃ (0.1 mL, 0.72 mmol) was added to the solution. After the yellow-coloured solids were completely dissolved, Cu(NO₃)₂·2.5H₂O (0.045 g, 0.19 mmol), a blue-coloured compound, was dissolved in EtOH (5 mL) and added to the solution. Upon addition of the two, the solution turned colour to a dark green. The solution was stirred at room temperature for 2 hours. Following completion of the reaction, dark green precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (103 mg, 96 % yield).

Melting Point Range: 203-205°C

IR: (cm⁻¹) 2890, 1896, 1740, 1589, 1531, 1496, 1434, 1355, 1298, 1254, 1168, 1114, 1028, 999, 867, 836, 788, 766, 733, 701, 675

UV-Vis: (DCM: 7.91×10^{-5} M) $\lambda_{\max 1} = 455$ nm ($\epsilon = 478$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 2} = 325$ nm ($\epsilon = 1.33 \times 10^4$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 3} = 299$ nm ($\epsilon = 1.32 \times 10^4$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 4} = 252$ nm ($\epsilon = 8.03 \times 10^3$ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₂₈H₃₂N₂O₆Cu: C 60.47 (60.66); H 5.80 (5.61); N 5.04 (5.08).

MS: (m/z) found (calc'd) [M]⁺ = 556.3 (556.1)

4.5.4 Chelation of **4d** to Copper metal to obtain **7d**

A round-bottom flask was charged with **4d** (0.149 g, 0.607 mmol) and EtOH (10 mL). After the yellow-coloured solids were completely dissolved, Cu(NO₃)₂·2.5 H₂O (0.069 g, 0.30 mmol), a bright blue-coloured compound, was dissolved in EtOH (5 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark green. NEt₃ (1.00 mL, 0.726 g, 7.17 mmol) was added as a base. The solution stirred at room temperature for 1 hour. Following completion of the reaction, green precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (150 mg, 91 % yield). EA is pending.

Melting Point Range: 218-220°C

IR: (cm⁻¹) 2961, 1740, 1590, 1561, 1526, 1488, 1457, 1352, 1308, 1191, 1129, 1095, 131, 992, 944, 881, 811, 764, 731, 700

UV-Vis: (DCM: 1.45×10^{-5} M) $\lambda_{\max 1} = 443$ nm ($\epsilon = 1.72 \times 10^3$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 2} = 325$ nm ($\epsilon = 4.04 \times 10^4$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 3} = 293$ nm ($\epsilon = 5.18 \times 10^4$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 4} = 250$ nm ($\epsilon = 3.74 \times 10^4$ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₃₀H₃₆N₂O₄Cu: C 65.26 (64.99); H 6.57 (6.50); N 5.07 (5.07).

MS: (m/z) found (calc'd) [M]⁺ = 552.3 (552.2)

4.5.5 Chelation of **4e** to Copper metal to obtain **7e**;

A round-bottom flask was charged with **4e** (0.499 g, 1.88 mmol) and EtOH (20 mL). After the yellow-coloured solids were completely dissolved, Cu(NO₃)₂·2.5 H₂O (0.213 g, 0.918 mmol), a bright blue-coloured compound, was dissolved in EtOH (10 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark green. The solution was stirred at room temperature for 18 hours. Following completion of the reaction, green precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (176 mg, 32 % yield).

Melting Point Range: 207-209 °C

IR: (cm⁻¹) 2963, 1589, 1529, 1483, 1457, 1355, 1305, 1179, 1114, 1030, 997, 942, 894, 799, 765, 708

UV-Vis: (DCM: 2.83×10^{-5} M) $\lambda_{\max 1} = 445$ nm ($\epsilon = 815$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 2} = 330$ nm ($\epsilon = 2.13 \times 10^4$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 3} = 301$ nm ($\epsilon = 2.16 \times 10^4$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 4} = 244$ nm ($\epsilon = 1.99 \times 10^4$ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₂₈H₃₀Cl₂N₂O₄Cu: C 56.71 (56.94); H 5.10 (5.16); N 4.72 (4.80).

MS: (m/z) found (calc'd) [M]⁺ = 594.1 (593.1)

4.5.6 Chelation of **4f** to Copper metal to obtain **7f**;

A round-bottom flask was charged with **4f** (0.450 g, 1.69 mmol) and EtOH (10 mL). After the yellow-coloured solids were completely dissolved, Cu(NO₃)₂·2.5 H₂O (0.196 g, 0.839 mmol), a bright blue-coloured compound, was dissolved in EtOH (10 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark green. The solution was stirred at room temperature for 16 hours. Following completion of the reaction, dark green precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (380 mg, 76 % yield).

Melting Point Range: >210 °C (decomp.)

IR: (cm⁻¹) 2959, 2113, 1750, 1593, 1563, 1529, 1466, 1352, 1307, 1198, 1102, 1031, 1002, 946, 878, 826, 807, 766, 731, 677

UV-Vis: (DCM: 4.72×10^{-5} M) $\lambda_{\text{max}1} = 436 \text{ nm}$ ($\epsilon = 788 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}2} = 330 \text{ nm}$ ($\epsilon = 1.81 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}3} = 299 \text{ nm}$ ($\epsilon = 1.88 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}4} = 247 \text{ nm}$ ($\epsilon = 1.79 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$).

EA: (%): calc'd. (found) for C₂₈H₃₀Cl₂N₂O₄Cu: C 56.71 (56.89); H 5.10 (5.01); N 4.72 (4.75).

MS: (m/z) found (calc'd) [C₁₄H₁₆ClNO₂] = 266.1 (265.7), [M]⁺ = 594.3 (593.1)

4.5.7 Chelation of **4g** to Copper metal to obtain **7g**;

A round-bottom flask was charged with **4g** (0.299 g, 1.12 mmol) and EtOH (10 mL). After the yellow-coloured solids were completely dissolved, Cu(NO₃)₂·2.5 H₂O (0.128 g, 0.55 mmol), a bright blue-coloured compound, was dissolved in EtOH (5 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark green. NEt₃ (0.20 mL, 0.14 g, 1.4 mmol) was added as a base. The solution stirred at room temperature for 1 hour. Following completion of the reaction, green precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (300 mg, 91 % yield).

Melting Point Range: 171-173°C

IR: (cm⁻¹) 3041, 2955, 2112, 1916, 1581, 1548, 1457, 1420, 1390, 1352, 1308, 1260, 1199, 1174, 1062, 1040, 1000, 935, 880, 771, 714, 669

UV-Vis: (DCM: 2.01 × 10⁻⁵M) λ_{max1} = 425 nm (ε = 1.21 × 10³ L·mol⁻¹·cm⁻¹), λ_{max2} = 293 nm (ε = 3.21 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max3} = 247 nm (ε = 3.33 × 10⁴ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₃₄H₃₂N₂O₄Cu: C 68.50 (68.47); H 5.41 (5.34); N 4.70 (4.78).

MS: (m/z) found (calc'd) [M]⁺ = 596.3 (596.2)

4.5.8 Chelation of **4h** to Copper metal to obtain **7h**

A round-bottom flask was charged with **4h** (0.128 g, 0.649 mmol) and EtOH (10 mL). After the yellow-coloured solids were completely dissolved, Cu(NO₃)₂·2.5 H₂O (0.0740 g, 0.318 mmol), a bright blue-coloured compound, was dissolved in EtOH (10 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark green. NEt₃ (0.20 mL, 1.4 mmol) was added to as a base. The solution was stirred at room temperature for 1 hour. Following completion of the reaction, green precipitates were collected which recrystallized in a mixture of 1:1 DCM/MeOH (73 mg, 50 % yield). Analyses of this complex are consistent with those in the literature.⁶⁶

Melting Point Range: 203-204 °C

IR: (cm⁻¹) 2958, 1590, 1527, 1474, 1415, 1349, 1297, 1249, 1197, 1153, 1036, 1003, 945, 890, 848, 803, 788, 729

UV-Vis: (DCM: 2.10 × 10⁻⁵M) λ_{max1} = 411 nm (ε = 1.11 × 10³ L·mol⁻¹·cm⁻¹), λ_{max2} = 309 nm (ε = 8.15 × 10³ L·mol⁻¹·cm⁻¹), λ_{max3} = 275 nm (ε = 2.38 × 10⁴ L·mol⁻¹·cm⁻¹)

EA: (%): calc'd. (found) for C₂₂H₃₆N₂O₄Cu: C 57.94 (58.17); H 7.96 (7.85); N 6.14 (5.92).

4.5.9 Chelation of **4i** to a Copper metal to obtain **7i**;

A round-bottom flask was charged with **4i** (0.122 g, 0.465 mmol) and EtOH (10 mL). After the yellow-coloured solids were completely dissolved, Cu(NO₃)₂·2.5 H₂O (0.0530 g, 0.228 mmol), a bright blue-coloured compound, was dissolved in EtOH (10 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark green. NEt₃ (1.0 mL, 7.2 mmol) was added to as a base. The solution was stirred at room temperature for 1 hour. Following completion of the reaction, green precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (120 mg, 90 % yield). Analyses of this complex are consistent with those in the literature.⁶⁶

Melting Point Range: 217-218 °C

IR: (cm⁻¹) 1584, 1535, 1514, 1484, 1341, 1299, 1250, 1181, 1108, 1031, 998, 946, 861, 834, 779, 752, 750

UV-Vis: (solv: 1.43 × 10⁻⁵M) λ_{max1} = 717 nm (ε = 550 L·mol⁻¹·cm⁻¹), λ_{max2} = 374 nm (ε = 1.78 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max3} = 306 nm (ε = 1.41 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max4} = 269 nm (ε = 3.11 × 10⁴ L·mol⁻¹·cm⁻¹)

EA: (%): calc'd. (found) for C₂₆H₂₆N₄O₈Cu: C 53.29 (53.17); H 4.47 (4.45); N 9.56 (9.45).

4.6 Synthesis and Characterization of **Cu** complexes made with **1b**

4.6.1 Chelation of **6a** to Copper metal to obtain **8a**

A round-bottom flask was charged with **6a** (0.755 g, 3.99 mmol) and MeOH (15 mL). After the pink-coloured solids were completely dissolved, CuBr₂ (0.560 g, 2.51 mmol) was added to the solution. Upon addition of the two, the solution turned colour from pink to a pastel green. The solution was stirred at room temperature for 3 hours. Upon completion of the reaction a green precipitate was collected *via* vacuum filtration (0.59 g, 70% yield). The synthesized substance is unable to be properly characterized due to its extreme limited solubility in only THF.

Melting Point Range: >260°C (decomp.)

IR: (cm⁻¹) 2890, 1605, 1578, 1540, 1487, 1454, 1414, 1360, 1328, 1263, 1200, 1110, 1063, 995, 951, 750, 689, 655

UV-Vis: (DCM: 1.36×10^{-4} M) $\lambda_{\text{max}1} = 320 \text{ nm}$ ($\epsilon = 7.17 \times 10^3 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}2} = 285 \text{ nm}$ ($\epsilon = 8.58 \times 10^3 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}3} = 233 \text{ nm}$ ($\epsilon = 8.45 \times 10^3 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$).

EA: (%): calc'd. (found) for C₂₂H₂₀N₂O₄Cu: C 60.06 (59.89); H 4.58 (4.76); N 6.37 (6.36).

MS: (m/z) found (calc'd): not detected (440.0)

4.6.2 Chelation of **6b** to Copper metal to obtain **8b**

A round-bottom flask was charged with **6b** (39 mg, 0.15 mmol) and EtOH (2 mL). After the yellow-coloured solids were completely dissolved, Cu(NO₃)₂·2.5 H₂O (17 mg, 0.073 mmol) dissolved in EtOH (3 mL) was added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark green. NEt₃ (0.10 mL, 73 mg, 72 mmol) was added to the reaction. The solution was stirred at room temperature for 1 hour. Upon completion of the reaction a pastel green-coloured precipitate was collected *via* vacuum filtration (30 mg, 71 % yield). The synthesized substance is unable to be properly characterized due to its extreme limited solubility in only THF. Characterization on this compound is pending. Crude characterization represents results like **8a**.

4.7 Synthesis and Characterization of **Co** complexes made with **1a**

4.7.1 Chelation of **4a** to a Cobalt metal to obtain **9a**

A round-bottom flask was charged with **4a** (1.01 g, 4.64 mmol), NEt₃ (3.00 mL, 2.18 g, 21.5 mmol), and EtOH (10 mL). After the yellow-coloured solids were completely dissolved, Co(NO₃)₂•6 H₂O (0.671 g, 2.31 mmol), a pink compound, was dissolved in EtOH (5 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a bright red. The solution was stirred at room temperature for 2 hours. Following completion of the reaction, red precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (0.927 g, 81 % yield). Analyses of this compound are consistent with those in the literature.⁶⁶

Melting Point Range: 198-201°C

IR: (cm⁻¹) 2986, 2889, 1740, 1589, 1570, 1520, 1485, 1471, 1407, 1353, 1304, 1172, 1103, 994, 952, 860, 840, 754, 695, 653

UV-Vis: (DCM: 5.70 × 10⁻⁵M) λ_{max1} = 534 nm (ε = 60.0 L·mol⁻¹·cm⁻¹), λ_{max2} = 314 nm (ε = 2.39 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max3} = 232 nm (ε = 2.14 × 10⁴ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₂₆H₂₈N₂O₄Co: C 63.54 (63.49); H 5.74 (5.72); N 5.70 (5.43).

MS: (m/z) found (calc'd) [M+H]⁺ = 492.3 (491.5)

4.7.2 Chelation of **4b** to a Cobalt metal to obtain **9b**

A round-bottom flask was charged with **4b** (0.533 g, 1.86 mmol) and MeOH (10 mL). After the yellow-coloured solids were completely dissolved, Co(NO₃)₂•6 H₂O (0.27 g, 0.92 mmol), a pink compound, was dissolved in MeOH (15 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a bright red. The solution was stirred at room temperature for 18 hours. Following completion of the reaction, orange precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (331 mg, 56 % yield). Analyses of this compound are consistent with those in the literature.⁶⁶

Melting Point Range: 223-225°C

IR: (cm⁻¹) 2960, 1739, 1581, 1524, 1461, 1428, 1354, 1303, 1256, 1178, 1141, 1102, 1025, 878, 766, 670

UV-Vis: (DCM: 2.54 × 10⁻⁵M) λ_{max1} = 538 nm (ε = 113 L·mol⁻¹·cm⁻¹), λ_{max2} = 321 nm (ε = 3.88 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max3} = 245 nm (ε = 3.03 × 10⁴ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₂₆H₂₄N₂O₄Cl₄Co: C 49.63 (49.48); H 3.84 (3.90); N 4.45 (4.40).

MS: (m/z) found (calc'd) [M+H]⁺ = 630.1 (629.2)

4.7.3 Chelation of **4c** to a Cobalt metal to obtain **9c**

A round-bottom flask was charged with **4c** (0.103 g, 0.42 mmol) and EtOH (15 mL). NEt₃ (0.10 mL, 0.073 g, 0.72 mmol) was added to the solution. After the yellow-coloured solids were completely dissolved, Co(NO₃)₂·6H₂O (0.058 g, 0.20 mmol), a pink-coloured compound, was dissolved in EtOH (5 mL) and added to the solution. Upon addition of the two, the solution turned colour to a dark red. The solution was stirred at room temperature for 18 hours. Following completion of the reaction, red precipitates were collected and recrystallized in a mixture of 1:1 DCM/MeOH (118 mg, 99 % yield). Analyses of this compound are consistent with those in the literature.⁶⁶

Melting Point Range: 243-245°C

IR: (cm⁻¹) 2969, 2882, 1610, 1584, 1560, 1524, 1498, 1396, 1355, 1300, 1253, 1190, 1167, 1113, 1026, 999, 842, 769, 680

UV-Vis: (DCM: 1.27 × 10⁻⁴M) λ_{max1} = 540 nm (ε = 30.0 L·mol⁻¹·cm⁻¹), λ_{max2} = 316 nm (ε = 1.14 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max3} = 263 nm (ε = 7.32 × 10³ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₂₈H₃₂N₂O₆Co: C 60.98 (60.90); H 5.85 (5.74); N 5.08 (5.10).

4.7.4 Chelation of **4d** to a Cobalt metal to obtain **9d**

A round-bottom flask was charged with **4d** (0.088 g, 0.34 mmol) and EtOH (3 mL). NEt₃ (0.20 mL, 0.15 g, 1.4 mmol) was added to the solution. After the yellow-coloured solids were completely dissolved, Co(NO₃)₂·6H₂O (0.048 g, 0.17 mmol), a pink-coloured compound, was dissolved in EtOH (2 mL) and added to the solution. Upon addition of the two, the solution turned colour to a dark red. The solution was stirred at room temperature for 1 hour. Following completion of the reaction, red precipitates were collected and recrystallized in a mixture of 1:1 DCM/MeOH (64 mg, 71 % yield).

Melting Point Range: 240°C (decomp.)

IR: (cm⁻¹) 2965, 1584, 1561, 1522, 1487, 1457, 1354, 1308, 1241, 1185, 1092, 1023, 991, 807, 768

UV-Vis: (DCM: 4.39×10^{-5} M) $\lambda_{\max 1} = 542$ nm ($\epsilon = 66.0$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 2} = 315$ nm ($\epsilon = 2.96 \times 10^4$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 3} = 257$ nm ($\epsilon = 1.78 \times 10^4$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 4} = 248$ nm ($\epsilon = 1.90 \times 10^4$ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₃₀H₃₆N₂O₄Co: C 63.71 (63.73); H 6.77 (6.57); N 4.95 (4.84).
- 1.0 H₂O found within lattice for elemental analysis

MS: (m/z) found (calc'd) [M+H]⁺ = 548.3 (547.6)

4.7.5 Chelation of **3e** to a Cobalt metal to obtain **9e**

To a round-bottom flask containing MeOH (25 mL), KOH (2.21 g, 39.4 mmol, 1.58 M) was added and stirred till all solids were dissolved. Once dissolved, **3e** (1.11 g, 2.65 mmol) was added. Separately, Co(NO₃)₂·6 H₂O (0.373 g, 1.28 mmol) was dissolved in MeOH (5 mL) and added to the swirling suspension. The resulting mixture was allowed the stir at room temperature for 6 hours. Following completion of the reaction all volatile components were removed *via* a rotary evaporator. The resulting solids were dissolved in DCM (50 mL) and washed with a solution of NaCl (0.20 mmol, 1.0 M). The organics were washed a second time with a solution of Na₂CO₃ (0.02 mmol, 0.1 M). The organic phase was dried (MgSO₄), filtered and the remaining volatile components were removed *via* a rotary evaporator. The crude orange-coloured precipitates were recrystallized in a mixture of 1:1 DCM/MeOH (0.519 g, 70 % yield)

Melting Point Range: 233-235°C

IR: (cm⁻¹) 2966, 1583, 1522, 1481, 1419, 1353, 1301, 1183, 1110, 1045, 1022, 995, 946, 890, 799, 767, 740, 731, 700

UV-Vis: (DCM: 6.80 × 10⁻⁵M) λ_{max1} = 533 nm (ε = 32.0 L·mol⁻¹·cm⁻¹), λ_{max2} = 317 nm (ε = 1.36 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max3} = 237 nm (ε = 1.07 × 10⁴ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₂₈H₃₀Cl₂N₂O₄Co: C 57.16 (56.89); H 5.14 (5.16); N 4.76 (4.80).

MS: (m/z) found (calc'd) [M]⁺ = 588.2 (588.3)

4.7.6 Chelation of **4f** to a Cobalt metal to obtain **9f**

A round-bottom flask was charged with **4f** (0.425 g, 1.60 mmol) and EtOH (15 mL). After the yellow-coloured solids were completely dissolved, Co(NO₃)₂·2.5 H₂O (0.227 g, 0.779 mmol), a pink-coloured compound, was dissolved in EtOH (10 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark red. The solution was stirred at room temperature for 18 hours. Following completion of the reaction, orange precipitates were collected and recrystallized in a mixture of 1:1 DCM/MeOH collected (459 mg, 61 % yield).

Melting Point Range: 234-236°C

IR: (cm⁻¹) 2967, 2114, 1738, 1583, 1562, 1521, 1470, 1424, 1354, 1306, 1198, 1178, 1148, 1100, 1045, 1023, 993, 946, 877, 831, 802, 767, 690

UV-Vis: (DCM: 1.04×10^{-4} M) $\lambda_{\text{max}1} = 536 \text{ nm}$ ($\epsilon = 41.0 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}2} = 317 \text{ nm}$ ($\epsilon = 1.96 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}3} = 292 \text{ nm}$ ($\epsilon = 1.56 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}4} = 247 \text{ nm}$ ($\epsilon = 1.59 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$).

EA: (%): calc'd. (found) for C₂₈H₃₀Cl₂N₂O₄Co: C 57.16 (56.88); H 5.14 (5.22); N 4.76 (4.77).

MS: (m/z) found (calc'd) [M]⁺ = 588.3 (588.3)

4.7.7 Chelation of **4g** to a Cobalt metal to obtain **9g**

A round-bottom flask was charged with **4g** (0.237 g, 0.887 mmol), NEt₃ (1.25 mL, 0.908 g, 8.97 mmol) and EtOH (10 mL). After the yellow-coloured solids were completely dissolved, Co(NO₃)₂·6 H₂O (0.127 g, 0.436 mmol), a bright pink-coloured compound, was dissolved in EtOH (5 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark red. The solution was stirred at room temperature for 1 hour. Following completion of the reaction, dark red precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (246 mg, 95 % yield). Analyses of this compound are consistent with those in the literature.⁶⁶

Melting Point Range: 201-202 °C

IR: (cm⁻¹) 3044, 2967, 2890, 1572, 1519, 1412, 1383, 1352, 1302, 1259, 1193, 1165, 1034, 989, 744, 666

UV-Vis: (DCM: 2.37×10^{-5} M) $\lambda_{\max 1} = 537$ nm ($\epsilon = 226$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 2} = 314$ nm ($\epsilon = 5.50 \times 10^4$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 3} = 297$ nm ($\epsilon = 6.13 \times 10^4$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 4} = 274$ nm ($\epsilon = 4.62 \times 10^4$ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₃₄H₃₂N₂O₄Co: C 69.03 (69.02); H 5.45 (5.44); N 4.74 (4.79).

MS: (m/z) found (calc'd) [M]⁺ = 592.3 (591.6)

4.7.8 Chelation of **4h** to a Cobalt metal to obtain **9h**

A round-bottom flask was charged with **4h** (0.119 g, 0.603 mmol) and EtOH (10 mL). After the yellow-coloured solids were completely dissolved, Co(NO₃)₂·6 H₂O (0.0850 g, 0.292 mmol), a bright pink-coloured compound, was dissolved in EtOH (10 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark red. NEt₃ (0.20 mL, 1.4 mmol) was added as a base. The solution was stirred at room temperature for 1 hour. Following completion of the reaction, pink precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (105 mg, 79 % yield). Analyses of this compound are consistent with those in the literature.⁶⁶

Melting Point Range: >220 °C (decomp.)

IR: (cm⁻¹) 2959, 1576, 1525, 1473, 1416, 1353, 1295, 1195, 1155, 1000, 948, 898, 803, 770, 729

UV-Vis: (solv: 1.77 × 10⁻⁵M) λ_{max1} = 537 nm (ε = 647 L·mol⁻¹·cm⁻¹), λ_{max2} = 321 nm (ε = 1.14 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max3} = 280 nm (ε = 8.37 × 10⁴ L·mol⁻¹·cm⁻¹)

EA: (%): calc'd. (found) for C₂₂H₃₆N₂O₄Co: C 58.53 (58.29); H 8.04 (8.07); N 6.21 (5.97).

4.7.9 Chelation of **4i** to a Cobalt metal to obtain **9i**

A round-bottom flask was charged with **4i** (0.120 g, 0.458 mmol) and EtOH (10 mL). After the yellow-coloured solids were completely dissolved, Co(NO₃)₂·6 H₂O (0.0670 g, 0.230 mmol), a bright pink-coloured compound, was dissolved in EtOH (10 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark red. NEt₃ (1.0 mL, 7.2 mmol) was added as a base. The solution was stirred at room temperature for 1 hour. Following completion of the reaction, orange precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH collected (97 mg, 73 % yield). Analyses of this compound are consistent with those in the literature.⁶⁶

Melting Point Range: > 240 °C (decomp.)

IR: (cm⁻¹) 2967, 1577, 1512, 1482, 1341, 1300, 1173, 1108, 1025, 992, 945, 862, 833, 777, 753, 703

UV-Vis: (solv: 6.88 × 10⁻⁶M) λ_{max1} = 363 nm (ε = 2.47 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max2} = 264 nm (ε = 3.33 × 10⁴ L·mol⁻¹·cm⁻¹)

EA: (%): calc'd. (found) for C₂₆H₂₆N₄O₈Co: C 51.02 (51.39); H 4.36 (4.28); N 8.98 (8.82).
- 0.5 DCM found within lattice for elemental analysis

4.7.10 Chelation of **4j** to a Cobalt metal to obtain **9j**

A round-bottom flask was charged with **4j** (0.503 g, 3.24 mmol), NEt₃ (1.00 mL, 0.726 g, 7.17 mmol), and EtOH (10 mL). Afterwards, Co(NO₃)₂·6 H₂O (0.435 g, 150 mmol), a bright pink-coloured compound, was dissolved in EtOH (5 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a bright pink. The solution was stirred at room temperature for 1 hour. Following completion of the reaction, bright pink precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (300 mg, 57 % yield).

Melting Point Range: >210 °C (decomp.)

IR: (cm⁻¹) 2964, 1577, 1508, 1405, 1351, 1297, 1221, 1190, 1165, 1045, 991, 932, 760.

UV-Vis: (DCM: 1.09×10^{-5} M) $\lambda_{\text{max}1} = 536 \text{ nm}$ ($\epsilon = 786 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}2} = 278 \text{ nm}$ ($\epsilon = 1.52 \times 10^5 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$)

EA: (%): calc'd. (found) for C₁₆H₂₄N₂O₄Co: C 52.32 (52.07); H 6.59 (6.37); N 7.63 (7.42).

4.7.11 Chelation of **4k** to a Cobalt metal to obtain **9k**

A round-bottom flask was charged with **4k** (0.502 g, 2.40 mmol) and EtOH (5 mL). After the transparent solids were completely dissolved, NEt₃ (1.00 mL, 0.726 g, 7.17 mmol) was added. Afterwards, Co(NO₃)₂·6 H₂O (0.317 g, 1.09 mmol), a bright pink-coloured compound, was dissolved in EtOH (5 mL) and added to the solution. Upon addition of the two, the solution turned colour from transparent to a bright pink. The solution was stirred at room temperature for 18 hours. Following completion of the reaction, bright pink precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (478 mg, 93 % yield).

Melting Point Range: 220 °C (decomp.)

IR: (cm⁻¹) 2975, 1608, 1560, 1463, 1360, 1314, 1250, 1173, 1126, 1028, 992, 971, 872, 751, 749, 699.

UV-Vis: (solv: 1.94 × 10⁻⁵M) λ_{max1} = 534 nm (ε = 389 L·mol⁻¹·cm⁻¹), λ_{max2} = 273 nm (ε = 7.38 × 10⁴ L·mol⁻¹·cm⁻¹)

EA: (%): calc'd. (found) for C₁₆H₁₈F₆N₂O₄Co: C 40.44 (40.54); H 3.82 (3.77); N 5.89 (5.98).

4.7.12 Chelation of **4I** to a Cobalt metal to obtain **9I**

A round-bottom flask was charged with **4I** (0.500 g, 1.93 mmol) and EtOH (5 mL). After the transparent solids were completely dissolved, NEt₃ (1.00 mL, 0.726 g, 7.17 mmol) was added. Afterwards, Co(NO₃)₂·6 H₂O (0.268 g, 0.920 mmol), a bright pink-coloured compound, was dissolved in EtOH (5 mL) and added to the solution. Upon addition of the two, the solution turned colour from transparent to a bright pink. The solution was stirred at room temperature for 18 hours. Following completion of the reaction, pale pink precipitates were collected which were recrystallized in a mixture of 1:1 DCM/MeOH (500 mg, 95 % yield).

Melting Point Range: 184 °C (decomp.)

IR: (cm⁻¹) 2970, 2895, 1596, 1524, 1458, 1352, 1281, 1199, 1031, 1005, 960, 936, 828, 802, 758, 671

UV-Vis: (solv: 2.10 × 10⁻⁴M) λ_{max1} = 536 nm (ε = 72.0 L·mol⁻¹·cm⁻¹), λ_{max2} = 489 nm (ε = 51.0 L·mol⁻¹·cm⁻¹), λ_{max3} = 285 nm (ε = 1.56 × 10⁴ L·mol⁻¹·cm⁻¹)

EA: (%): calc'd. (found) for C₁₆H₁₈F₆N₂O₄Co: C 40.44 (33.48); H 3.82 (3.77); N 5.89 (5.98).

4.8 Synthesis and Characterization of **Co** complexes made with **1a**

4.8.1 Chelation of Cobalt metal to obtain **10a**

A round-bottom flask was charged with compound **6a** (0.764 g, 4.04 mmol) and MeOH (15 mL). After the pinkish solid completely dissolved $\text{CoCl}_2 \cdot 6 \text{H}_2\text{O}$ (0.598 g, 2.51 mmol) was added to the solution. Upon addition of the two compounds the solution turned colour from pink to a wine red. The solution was stirred at room temperature for 3 hours. Upon completion of the reaction a tan-coloured precipitate was collected *via* vacuum filtration (0.587 g). The crude complex was recrystallized from a 1:1 mixture of warm CHCl_3 and diffusing with room temperature diethyl ether. The pure complex was obtained as a bright red solid (0.35 g, 39 % yield).

Melting Point Range: 240°C (decomp.)

IR: (cm^{-1}) 2968, 2903, 1591, 1569, 1527, 1486, 1418, 1354, 1261, 1192, 1108, 989, 945, 865, 746, 696, 668

UV-Vis: (DCM: $9.19 \times 10^{-5} \text{M}$) $\lambda_{\text{max}1} = 527 \text{ nm}$ ($\epsilon = 58.0 \text{ L} \cdot \text{mol}^{-1} \cdot \text{cm}^{-1}$), $\lambda_{\text{max}2} = 311 \text{ nm}$ ($\epsilon = 1.63 \times 10^4 \text{ L} \cdot \text{mol}^{-1} \cdot \text{cm}^{-1}$), $\lambda_{\text{max}3} = 282 \text{ nm}$ ($\epsilon = 1.52 \times 10^4 \text{ L} \cdot \text{mol}^{-1} \cdot \text{cm}^{-1}$), $\lambda_{\text{max}4} = 236 \text{ nm}$ ($\epsilon = 1.82 \times 10^4 \text{ L} \cdot \text{mol}^{-1} \cdot \text{cm}^{-1}$).

EA: (%): calc'd. (found) for $\text{C}_{22}\text{H}_{20}\text{N}_2\text{O}_4\text{Co}$: C 60.70 (60.49); H 4.63 (4.70); N 6.43 (6.49).

MS: (m/z) found (calc'd) $[\text{M}+\text{H}]^+ = 436.1$ (435.3)

4.8.1.1 Oxidation of **10a** to **11a**

A round-bottom flask was charged with compound **6a** (0.764 g, 4.04 mmol) and MeOH (15 mL). After the pink-solids completely dissolved $\text{CoCl}_2 \cdot 6 \text{H}_2\text{O}$ (0.598 g, 2.51 mmol) was added to the solution. Upon addition of the two compounds the solution turned colour from pink to a black-red. The solution was stirred at room temperature for 3 hours. Upon completion of the reaction a tan-coloured precipitate was collected *via* vacuum filtration (0.114 g). The crude complex (0.034 g) was recrystallized from a 1:1 mixture of hot DCM and layering on top cool MeOH, which led to oxidation of the metal forming the octahedral complex **11a** (0.032 mg). The pure complex was obtained as a blackish-green solid (49 %) and structurally determined *via* x-ray crystallography.

Melting Point Range: 235°C (decomp.)

IR: (cm^{-1}) 2921, 2108, 1612, 1526, 1478, 1454, 1420, 1353, 1317, 1266, 186, 1109, 1066, 990, 948, 871, 791, 736, 695

UV-Vis: (DCM: $5.77 \times 10^{-6}\text{M}$) $\lambda_{\text{max}1} = 556 \text{ nm}$ ($\epsilon = 556 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}2} = 441 \text{ nm}$ ($\epsilon = 1.31 \times 10^3 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}3} = 347 \text{ nm}$ ($\epsilon = 2.28 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}4} = 268 \text{ nm}$ ($\epsilon = 8.99 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}5} = 262 \text{ nm}$ ($\epsilon = 9.72 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}6} = 258 \text{ nm}$ ($\epsilon = 9.37 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}7} = 245 \text{ nm}$ ($\epsilon = 8.71 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$).

EA: (%): calc'd. (found) for $\text{C}_{33}\text{H}_{30}\text{N}_3\text{O}_6\text{Co}$: C 60.41 (61.05); H 4.96 (4.69); N 6.31 (5.99).

- 0.5 DCM found within lattice for elemental analysis

4.8.2 Chelation of Cobalt metal to obtain **10b**

A round-bottom flask was charged with **6b** (42 mg, 0.16 mmol) and EtOH (2 mL). After the yellow-coloured solids were completely dissolved, Co(NO₃)₂·6 H₂O (23 mg, 0.079 mmol) dissolved in EtOH (3 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark orange. NEt₃ (0.1 mL, 73 mg, 72 mmol) was added to the reaction. The solution was stirred at room temperature for 1 hour. Upon completion of the reaction a tan-coloured precipitate was collected *via* vacuum filtration. The crude complex was recrystallized from a 1:1 mixture of warm CHCl₃ and diffusing with room temperature diethyl ether. The pure complex was obtained as a bright red solid (36 mg, 79 % yield).

Melting Point Range: 221 °C (decomp.)

IR: (cm⁻¹) 1592, 1529, 1462, 1437, 1359, 1280, 1240, 1194, 1134, 1107, 1058, 1025, 995, 949, 899, 801, 761, 677

UV-Vis: (DCM: 7.00 × 10⁻⁵M) λ_{max1} = 524 nm (ε = 94.0 L·mol⁻¹·cm⁻¹), λ_{max2} = 321 nm (ε = 2.11 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max3} = 296 nm (ε = 1.51 × 10⁴ L·mol⁻¹·cm⁻¹), λ_{max4} = 243 nm (ε = 1.91 × 10⁴ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₂₂H₁₆Cl₄N₂O₄Co: C 43.90 (44.43); H 2.75 (2.78); N 4.55 (4.66).
- 0.5 DCM found within lattice for elemental analysis

4.8.3 Chelation of Cobalt metal to obtain **10c**

A round-bottom flask was charged with **6c** (0.630 g, 2.87 mmol) and EtOH (10 mL). After the yellow-coloured solids were completely dissolved, Co(NO₃)₂·6 H₂O (0.412 g, 1.42 mmol) dissolved in EtOH (10 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark orange. NEt₃ (0.60 mL, 0.44 g, 4.3 mmol) was added to the reaction. The solution was stirred at room temperature for 1 hour. Upon completion of the reaction a tan-coloured precipitate was collected *via* vacuum filtration. The crude complex was recrystallized from a 1:1 mixture of warm DCM and diffusing with room temperature diethyl ether. The pure complex was obtained as a bright red solid (595 mg, 85 % yield).

Melting Point Range: 250 °C (decomp.)

IR: (cm⁻¹) 2960, 1587, 1566, 1531, 1498, 1396, 1358, 1321, 1301, 1245, 1194, 1170, 1113, 1056, 1028, 992, 948, 869, 838, 770, 672

UV-Vis: (DCM: 1.94×10^{-5} M) $\lambda_{\max 1} = 547$ nm ($\epsilon = 73.0$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 2} = 317$ nm ($\epsilon = 4.04 \times 10^4$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 3} = 263$ nm ($\epsilon = 2.32 \times 10^4$ L·mol⁻¹·cm⁻¹), $\lambda_{\max 4} = 242$ nm ($\epsilon = 2.06 \times 10^4$ L·mol⁻¹·cm⁻¹).

EA: (%): calc'd. (found) for C₂₄H₂₄N₂O₆Co: C 49.18 (49.12); H 4.37 (4.80); N 4.50 (5.17).
- 1.5 DCM found within lattice for elemental analysis

MS: (m/z) found (calc'd) [M+H]⁺ = 495.4 (496.2)

4.8.3.1 Oxidation of **10c** to **11c**

A round-bottom flask was charged with **6c** (0.630 g, 2.87 mmol) and EtOH (10 mL). After the yellow-coloured solids were completely dissolved, $\text{Co}(\text{NO}_3)_2 \cdot 6 \text{H}_2\text{O}$ (0.412 g, 1.42 mmol) dissolved in EtOH (10 mL) and added to the solution. Upon addition of the two, the solution turned colour from yellow to a dark orange. NEt_3 (0.60 mL, 0.43 g, 4.30 mmol) was added to the reaction. The solution was stirred at room temperature for 1 hour. Upon completion of the reaction a tan-coloured precipitate was collected *via* vacuum filtration (595 mg, 85 % yield). The precipitates (40 mg) were dissolved in a vial containing only DCM, which led to oxidation of the metal forming the octahedral complex **11c**. The pure complex was obtained as a blackish-green crystalline solid (50 mg, 86 % yield.).

Melting Point Range: >230 °C (decomp.)

IR: (cm^{-1}) 2923, 1599, 1529, 1495, 1437, 1358, 1319, 1299, 1244, 1167, 1111, 1071, 126, 992, 950, 876, 837, 756, 723

UV-Vis: (DCM: $5.61 \times 10^{-6}\text{M}$) $\lambda_{\text{max}1} = 553 \text{ nm}$ ($\epsilon = 683 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}2} = 429 \text{ nm}$ ($\epsilon = 1.34 \times 10^3 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}3} = 343 \text{ nm}$ ($\epsilon = 2.06 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}4} = 276 \text{ nm}$ ($\epsilon = 6.38 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}5} = 269 \text{ nm}$ ($\epsilon = 7.43 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}6} = 265 \text{ nm}$ ($\epsilon = 7.15 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$), $\lambda_{\text{max}7} = 244 \text{ nm}$ ($\epsilon = 4.46 \times 10^4 \text{ L}\cdot\text{mol}^{-1}\cdot\text{cm}^{-1}$)

EA: (%): calc'd. (found) for $\text{C}_{36}\text{H}_{36}\text{N}_3\text{O}_9\text{Co}$: C 59.47 (59.84); H 5.20 (5.54); N 5.78 (5.04).
- 0.75 H_2O found within lattice for elemental analysis

4.9 Attempted Oxidation of **Co** complexes made with **1a**

4.9.1 Attempted oxidation of **9a** to **9aX**;

For quantities and general procedure, see below;

To a 25-mL round-bottom flask, **9a** and solvent were added. X_2 was added to the flask. The contents of the flask were stirred at room temperature for (mins). At the termination of the reaction, the solvent was removed *in vacuo*.

For all bromine reactions, removal of the excess Br_2 revealed a bright blue-green gum. The dried gum was insoluble in most common laboratory solvents with the exclusion of acetone and water. Once dissolved the solution turned from a teal colour to a bright blue. NMR spectral analysis turned no positive results as the lack of any ligand signals. Dissolving commercially available $CoBr_2 \cdot 6H_2O$ and the synthesized product in water confirmed the formation of $CoBr_2$ occurred.

For all iodine reactions, the resultant red power was vacuum filtered, washed with cold methanol then left to dry. Further analysis revealed the persistency of the starting material **9a** with absence of any iodine. [See **Sec. 2.5.3.1** for full discussion]

Table 15: Trials for oxidative addition of halogens to **9a**

Trial	$Co(L_2)_2$ (mg)	X_2	Molar Equiv. ($X_2:Co$)	Solvent	Time (mins)	light? (Y/N)
1	251	Br	1:1	neat	60	Y
2	507	Br	1:2	neat	15	N
3	108	I	1:1	toluene	60	Y
4	105	I	1:1	acetone	60	Y

4.9.2 Attempted oxidation of **9h** to **9h-1** and then **9h-2**

In a 100-mL round-bottom flask, **9h** (445 mg, 0.986 mmol), a bright pink compound, was dissolved in acetonitrile (20 mL). Upon mixing, the solution turned to a dark purple suggesting coordination of either one or two ACN molecules. To this, bromine (0.070 mL, 0.22 g, 1.3 mmol) was added dropwise. The solution was stirred at room temperature in the dark (wrapped in aluminum foil) for 1 hour. After completion of the reaction, volatile components were removed *in vacuo*. What was left behind was a green resin. This green resin is water soluble and insoluble in most organic solvents. Thus, it was determined not to be the target complex.

4.10 Using **9h** in ATRP

4.10.1 Polymerization of MMA with **9h** to PMMA

Into a flame dried 10-mL Schlenk flask, a mixture of dry solvent (3 mL), TsCl (0.3 mol %) and **9h** (2 mol %) were added. To the flask, MMA (1.70mL, 1.60 g, 16.0 mmol) was added. The mixture of materials was sparged for 30 minutes with N_{2(g)}. The reaction was left to stir at 60 °C for 24 hours. After completion of the reaction, the solvent was removed *in vacuo*. The polymer was diluted with a small amount of THF (~1 mL) and dropped into rapidly stirring -26 °C MeOH. This cleaning procedure was repeated at 3-4 times to ensure complete removal of all starting materials. The catalyst (**9h**) was quantitatively recovered by removal of MeOH by rotary evaporator. PMMA was isolated by removing excess solvent (MeOH/THF) *in vacuo* till dry (658 mg, 42 % conversion). GPC analysis revealed a PDI of 1.19 and a molecular weight for PMMA of 50,000 Da. Conversions are similar to other Co catalysts.⁷⁴

¹H-NMR: (400 MHz, CDCl₃) δ **0.80-0.97** (m, 3H, **2-syndio/isotactic**), **1.38** (m, 2H, **1-isotactic**), **1.76-1.92** (m, 2H, **1-syndiotactic**), **3.54** (m, 3H, **5-syndiotactic**), **3.69** (m, 3H, **5-isotactic**)

¹³C-NMR: (101 MHz, CDCl₃) δ 16.5 (**2-syndio**); 18.7 (**2-syndio**); 25.6 (**1-iso**); 30.3 (**1-syndio**); 44.5 (**3-iso**); 44.9 (**3-syndio**); 51.8 (**5-syndio**); 54.1 (**1-iso**); 54.4 (**1-syndio**); 67.8 (**5-iso**); 176.9 (**4-iso**); 177.7 (**4-syndio**);

4.10.2 Attempted polymerization of THF with **9h**

Into a flame dried 10-mL Schlenk flask, a mixture of dry THF (5 mL), TsCl (11 mg, 0.058 mmol) and **9h** (101 mg, 0.226 mmol) were added. The mixture of materials was sparged for 30 minutes with N_{2(g)}. The reaction was left to stir at 60 °C, under static vacuum, for 24. After the completion of the reaction, purification yielded no trace of poly(THF).

Trial	Mol% cat.	Solvent	Time	Temperature (°C)	Initiator : mol %	Mw : PDI	% Conversion
1	65	THF	24	60 (static vac.)	TsCl : 16	N/A	0

4.11 Synthesis of Other Group 9 Complexes

4.11.1 Synthesis of Vaska's complex ($\text{IrCl}(\text{CO})(\text{PPh}_3)_2$) (**12**)

Analyses of this compound are consistent with those in the literature.⁸⁹

^{31}P -NMR: (161 MHz, C_6D_6) δ +24 ppm, +5 ppm

^1H -NMR: (400 MHz, CDCl_3) δ 7.36-7.40 (m, 18H), 7.72-7.73 (m, 12H)

TLC: 20 % EtOAc/*n*-hexanes R_f = 0.29

IR: (cm^{-1}) 2109, 1948($\nu = \text{CO}$), 1477, 1433, 1306, 1181, 1095, 1027, 997, 844, 744, 689.

4.11.2 Synthesis of bis(1,5-cyclooctadiene)diiridium(I)dichloride (**14**)

Analyses of this compound are consistent with those in the literature.⁵⁶

^1H -NMR: (400 MHz, CDCl_3) δ **1.52** (q, J = 8.0 Hz, 4H), **2.24-2.26** (m, 4H), **4.22** (s, 4H)

^{13}C -NMR: (101 MHz, CDCl_3) δ 31.9; 62.3

4.11.3 Substitution of **14** with **4a** to give **15**

A 50-mL three-necked round-bottom flask was charged with both **14** (0.608 mg, 0.905 mmol) and **4a** (0.512 mg, 2.36 mmol). This flask was purged and back filled with an N_{2(g)} atmosphere. To the flask, toluene (6 mL) and NEt₃ (1.0 mL, 0.726 g, 7.2 mmol) were added. As soon as the base was added, the reaction mixture immediately changed colour from dark red-orange to a bright yellow. The reaction was left to stir at room temperature for 1 hour under a constant nitrogen atmosphere. Once completed, the solvent mixture was removed *in vacuo* leaving behind a yellow powder. The powder was washed with ice cold MeOH (~5 mL × 5) and the solution decanted from the yellow solids. The solids were dried over vacuum after being washed one last time with ice cold MeOH. The fine yellow powder was determined to be the chelated product, **15** (894 mg, 90 % yield).

Melting Point: >150 °C (decomp.)

TLC: 30 % EtOAc/*n*-hexanes R_f = 0.30

IR: (cm⁻¹) 2988, 2927, 2877, 2832, 1586, 1560, 1533, 1486, 1357, 1315, 1255, 1196, 1176, 1155, 1114, 1019, 969, 936, 907, 866, 834, 798, 774, 738, 696.

¹H-NMR: (400 MHz, CDCl₃) δ **1.40** (s, 6H, **1**), **1.50-1.58** (m, 2H, **12,13**), **1.69-1.76** (m, 2H, **12,13**), **2.16-2.25** (m, 2H, **12,13**), **2.29-2.39** (m, 2H, **12,13**), **3.89** (s, 2H, **2**), **4.27** (t, *J* = 3.0 Hz, 2H, **11,14**), **4.35** (t, *J* = 3.0 Hz, 2H, **11,14**), **5.59** (s, 1H, **5**), **7.30-7.36** (m, 3H, **9,10**), **7.75-7.78** (m, 2H, **8**),

¹³C-NMR: (101 MHz, CDCl₃) δ 27.5 (**1**); 28.5 (**12-13**); 33.5 (**12-13**); 53.0 (**11,14**); 65.7 (**11,14**); 72.0 (**3**); 79.4 (**2**); 81.7 (**5**); 126.7 (**8**); 128.1 (**9**); 129.8 (**10**); 138.6 (**7**); 168.6 (**4**); 176.5 (**6**)

EA: (%): calc'd. (found) for C₂₁H₂₆NO₂Ir: C 48.82 (49.11); H 5.07 (5.20); N 2.71 (2.79).

4.11.4 Oxidative addition of methyl iodide to **15**

In a 25-mL round-bottom flask, dry toluene (10 mL) was added to dissolve **15** (201 mg, 0.368 mmol). Once fully dissolved, MeI (0.11 mL, 1.8 mmol) was added dropwise. The solution was stirred under an atmosphere of N_{2(g)} for 24 hours. The yellow coloured solution turned to a darker orange. Upon completion of the reaction, excess toluene was removed *via* a rotary evaporator where the mixture solidified. To purify, the crude solids were dissolved in a minimal amount of toluene and dropped into -26 °C *n*-hexanes. This revealed the product as light brown solids which then the target product was isolated *via* vacuum filtration. The solids were recrystallized from a mixture of 1:1 C₆D₆/*n*-hexanes to give brown spherical plates (101 mg, 42 % yield).

TLC: 30 % EtOAc/*n*-hexanes R_f = 0.33

IR: (cm⁻¹) 2925, 2870, 1591, 1543, 1486, 1448, 1361, 1307, 1249, 1190, 1117, 1068, 1037, 937, 870, 839, 776, 726, 697.

¹H-NMR: (400 MHz, CDCl₃) δ **1.38** (d, *J* = 18.4 Hz, 6H, **1,1a**), **1.65-1.74** (m, 1H, **(11/14)a**), **2.03-2.11** (m, 1H, **(11/14)a**), **2.22-2.28** (m, 1H, **(11/14)a**), **2.33** (s, 3H, **15**), **2.42-2.51** (m, 2H, **(11/14)a**), **2.62-2.72** (m, 1H, **(11/14)a**), **3.01-3.10** (m, 1H, **(11/14)a**), **3.27-3.37** (m, 1H, **(11/14)a**), **3.89** (ABq, *J* = 8.0, 2H, **2,2a**), **4.43** (td, *J* = 3.8, 8.5 Hz, 1H, **(12-13)a**), **4.70-4.73** (m, 1H, **(12-13)a**), **5.46-5.50** (m, 2H, **5** and **(12-13)a**), **5.71** (td, *J* = 3.4, 8.4 Hz, 1H **(12-13)a**), **7.29-7.34** (m, 3H, **9,10**), **7.64-7.66** (m, 2H, **8**)

¹³C-NMR: (101 MHz, CDCl₃) δ 11.3 (**15**); 25.1 (**(11/14)a**); 25.3 (**1,1a**); 28.6 (**1,1a**); 31.6 (**(11/14)a**); 32.1 (**(11/14)a**); 34.7 (**(11/14)a**); 71.2 (**(12-13)a**); 71.6 (**3**); 74.0 (**(12-13)a**); 79.6 (**2,2a**); 82.4 (**5**); 86.9 (**(12-13)a**); 88.0 (**(12-13)a**); 126.7 (**8**); 128.0 (**9**); 129.8 (**10**); 138.7 (**7**); 165.6 (**4**); 174.5 (**6**)

EA: (%): calc'd. (found) for C₂₂H₂₉INO₂Ir: C 40.12 (39.99); H 4.44 (4.50); N 2.13 (2.11).

4.11.5 Synthesis and oxidative addition of triphenylstannane to **15**

In the glovebox, Et₂O (100 mL) was combined with ClSnPh₃ (7.261 g, 18.84 mmol) in a Schlenk flask and cooled to 0 °C. LiAlH₄ (18.00 mL, 18.00 mmol) was added and stirred at 0 °C for 4 hours. The reaction was quenched with degassed dH₂O (50 mL). The organic components were extracted, dried (MgSO₄), filtered and all volatile components were removed *in vacuo* to reveal HSnPh₃ (6.61 g, 94 % yield). In a 25-mL round-bottom flask in the glovebox, HSnPh₃ (149 mg, 0.425 mmol) and **15** (100 mg, 0.183 mmol) were weighed out and dissolved in C₆D₆ (1.66 mL). The mixture was stirred at room temperature under a constant nitrogen atmosphere for 20 hours. The mixture is concentrated under vacuum, DCM (0.5 mL) is added to dissolve the complex then dry *n*-hexanes (10 mL) was added. The complex, **18**, is collected by vacuum filtration as a yellow powder (128 mg, 80 % yield).

IR: (cm⁻¹) 2956, 2884, 1600, 1574, 1537, 1424, 1359, 1307, 1254, 1191, 1112, 1064, 1011, 922, 861, 726, 693

¹H-NMR: (400 MHz, C₆D₆) δ -17.40 (s, *J*_{Sn-H} = 23.0 Hz, 1H, **19**), 0.85 (d, *J* = 34.0 Hz, 6H, **1,1a**), 1.01-1.12 (m, 2H, **(12-13)a**), 1.58-1.68 (m, 2H, **(12-13)a**), 2.00-2.08 (m, 1H, **(12-13)a**), 2.15-2.35 (m, 2H, **(12-13)a**), 2.75-2.80 (m, 1H, **(12-13)a**), 3.11 (ABq, *J* = 7.8 Hz, 2H, **2,2a**), 4.52-4.57 (m, 1H, **(11-14)a**), 4.74-4.78 (m, 2H, **(11-14)a**), 4.99-5.08 (m, 2H, **(11-14)a**), 5.74 (s, 1H, **5**), 7.18-7.25 (m, 12H, **9-10/17-18**), 7.88-7.98 (m, 6H, **16**), 8.04-8.05 (m, 2H, **8**),

¹³C-NMR: (101 MHz, C₆D₆) δ 26.7 (**(12-13)a**); 27.2 (**(12-13)a**); 27.3 (**1/1a**); 27.7 (**1/1a**); 28.8 (**(12-13)a**); 41.2 (**(12-13)a**); 68.3 (**3**); 70.0 (**(11/14)a**); 76.0 (**(11/14)a**); 76.8 (**2/2a**); 80.6 (**5**); 95.3 (**(11/14)a**); 99.8 (**(11/14)a**); 126.5 (**8**); 127.6 (**9**); 127.8 (**17/18**); 128.2 (**10**); 129.5 (**17/18**); 137.7 (*J*_{Sn-C} = 17.0 Hz, **(16)**); 139.9 (**15**); 144.1 (**7**); 162.8 (**4**); 175.6 (**6**);

¹¹⁹Sn-NMR: (149 MHz, C₆D₆) δ -178.1, (149 MHz, CDCl₃) δ

EA: (%): calc'd. (found) for C₃₉H₄₂SnNO₂Ir: C 52.13 (52.18); H 4.76 (4.58); N 1.54 (1.41).

Appendix

Appendix A: Nuclear Magnetic Resonance Spectroscopy (NMR)

A1: ^1H -NMR of 6a in CDCl_3^*	158
A2: ^{13}C -NMR of 6a in CDCl_3^*	159
A3: COSY NMR of 6a in CDCl_3	160
A4: HSQC NMR of 6a in CDCl_3	161
A5: ^1H -NMR of 3a in CDCl_3^*	162
A6: ^{13}C -NMR of 3a in CDCl_3^*	163
A7: ^1H -NMR of 4a in CDCl_3^*	164
A8: ^1H -NMR of 4a in C_6D_6	165
A9: ^{13}C -NMR of 4a in CDCl_3^*	166
A10: ^1H -NMR of 3b in CDCl_3^*	167
A11: ^{13}C -NMR of 3b in CDCl_3^*	168
A12: ^1H -NMR of 4b in CDCl_3^*	169
A13: ^{13}C -NMR of 4b in CDCl_3^*	170
A14: COSY NMR of 4b in CDCl_3	171
A15: HSQC NMR of 4b in CDCl_3	172
A16: ^1H -NMR of 3c in CDCl_3^*	173
A17: ^{13}C -NMR of 3c in CDCl_3^*	174
A18: ^1H -NMR of 4c in CDCl_3^*	175
A19: ^{13}C -NMR of 4c in CDCl_3^*	176
A20: COSY NMR of 4c in CDCl_3	177
A21: HSQC NMR of 4c in CDCl_3	178
A22: ^1H -NMR of 3d in CDCl_3^*	179

A23: ^{13}C -NMR of 3d in CDCl_3^*	180
A24: ^1H -NMR of 4d in CDCl_3^*	181
A25: ^{13}C -NMR of 4d in CDCl_3^*	182
A26: COSY NMR of 4d in CDCl_3	183
A27: HSQC NMR of 4d in CDCl_3	184
A28: ^1H -NMR of 3e in CDCl_3^*	185
A29: ^{13}C -NMR of 3e in CDCl_3^*	186
A30: ^1H -NMR of 4e in CDCl_3^*	187
A31: ^{13}C -NMR of 4e in CDCl_3^*	188
A32: COSY NMR of 4e in CDCl_3	189
A33: HSQC NMR of 4e in CDCl_3	190
A34: ^1H -NMR of 3f in CDCl_3^*	191
A35: ^{13}C -NMR of 3f in CDCl_3^*	192
A36: ^1H -NMR of 4f in CDCl_3^*	193
A37: ^{13}C -NMR of 4f in CDCl_3^*	194
A38: COSY NMR of 4f in CDCl_3	195
A39: HSQC NMR of 4f in CDCl_3	196
A42: ^1H -NMR of 4g in CDCl_3^*	197
A43: ^{13}C -NMR of 4g in CDCl_3^*	198
A44: COSY NMR of 4g in CDCl_3	199
A45: HSQC NMR of 4g in CDCl_3	200
A46: ^1H -NMR of 6b in CDCl_3^*	201
A47: ^{13}C -NMR of 6b in CDCl_3^*	202

A48: COSY NMR of 6b in CDCl ₃	203
A49: HSQC NMR of 6b in CDCl ₃ *	204
A50: ¹ H-NMR of 6c in CDCl ₃ *	205
A51: ¹³ C-NMR of 6c in CDCl ₃ *	206
A52: COSY NMR of 6c in CDCl ₃	207
A53: HSQC NMR of 6c in CDCl ₃	208
A54: ¹ H-NMR of 3h in CDCl ₃ *	209
A55: ¹³ C-NMR of 3h in CDCl ₃ *	210
A56: ¹ H-NMR of 4h in CDCl ₃ *.....	211
A57: ¹³ C-NMR of 4h in CDCl ₃ *	212
A58: COSY NMR of 4h in CDCl ₃	213
A59: HSQC NMR of 4h in CDCl ₃	214
A60: ¹ H-NMR of 4j in CDCl ₃ *	215
A61: ¹³ C-NMR of 4j in CDCl ₃ *	216
A62: COSY NMR of 4j in CDCl ₃	217
A63: HSQC-NMR of 4j in CDCl ₃	218
A64: ¹ H-NMR of 4k in CDCl ₃ *	219
A65: ¹³ C-NMR of 4k in CDCl ₃ *	220
A66: ¹⁹ F-NMR of 4k in CDCl ₃ *	221
A67: COSY NMR of 4k in CDCl ₃	222
A68: HSQC NMR of 4k in CDCl ₃	223
A69: ¹ H-NMR of 4l in CDCl ₃ *	224
A70: ¹³ C-NMR of 4l in CDCl ₃ *	225

A71: COSY NMR of 41 in CDCl ₃	226
A72: HSQC NMR of 41 in CDCl ₃	227
A73: ¹ H-NMR of 14 in CDCl ₃ *	228
A74: ¹³ C-NMR of 14 in CDCl ₃ *	229
A75: ¹ H-NMR of 15 in CDCl ₃ *	230
A76: ¹³ C-NMR of 15 in CDCl ₃ *	231
A77: NOESY NMR of 15 in CDCl ₃	232
A78: COSY NMR of 15 in CDCl ₃	233
A79: HSQC NMR of 15 in CDCl ₃	234
A80: ¹ H-NMR of 16 in CDCl ₃ *	235
A81: ¹³ C-NMR of 16 in CDCl ₃ *	236
A82: COSY NMR of 16 in CDCl ₃	237
A83: NOESY NMR of 16 in CDCl ₃	238
A84: HSQC NMR of 16 in CDCl ₃	239
A85: ¹ H-NMR of HSnPh₃ in C ₆ D ₆ *	240
A86: ¹¹⁹ Sn-NMR of HSnPh₃ in C ₆ D ₆ *	241
A87: ¹ H-NMR of 18 in C ₆ D ₆ *	242
A88: ¹³ C-NMR of 18 in C ₆ D ₆ *	243
A89: ¹¹⁹ Sn-NMR of 18 in C ₆ D ₆	244
A90: COSY NMR of 18 in C ₆ D ₆	245
A91: NOESY NMR of 18 in C ₆ D ₆	246
A92: HSQC NMR of 18 in C ₆ D ₆	247
A94: ¹ H-NMR of PMMA in CDCl ₃ *	248

A95: ^{13}C -NMR of PMMA in CDCl_3^*	249

A1: ¹H-NMR of **6a** in CDCl₃*

851

A2: ¹³C-NMR of **6a** in CDCl₃*

159

A3: COSY NMR of **6a** in CDCl₃

160

A4: HSQC NMR of **6a** in CDCl₃

A5: ¹H-NMR of **3a** in CDCl₃*

A6: ¹³C-NMR of **3a** in CDCl₃*

A7: ¹H-NMR of **4a** in CDCl₃*

A8: ¹H-NMR of **4a** in C₆D₆

KM-03-25
Ligand-2
13C in CDCl3

—187.33

—169.56

—140.00

—130.55

—128.14

—126.82

—78.90

—73.95

—58.51

—27.05

165

A9: ^{13}C -NMR of **4a** in CDCl_3^*

KM-03-33
Int-3
1H in CDCl₃

A10: ¹H-NMR of **3b** in CDCl₃*

L91

A11: ¹³C-NMR of **3b** in CDCl₃*

A12: ¹H-NMR of **4b** in CDCl₃*

A13: ^{13}C -NMR of **4b** in CDCl_3 *

170

A14: COSY NMR of **4b** in CDCl₃

A15: HSQC NMR of **4b** in CDCl₃

172

KM-03-118
Int-4
1H in CDCl₃

A16: ¹H-NMR of 3c in CDCl₃*

A17: ¹³C-NMR of 3c in CDCl₃*

A18: ¹H-NMR of **4c** in CDCl₃*

KM-03-43
Ligand-4
13C in CDCl₃

—186.81

—169.40

—161.66

—132.68

—128.63

—113.34

—78.87

—73.31

—58.44

—55.31

—27.11

175

A19: ¹³C-NMR of 4c in CDCl₃*

A20: COSY NMR of **4c** in CDCl₃

177

A21: HSQC NMR of **4c** in CDCl₃

A22: ¹H-NMR of **3d** in CDCl₃*

A24: ¹H-NMR of **4d** in CDCl₃*

KM-03-48
Ligand-5
 ^{13}C in CDCl_3

—187.67

—169.46

—139.53

—137.69

—136.26

—129.44

—128.12

—124.39

—78.88

—73.79

—58.48

—27.13

—19.87

—19.80

A25: ^{13}C -NMR of **4d** in CDCl_3 *

A26: COSY NMR of **4d** in CDCl₃

183

KM-03-48
Ligand-5
HSQC in CDCl₃

A27: HSQC NMR of **4d** in CDCl₃

A28: ¹H-NMR of **3e** in CDCl₃*

A29: ¹³C-NMR of **3e** in CDCl₃*

A30: ^1H -NMR of **4e** in CDCl_3 *

A32: COSY NMR of **4e** in CDCl₃

189

A33: HSQC NMR of **4e** in CDCl₃

KM-03-69
Int-7
1H in CDCl3

0.6 T

A34: ¹H-NMR of 3f in CDCl₃*

A35: ¹³C-NMR of 3f in CDCl₃*

A36: ¹H-NMR of **4f** in CDCl₃*

A37: ¹³C-NMR of **4f** in CDCl₃*

194

A38: COSY NMR of **4f** in CDCl₃

A39: HSQC NMR of **4f** in CDCl₃

A40: ¹H-NMR of **4g** in CDCl₃*

A41: ^{13}C -NMR of **4g** in CDCl_3 *

A42: COSY NMR of **4g** in CDCl₃

A43: HSQC NMR of **4g** in CDCl₃

A44: ¹H-NMR of **6b** in CDCl₃*

KM-03-94
Ligand-9
 ^{13}C in CDCl_3

—184.69

—171.04

—139.88

—134.57

—132.50

—130.17

—128.99

—126.04

77.37

77.05

76.73

74.13

67.64

—43.07

A45: ^{13}C -NMR of **6b** in CDCl_3 *

A46: COSY NMR of **6b** in CDCl₃

A47: HSQC NMR of **6b** in CDCl₃*

A48: ¹H-NMR of **6c** in CDCl₃*

KM-03-93
 Ligand-10
¹³C in CDCl₃

—187.00

—170.53

—161.70

—132.67

—128.69

—113.36

—73.53

—67.32

—55.33

—43.06

8/12

9/11

*

5

2

13

1

6

3

7

10

A49: ¹³C-NMR of **6c** in CDCl₃*

A50: COSY NMR of **6c** in CDCl₃

A51: HSQC NMR of **6c** in CDCl₃

KM-03-132
Int-12
1H in CDCl₃

A52: ¹H-NMR of **3h** in CDCl₃*

A53: ^{13}C -NMR of **3h** in CDCl_3 *

A54: ¹H-NMR of **4h** in CDCl₃*

KM-03-105
Ligand-12
1H in CDCl₃

A55: ¹³C-NMR of **4h** in CDCl₃*

A56: COSY NMR of **4h** in CDCl₃

A57: HSQC NMR of **4h** in CDCl₃

KM-03-180
L14
1H in CDCl3

A58: ¹H-NMR of **4j** in CDCl₃*

A59: ¹³C-NMR of **4j** in CDCl₃*

A60: COSY NMR of **4j** in CDCl₃

KM-03-174
Ligand-15
1H in CDCl3

A62: ^1H -NMR of **4k** in CDCl_3 *

A63: ^{13}C -NMR of **4k** in CDCl_3 *

KM-03-174
Ligand-15
 ^{19}F in CDCl_3

-76.35

9

220

A64: ^{19}F -NMR of 4k in CDCl_3^*

A65: COSY NMR of **4k** in CDCl₃

A66: HSQC NMR of **4k** in CDCl₃

KM-03-175
Ligand-16
 ^1H in CDCl_3

A67: ^1H -NMR of **41** in CDCl_3^*

KM-03-175
Ligand-16
13C in CDCl₃

—181.10

—170.95

—97.18

—79.83

—69.84

—59.08

—27.18

A68: ¹³C-NMR of **41** in CDCl₃*

A69: COSY NMR of **4l** in CDCl₃

A70: HSQC NMR of **4I** in CDCl₃

KM-04-30
[IrCODCl]₂
1H in CDCl₃

A71: ¹H-NMR of **14** in CDCl₃*

KM-04-30
[IrCODCl]₂
¹³C in CDCl₃

—62.18

—31.78

*

1

2

A72: ¹³C-NMR of **14** in CDCl₃*

A73: ¹H-NMR of **15** in CDCl₃*

A74: ¹³C-NMR of **15** in CDCl₃*

A75: NOESY NMR of **15** in CDCl₃

A76: COSY NMR of **15** in CDCl₃

A77: HSQC NMR of **15** in CDCl₃

A78: ¹H-NMR of **16** in CDCl₃*

A79: ¹³C-NMR of **16** in CDCl₃*

KM-04-26
(COD)Ir(L2)MeI
COSY in CDCl₃

A80: COSY NMR of **16** in CDCl₃

KM-04-26
(COD)Ir(L2)MeI
NOESY in CDCl₃

A81: NOESY NMR of **16** in CDCl₃

KM-04-26
(COD)Ir(L2)MeI
HSQC in CDCl₃

A82: HSQC NMR of **16** in CDCl₃

A83: ¹H-NMR of HSnPh₃ in C₆D₆*

KM-04-32
H₃SnPh
¹¹⁹Sn in C₆D₆

-162.82

240

A84: ¹¹⁹Sn-NMR of H₃SnPh in C₆D₆*

KM-04-33
(COD)Ir(L2)HSnPh₃
13C in C₆D₆

A86: ¹³C-NMR of 18 in C₆D₆*

KM-04-33
(COD)Ir(L2)HSnPh₃
119Sn in C₆D₆

---178.05

A87: ¹¹⁹Sn-NMR of **18** in C₆D₆

A88: COSY NMR of **18** in C_6D_6

A89: NOESY NMR of **18** in C_6D_6

KM-04-33
(COD)Ir(L2)HSnPh3
HSQC in C6D6

A90: HSQC NMR of **18** in C₆D₆

A92: ¹³C-NMR of PMMA in CDCl₃*

Appendix B: Infrared Spectroscopy (IR)

B1: IR spectrum for 10a	264
B2: IR spectrum for 11a	267
B3: IR spectrum for 9a	252
B4: IR spectrum for 9b	253
B5: IR spectrum for 9c	254
B6: IR spectrum for 9d	255
B7: IR spectrum for 9e	256
B8: IR spectrum for 9f	257
B9: IR spectrum for 9g	258
B10: IR spectrum for 10b	265
B11: IR spectrum for 10c	266
B12: IR spectrum for 11c	268
B13: IR spectrum for 9h	259
B14: IR spectrum for 9i	260
B15: IR spectrum for 9j	261
B16: IR spectrum for 9k	262
B17: IR spectrum for 9l	263
B18: IR spectrum for 8a	269
B19: IR spectrum for 7a (brown)	270
B20: IR spectrum for 7a (green)	271
B21: IR spectrum for 7b	272
B22: IR spectrum for 7c	273

B23: IR spectrum for 7d	274
B24: IR spectrum for 7e	275
B25: IR spectrum for 7f	276
B26: IR spectrum for 7g	277
B27: IR spectrum for 8b	278
B28: IR spectrum for 7h	279
B29: IR spectrum for 7i	280
B30: IR spectrum for 12	281
B31: IR spectrum for 15	282
B32: IR spectrum for 16	283
B33: IR spectrum for 18	284

Agilent Resolutions Pro

Name
Co (oxme) 2 - L2

B1: IR spectrum for 9a

Agilent Resolutions Pro

252

Name
Co(Cl ₂) ₂ - L3

B2: IR spectrum for **9b**

Agilent Resolutions Pro

Name
Co (OMe) 2 - L4

Agilent Resolutions Pro

B4: IR spectrum for **9d**

Name
Co (Me ₂) ₂ - L5

Agilent Resolutions Pro

B5: IR spectrum for 9e

Co(pCl3Me)2 - L6

Agilent Resolutions Pro

B6: IR spectrum for 9f

Agilent Resolutions Pro

257

Name
Co(nap)2 - L8

B7: IR spectrum for 9g

Agilent Resolutions Pro

Name
Co (L12) 2

B8: IR spectrum for 9h

Agilent Resolutions Pro

259

Name
Co (L13) 2

B9: IR spectrum for 9i

Agilent Resolutions Pro

Name
Co (Me) 2

B10: IR spectrum for 9j

Agilent Resolutions Pro

Name
Co (CF ₃) ₂

B11: IR spectrum for 9k

B12: IR spectrum for **9l**

B13: IR spectrum for **10a**

Agilent Resolutions Pro

Name
Co (L9) 2

B14: IR spectrum for **10b**

Agilent Resolutions Pro

Name
Co (L10) 2

B15: IR spectrum for 10c

Agilent Resolutions Pro

Name
Co (ox) 3 - L1

B16: IR spectrum for **11a**

Agilent Resolutions Pro

Name
Co (L10) 3

B17: IR spectrum for **11c**

Agilent Resolutions Pro

Name
Cu (ox) 2 - L1

B18: IR spectrum for **8a**

Agilent Resolutions Pro

Name
Cu(oxme) ₂ - L2

B19: IR spectrum for **7a (brown)**

Agilent Resolutions Pro

Name
Cu (oxme) 2 - L2

B20: IR spectrum for **7a (green)**

Agilent Resolutions Pro

Name
Cu(Cl ₂) ₂ - L3

B21: IR spectrum for **7b**

Agilent Resolutions Pro

Name
Cu(OMe) ₂ - L4

B22: IR spectrum for **7c**

Agilent Resolutions Pro

Name
Cu (Me ₂) ₂ - L5

B23: IR spectrum for **7d**

Agilent Resolutions Pro

Name
$\text{Cu}(\text{pCl}_3\text{Me})_2 - \text{L6}$

B24: IR spectrum for **7e**

B25: IR spectrum for **7f**

Agilent Resolutions Pro

Name
Cu(nap)2 - L8

B26: IR spectrum for 7g

Agilent Resolutions Pro

Name
Cu (L9) 2

B27: IR spectrum for **8b**

Agilent Resolutions Pro

Name
Cu (L12) 2

B28: IR spectrum for **7h**

Agilent Resolutions Pro

Name
Cu (L13) 2

B29: IR spectrum for **7i**

Agilent Resolutions Pro

Name
Kathy - IrCl (CO) (PPh ₃) ₂

B30: IR spectrum for **12**

Agilent Resolutions Pro

Name
(COD) Ir (L2)

B31: IR spectrum for 15

Agilent Resolutions Pro

Name
(COD) Ir (L2) MeI

B32: IR spectrum for 16

Agilent Resolutions Pro

Name
(COD) Ir (L2) HSnPh3

B33: IR spectrum for **18**

Appendix C: Ultra-Violet-Visible Spectroscopy (UV-Vis)

C1: UV-Vis Trace of 10a	299
C2: UV-Vis Trace of 11a	302
C3: UV-Vis Trace of 9a	287
C4: UV-Vis Trace of 9b	288
C5: UV-Vis Trace of 9c	289
C6: UV-Vis Trace of 9d	290
C7: UV-Vis Trace of 9e	291
C8: UV-Vis Trace of 9f	292
C9: UV-Vis Trace of 9g	293
C10: UV-Vis Trace of 10b	300
C11: UV-Vis Trace of 10c	301
C12: UV-Vis Trace of 11c	303
C13: UV-Vis Trace of 9h	294
C14: UV-Vis Trace of 9i	295
C15: UV-Vis Trace of 9j	296
C16: UV-Vis Trace of 9k	297
C17: UV-Vis Trace of 9l	298
C18: UV-Vis Trace of 7a (brown)	304
C19: UV-Vis Trace of 7a (green)	305
C20: UV-Vis Trace of 7b	306
C21: UV-Vis Trace of 7c	307
C22: UV-Vis Trace of 7d	308

C23: UV-Vis Trace of 7e	309
C24: UV-Vis Trace of 7f	310
C25: UV-Vis Trace of 7g	311
C26: UV-Vis Trace for 8b	312
C27: UV-Vis Trace of 7h	313
C28: UV-Vis Trace of 7i	314

C1: UV-Vis Trace of **9a**

C3: UV-Vis Trace of **9c**

C4: UV-Vis Trace of **9d**

C7: UV-Vis Trace of **9g**

C8: UV-Vis Trace of **9h**

C9: UV-Vis Trace of **9i**

C10: UV-Vis Trace of **9j**

C11: UV-Vis Trace of **9k**

C13: UV-Vis Trace of **10a**

C14: UV-Vis Trace of **10b**

C15: UV-Vis Trace of **10c**

C16: UV-Vis Trace of **11a**

C17: UV-Vis Trace of **11c**

C20: UV-Vis Trace of **7b**

C21: UV-Vis Trace of **7c**

C22: UV-Vis Trace of **7d**

C23: UV-Vis Trace of **7e**

C24: UV-Vis Trace of **7f**

C27: UV-Vis Trace of **7h**

C28: UV-Vis Trace of **7i**

Appendix D: Cyclic Voltammetry (CV)

D1: Voltammogram of 10a	329
D2: Voltammogram of 11a	331
D3: Voltammogram of 9a	317
D4: Voltammogram of 9b	318
D5: Voltammogram of 9c	319
D6: Voltammogram of 9d	320
D7: Voltammogram of 9e	321
D8: Voltammogram of 9f	322
D9: Voltammogram of 9g	323
D10: Voltammogram of 10c	330
D11: Voltammogram of 11c	332
D12: Voltammogram of 9h	324
D13: Voltammogram of 9i	325
D14: Voltammogram of 9j	326
D15: Voltammogram of 9k	327
D16: Voltammogram of 9l	328
D17: Voltammogram of 7a (brown)	333
D18: Voltammogram of 7a (green)	334
D19: Voltammogram of 7b	335
D20: Voltammogram of 7c	336
D21: Voltammogram of 7d	337
D22: Voltammogram of 7e	338

D23: Voltammogram of 7f	339
D24: Voltammogram of 7g	340
D25: Voltammogram of 7h	341
D26: Voltammogram of 7i	342

D1: Voltammogram of **9a**

D2: Voltammogram of **9b**

D3: Voltammogram of **9c**

D4: Voltammogram of **9d**

D5: Voltammogram of **9e**

D6: Voltammogram of **9f**

D7: Voltammogram of **9g**

D8: Voltammogram of **9h**

D9: Voltammogram of **9i**

D10: Voltammogram of **9j**

D11: Voltammogram of **9k**

D12: Voltammogram of 9I

D13: Voltammogram of **10a**

D14: Voltammogram of **10c**

D15: Voltammogram of **11a**

D16: Voltammogram of **11c**

D17: Voltammogram of **7a** (brown)

D18: Voltammogram of **7a** (green)

D19: Voltammogram of **7b**

D20: Voltammogram of **7c**

D21: Voltammogram of **7d**

D22: Voltammogram of **7e**

D23: Voltammogram of **7f**

D24: Voltammogram of **7g**

D25: Voltammogram of **7h**

D26: Voltammogram of **7i**

Appendix E: Gel-Permeation Chromatography (GPC)

E1: GPC Trace of PMMA (approximate $R_t = 15.5$ mins)	344

E1: GPC Trace of **PMMA** (approximate Rt = 15.5 mins)

Appendix F: Single Crystal X-Ray Diffraction (X-Ray Crystallography)

F1: Crystal data and structure refinement for d1759_a (10a).	346
F2: Crystal data and structure refinement for d1747_a (11a).	358
F3: Crystal data and structure refinement for d1842_a (11c).	375
F4: Crystal data and structure refinement for d1870ra_b (9j).	395
F5: Crystal data and structure refinement for d1867_a (9k).	407
F6: Crystal data and structure refinement for d1889a_a (9l).	419
F7: Crystal data and structure refinement for d1825_a (7g).	429
F8: Crystal data and structure refinement for d1871_a (7h).	440
F9: Crystal data and structure refinement for d1804_a (15).	467
F10: Crystal data and structure refinement for d1872_a (16).	479
F11: Crystal data and structure refinement for 18	492

F1: Crystal data and structure refinement for d1759_a (**10a**).

Identification code	d1759_a (10a₂)	
Empirical formula	C ₂₂ H ₂₀ Co N ₂ O ₄	
Formula weight	435.33	
Temperature	150(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P2 ₁ /n	
Unit cell dimensions	a = 12.3954(7) Å	a = 90°.
	b = 10.2361(5) Å	b = 103.402(2)°.
	c = 15.3565(8) Å	g = 90°.
Volume	1895.38(17) Å ³	
Z	4	
Density (calculated)	1.526 Mg/m ³	
Absorption coefficient	0.938 mm ⁻¹	
F(000)	900	
Crystal size	0.300 x 0.250 x 0.040 mm ³	
Theta range for data collection	1.909 to 27.543°.	
Index ranges	-16 ≤ h ≤ 16, -13 ≤ k ≤ 13, -19 ≤ l ≤ 19	
Reflections collected	27284	

Independent reflections	4360 [R(int) = 0.0392]
Completeness to theta = 25.242°	100.0 %
Absorption correction	Semi-empirical from equivalents
Max. and min. transmission	0.7456 and 0.6452
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	4360 / 0 / 262
Goodness-of-fit on F ²	1.041
Final R indices [I>2sigma(I)]	R1 = 0.0290, wR2 = 0.0694
R indices (all data)	R1 = 0.0430, wR2 = 0.0786
Extinction coefficient	n/a
Largest diff. peak and hole	0.348 and -0.315 e.Å ⁻³

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d1759_a. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	$U(\text{eq})$
Co(1)	7235(1)	7531(1)	7396(1)	22(1)
O(1)	6093(1)	8841(1)	7040(1)	27(1)
O(2)	5406(1)	6725(1)	9239(1)	28(1)
O(3)	7133(1)	6296(1)	6425(1)	25(1)
O(4)	10290(1)	8143(1)	6866(1)	26(1)
N(1)	6688(1)	6895(1)	8419(1)	22(1)
N(2)	8760(1)	8058(1)	7415(1)	21(1)
C(1)	7151(2)	5918(2)	9113(1)	28(1)
C(2)	6216(2)	5717(2)	9603(1)	30(1)
C(3)	5746(1)	7313(2)	8555(1)	21(1)
C(4)	5025(1)	8280(2)	8080(1)	21(1)
C(5)	5225(1)	8977(2)	7363(1)	20(1)
C(6)	4393(1)	9946(2)	6876(1)	21(1)
C(7)	4569(2)	10534(2)	6102(1)	29(1)
C(8)	3794(2)	11394(2)	5609(1)	36(1)
C(9)	2838(2)	11692(2)	5889(1)	34(1)
C(10)	2659(2)	11121(2)	6659(1)	35(1)
C(11)	3423(2)	10256(2)	7148(1)	29(1)
C(12)	9513(1)	8955(2)	8020(1)	25(1)
C(13)	10480(1)	9127(2)	7566(1)	27(1)
C(14)	9266(1)	7623(2)	6821(1)	19(1)
C(15)	8896(1)	6698(2)	6130(1)	21(1)
C(16)	7885(1)	6072(2)	5984(1)	18(1)
C(17)	7572(1)	5073(2)	5256(1)	18(1)
C(18)	6470(1)	4665(2)	4998(1)	23(1)
C(19)	6141(1)	3721(2)	4339(1)	26(1)
C(20)	6912(2)	3181(2)	3923(1)	26(1)
C(21)	8011(2)	3570(2)	4172(1)	29(1)
C(22)	8337(1)	4508(2)	4835(1)	26(1)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for d1759_a.

Co(1)-O(1)	1.9348(12)
Co(1)-O(3)	1.9360(12)
Co(1)-N(2)	1.9588(14)
Co(1)-N(1)	1.9621(14)
O(1)-C(5)	1.2923(19)
O(2)-C(3)	1.359(2)
O(2)-C(2)	1.458(2)
O(3)-C(16)	1.2926(19)
O(4)-C(14)	1.3632(19)
O(4)-C(13)	1.452(2)
N(1)-C(3)	1.305(2)
N(1)-C(1)	1.476(2)
N(2)-C(14)	1.301(2)
N(2)-C(12)	1.474(2)
C(1)-C(2)	1.535(3)
C(1)-H(1A)	0.9900
C(1)-H(1B)	0.9900
C(2)-H(2A)	0.9900
C(2)-H(2B)	0.9900
C(3)-C(4)	1.417(2)
C(4)-C(5)	1.382(2)
C(4)-H(4A)	0.9500
C(5)-C(6)	1.499(2)
C(6)-C(7)	1.395(2)
C(6)-C(11)	1.397(2)
C(7)-C(8)	1.391(3)
C(7)-H(7A)	0.9500
C(8)-C(9)	1.385(3)
C(8)-H(8A)	0.9500
C(9)-C(10)	1.383(3)
C(9)-H(9A)	0.9500
C(10)-C(11)	1.384(3)
C(10)-H(10A)	0.9500
C(11)-H(11A)	0.9500

C(12)-C(13)	1.529(2)
C(12)-H(12A)	0.9900
C(12)-H(12B)	0.9900
C(13)-H(13A)	0.9900
C(13)-H(13B)	0.9900
C(14)-C(15)	1.417(2)
C(15)-C(16)	1.378(2)
C(15)-H(15A)	0.9500
C(16)-C(17)	1.499(2)
C(17)-C(22)	1.391(2)
C(17)-C(18)	1.396(2)
C(18)-C(19)	1.391(2)
C(18)-H(18A)	0.9500
C(19)-C(20)	1.382(2)
C(19)-H(19A)	0.9500
C(20)-C(21)	1.385(3)
C(20)-H(20A)	0.9500
C(21)-C(22)	1.390(2)
C(21)-H(21A)	0.9500
C(22)-H(22A)	0.9500
O(1)-Co(1)-O(3)	108.80(5)
O(1)-Co(1)-N(2)	116.97(6)
O(3)-Co(1)-N(2)	94.75(5)
O(1)-Co(1)-N(1)	95.14(6)
O(3)-Co(1)-N(1)	115.29(6)
N(2)-Co(1)-N(1)	126.20(6)
C(5)-O(1)-Co(1)	125.93(11)
C(3)-O(2)-C(2)	107.50(13)
C(16)-O(3)-Co(1)	126.28(11)
C(14)-O(4)-C(13)	107.31(13)
C(3)-N(1)-C(1)	108.65(14)
C(3)-N(1)-Co(1)	120.48(12)
C(1)-N(1)-Co(1)	130.84(11)
C(14)-N(2)-C(12)	108.45(14)
C(14)-N(2)-Co(1)	121.03(11)

C(12)-N(2)-Co(1)	130.50(11)
N(1)-C(1)-C(2)	103.48(14)
N(1)-C(1)-H(1A)	111.1
C(2)-C(1)-H(1A)	111.1
N(1)-C(1)-H(1B)	111.1
C(2)-C(1)-H(1B)	111.1
H(1A)-C(1)-H(1B)	109.0
O(2)-C(2)-C(1)	104.36(14)
O(2)-C(2)-H(2A)	110.9
C(1)-C(2)-H(2A)	110.9
O(2)-C(2)-H(2B)	110.9
C(1)-C(2)-H(2B)	110.9
H(2A)-C(2)-H(2B)	108.9
N(1)-C(3)-O(2)	115.26(15)
N(1)-C(3)-C(4)	128.92(16)
O(2)-C(3)-C(4)	115.82(15)
C(5)-C(4)-C(3)	124.23(15)
C(5)-C(4)-H(4A)	117.9
C(3)-C(4)-H(4A)	117.9
O(1)-C(5)-C(4)	124.66(16)
O(1)-C(5)-C(6)	114.71(14)
C(4)-C(5)-C(6)	120.61(15)
C(7)-C(6)-C(11)	118.19(16)
C(7)-C(6)-C(5)	119.02(16)
C(11)-C(6)-C(5)	122.75(16)
C(8)-C(7)-C(6)	120.83(18)
C(8)-C(7)-H(7A)	119.6
C(6)-C(7)-H(7A)	119.6
C(9)-C(8)-C(7)	120.21(19)
C(9)-C(8)-H(8A)	119.9
C(7)-C(8)-H(8A)	119.9
C(10)-C(9)-C(8)	119.42(18)
C(10)-C(9)-H(9A)	120.3
C(8)-C(9)-H(9A)	120.3
C(9)-C(10)-C(11)	120.61(19)
C(9)-C(10)-H(10A)	119.7

C(11)-C(10)-H(10A)	119.7
C(10)-C(11)-C(6)	120.73(18)
C(10)-C(11)-H(11A)	119.6
C(6)-C(11)-H(11A)	119.6
N(2)-C(12)-C(13)	103.51(13)
N(2)-C(12)-H(12A)	111.1
C(13)-C(12)-H(12A)	111.1
N(2)-C(12)-H(12B)	111.1
C(13)-C(12)-H(12B)	111.1
H(12A)-C(12)-H(12B)	109.0
O(4)-C(13)-C(12)	104.36(13)
O(4)-C(13)-H(13A)	110.9
C(12)-C(13)-H(13A)	110.9
O(4)-C(13)-H(13B)	110.9
C(12)-C(13)-H(13B)	110.9
H(13A)-C(13)-H(13B)	108.9
N(2)-C(14)-O(4)	115.19(14)
N(2)-C(14)-C(15)	128.96(15)
O(4)-C(14)-C(15)	115.84(14)
C(16)-C(15)-C(14)	123.83(15)
C(16)-C(15)-H(15A)	118.1
C(14)-C(15)-H(15A)	118.1
O(3)-C(16)-C(15)	124.87(15)
O(3)-C(16)-C(17)	114.71(14)
C(15)-C(16)-C(17)	120.40(14)
C(22)-C(17)-C(18)	118.12(15)
C(22)-C(17)-C(16)	123.01(15)
C(18)-C(17)-C(16)	118.85(14)
C(19)-C(18)-C(17)	121.11(16)
C(19)-C(18)-H(18A)	119.4
C(17)-C(18)-H(18A)	119.4
C(20)-C(19)-C(18)	119.79(16)
C(20)-C(19)-H(19A)	120.1
C(18)-C(19)-H(19A)	120.1
C(19)-C(20)-C(21)	119.95(16)
C(19)-C(20)-H(20A)	120.0

C(21)-C(20)-H(20A)	120.0
C(20)-C(21)-C(22)	120.04(17)
C(20)-C(21)-H(21A)	120.0
C(22)-C(21)-H(21A)	120.0
C(21)-C(22)-C(17)	120.97(16)
C(21)-C(22)-H(22A)	119.5
C(17)-C(22)-H(22A)	119.5

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d1759_a. The anisotropic displacement factor exponent takes the form: $-2p^2[h^2 a^{*2}U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Co(1)	20(1)	26(1)	23(1)	-4(1)	10(1)	1(1)
O(1)	26(1)	30(1)	30(1)	5(1)	14(1)	4(1)
O(2)	26(1)	33(1)	26(1)	8(1)	11(1)	4(1)
O(3)	20(1)	31(1)	27(1)	-8(1)	12(1)	-4(1)
O(4)	19(1)	27(1)	35(1)	-12(1)	10(1)	-4(1)
N(1)	22(1)	23(1)	22(1)	-2(1)	6(1)	1(1)
N(2)	21(1)	20(1)	22(1)	-3(1)	5(1)	1(1)
C(1)	27(1)	28(1)	27(1)	1(1)	6(1)	6(1)
C(2)	32(1)	28(1)	29(1)	5(1)	8(1)	5(1)
C(3)	21(1)	24(1)	18(1)	-3(1)	5(1)	-4(1)
C(4)	17(1)	25(1)	22(1)	-2(1)	6(1)	0(1)
C(5)	18(1)	19(1)	22(1)	-5(1)	5(1)	-3(1)
C(6)	22(1)	18(1)	22(1)	-4(1)	3(1)	-3(1)
C(7)	30(1)	30(1)	29(1)	2(1)	8(1)	-1(1)
C(8)	43(1)	33(1)	30(1)	8(1)	4(1)	-2(1)
C(9)	31(1)	25(1)	40(1)	4(1)	-4(1)	1(1)
C(10)	28(1)	30(1)	47(1)	4(1)	9(1)	6(1)
C(11)	28(1)	27(1)	34(1)	5(1)	10(1)	3(1)
C(12)	23(1)	25(1)	26(1)	-8(1)	6(1)	0(1)
C(13)	23(1)	24(1)	32(1)	-11(1)	5(1)	-2(1)
C(14)	15(1)	19(1)	23(1)	1(1)	4(1)	2(1)
C(15)	18(1)	22(1)	22(1)	-4(1)	8(1)	1(1)
C(16)	18(1)	19(1)	18(1)	1(1)	6(1)	3(1)
C(17)	19(1)	18(1)	18(1)	2(1)	4(1)	1(1)
C(18)	19(1)	27(1)	23(1)	0(1)	6(1)	0(1)
C(19)	20(1)	29(1)	26(1)	0(1)	2(1)	-3(1)
C(20)	30(1)	22(1)	24(1)	-6(1)	2(1)	-2(1)
C(21)	26(1)	31(1)	32(1)	-10(1)	11(1)	1(1)
C(22)	17(1)	31(1)	31(1)	-9(1)	8(1)	-2(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for d1759_a.

	x	y	z	U(eq)
H(1A)	7830	6252	9526	33
H(1B)	7328	5093	8840	33
H(2A)	5886	4835	9482	36
H(2B)	6494	5826	10257	36
H(4A)	4361	8461	8267	25
H(7A)	5227	10346	5909	35
H(8A)	3920	11777	5077	43
H(9A)	2311	12285	5555	41
H(10A)	2005	11323	6854	42
H(11A)	3287	9868	7675	35
H(12A)	9768	8569	8624	30
H(12B)	9146	9801	8072	30
H(13A)	10476	10015	7308	32
H(13B)	11198	8982	7996	32
H(15A)	9372	6496	5745	25
H(18A)	5935	5039	5278	27
H(19A)	5388	3448	4175	31
H(20A)	6689	2543	3468	31
H(21A)	8542	3195	3889	35
H(22A)	9092	4768	5003	31

Table 6. Torsion angles [°] for d1759_a.

C(3)-N(1)-C(1)-C(2)	7.36(19)
Co(1)-N(1)-C(1)-C(2)	-170.37(12)
C(3)-O(2)-C(2)-C(1)	6.99(18)
N(1)-C(1)-C(2)-O(2)	-8.50(18)
C(1)-N(1)-C(3)-O(2)	-3.3(2)
Co(1)-N(1)-C(3)-O(2)	174.70(11)
C(1)-N(1)-C(3)-C(4)	177.69(17)
Co(1)-N(1)-C(3)-C(4)	-4.3(2)
C(2)-O(2)-C(3)-N(1)	-2.61(19)
C(2)-O(2)-C(3)-C(4)	176.52(14)
N(1)-C(3)-C(4)-C(5)	-0.6(3)
O(2)-C(3)-C(4)-C(5)	-179.59(15)
Co(1)-O(1)-C(5)-C(4)	6.6(2)
Co(1)-O(1)-C(5)-C(6)	-171.60(10)
C(3)-C(4)-C(5)-O(1)	-0.4(3)
C(3)-C(4)-C(5)-C(6)	177.60(15)
O(1)-C(5)-C(6)-C(7)	4.6(2)
C(4)-C(5)-C(6)-C(7)	-173.60(16)
O(1)-C(5)-C(6)-C(11)	-177.68(16)
C(4)-C(5)-C(6)-C(11)	4.1(2)
C(11)-C(6)-C(7)-C(8)	-0.7(3)
C(5)-C(6)-C(7)-C(8)	177.06(16)
C(6)-C(7)-C(8)-C(9)	0.9(3)
C(7)-C(8)-C(9)-C(10)	-0.6(3)
C(8)-C(9)-C(10)-C(11)	0.0(3)
C(9)-C(10)-C(11)-C(6)	0.2(3)
C(7)-C(6)-C(11)-C(10)	0.2(3)
C(5)-C(6)-C(11)-C(10)	-177.54(17)
C(14)-N(2)-C(12)-C(13)	9.02(18)
Co(1)-N(2)-C(12)-C(13)	-169.80(12)
C(14)-O(4)-C(13)-C(12)	8.85(18)
N(2)-C(12)-C(13)-O(4)	-10.62(18)
C(12)-N(2)-C(14)-O(4)	-3.8(2)
Co(1)-N(2)-C(14)-O(4)	175.11(11)

C(12)-N(2)-C(14)-C(15)	176.83(16)
Co(1)-N(2)-C(14)-C(15)	-4.2(2)
C(13)-O(4)-C(14)-N(2)	-3.51(19)
C(13)-O(4)-C(14)-C(15)	175.90(14)
N(2)-C(14)-C(15)-C(16)	-0.8(3)
O(4)-C(14)-C(15)-C(16)	179.93(15)
Co(1)-O(3)-C(16)-C(15)	-0.7(2)
Co(1)-O(3)-C(16)-C(17)	-179.18(10)
C(14)-C(15)-C(16)-O(3)	3.6(3)
C(14)-C(15)-C(16)-C(17)	-178.03(15)
O(3)-C(16)-C(17)-C(22)	-166.48(16)
C(15)-C(16)-C(17)-C(22)	15.0(2)
O(3)-C(16)-C(17)-C(18)	12.1(2)
C(15)-C(16)-C(17)-C(18)	-166.41(16)
C(22)-C(17)-C(18)-C(19)	0.0(3)
C(16)-C(17)-C(18)-C(19)	-178.69(15)
C(17)-C(18)-C(19)-C(20)	-0.6(3)
C(18)-C(19)-C(20)-C(21)	0.7(3)
C(19)-C(20)-C(21)-C(22)	-0.4(3)
C(20)-C(21)-C(22)-C(17)	-0.2(3)
C(18)-C(17)-C(22)-C(21)	0.4(3)
C(16)-C(17)-C(22)-C(21)	179.02(16)

Symmetry transformations used to generate equivalent atoms:

F2: Crystal data and structure refinement for d1747_a (**11a**).

Identification code	d1747_a (11a)	
Empirical formula	C ₃₃ H ₃₀ Co N ₃ O ₆	
Formula weight	623.53	
Temperature	150(2) K	
Wavelength	1.54178 Å	
Crystal system	Triclinic	
Space group	P-1	
Unit cell dimensions	$a = 10.7799(4) \text{ Å}$	$\alpha = 72.051(2)^\circ$
	$b = 10.7875(4) \text{ Å}$	$\beta = 84.462(2)^\circ$
	$c = 13.4524(5) \text{ Å}$	$\gamma = 68.638(2)^\circ$
Volume	$1385.76(9) \text{ Å}^3$	
Z	2	
Density (calculated)	1.494 Mg/m^3	
Absorption coefficient	5.301 mm^{-1}	
F(000)	648	
Crystal size	$0.160 \times 0.050 \times 0.040 \text{ mm}^3$	
Theta range for data collection	3.454 to 67.549°	
Index ranges	$-12 \leq h \leq 12$, $-12 \leq k \leq 12$, $-16 \leq l \leq 16$	
Reflections collected	34303	
Independent reflections	4927 [R(int) = 0.0638]	
Completeness to theta = 67.549°	98.6 %	
Absorption correction	Semi-empirical from equivalents	

Max. and min. transmission	0.7529 and 0.5658
Refinement method	Full-matrix least-squares on F^2
Data / restraints / parameters	4927 / 0 / 388
Goodness-of-fit on F^2	1.022
Final R indices [$I > 2\sigma(I)$]	$R_1 = 0.0401$, $wR_2 = 0.0984$
R indices (all data)	$R_1 = 0.0479$, $wR_2 = 0.1032$
Extinction coefficient	n/a
Largest diff. peak and hole	0.506 and -0.411 $e.\text{\AA}^{-3}$

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\approx^2 \times 10^3$) for d1747_a. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	U(eq)
Co(1)	5074(1)	7355(1)	7297(1)	22(1)
O(1)	3619(2)	6986(2)	8080(1)	27(1)
O(2)	2799(2)	7544(2)	4971(1)	40(1)
O(3)	5932(2)	5406(2)	7486(1)	26(1)
O(4)	8708(2)	7371(2)	6154(1)	35(1)
O(5)	4095(2)	9288(2)	7038(1)	26(1)
O(6)	5751(2)	7681(2)	10102(1)	34(1)
N(1)	4335(2)	7463(2)	6015(2)	24(1)
N(2)	6574(2)	7777(2)	6608(2)	26(1)
N(3)	5751(2)	7134(2)	8629(2)	26(1)
C(1)	2440(2)	7329(2)	7708(2)	26(1)
C(2)	2113(2)	7669(3)	6671(2)	29(1)
C(3)	3119(2)	7570(2)	5914(2)	26(1)
C(4)	4024(3)	7217(3)	4395(2)	33(1)
C(5)	5078(2)	7210(3)	5082(2)	32(1)
C(6)	1404(2)	7286(3)	8528(2)	29(1)
C(7)	443(3)	6732(3)	8483(2)	39(1)
C(8)	-481(3)	6669(3)	9273(3)	50(1)
C(9)	-462(3)	7182(4)	10097(2)	51(1)
C(10)	483(3)	7757(3)	10135(2)	45(1)
C(11)	1427(3)	7781(3)	9360(2)	36(1)
C(12)	7164(2)	4773(3)	7303(2)	25(1)
C(13)	8088(2)	5396(3)	6911(2)	30(1)
C(14)	7746(2)	6844(3)	6579(2)	27(1)
C(15)	8120(3)	8866(3)	5924(2)	37(1)
C(16)	6657(3)	9163(3)	6157(2)	40(1)
C(17)	7567(2)	3237(3)	7570(2)	26(1)
C(18)	8639(3)	2445(3)	7087(2)	33(1)
C(19)	9008(3)	1024(3)	7359(2)	38(1)
C(20)	8316(3)	347(3)	8108(2)	39(1)
C(21)	7245(3)	1112(3)	8586(2)	35(1)

C(22)	6876(2)	2542(3)	8319(2)	29(1)
C(23)	4012(2)	10027(3)	7666(2)	26(1)
C(24)	4583(2)	9519(3)	8649(2)	27(1)
C(25)	5359(2)	8110(3)	9083(2)	26(1)
C(26)	6466(3)	6182(3)	10386(2)	32(1)
C(27)	6416(3)	5784(3)	9398(2)	33(1)
C(28)	3254(2)	11538(3)	7217(2)	26(1)
C(29)	3593(3)	12550(3)	7461(2)	33(1)
C(30)	2956(3)	13948(3)	6989(2)	40(1)
C(31)	1934(3)	14381(3)	6263(2)	39(1)
C(32)	1583(2)	13406(3)	6008(2)	35(1)
C(33)	2238(2)	11998(3)	6481(2)	33(1)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for d1747_a.

Co(1)-O(5)	1.8983(16)
Co(1)-O(1)	1.8993(16)
Co(1)-O(3)	1.9072(16)
Co(1)-N(3)	1.9144(19)
Co(1)-N(2)	1.915(2)
Co(1)-N(1)	1.9219(19)
O(1)-C(1)	1.292(3)
O(2)-C(3)	1.356(3)
O(2)-C(4)	1.449(3)
O(3)-C(12)	1.292(3)
O(4)-C(14)	1.357(3)
O(4)-C(15)	1.444(3)
O(5)-C(23)	1.307(3)
O(6)-C(25)	1.360(3)
O(6)-C(26)	1.457(3)
N(1)-C(3)	1.291(3)
N(1)-C(5)	1.467(3)
N(2)-C(14)	1.304(3)
N(2)-C(16)	1.463(3)
N(3)-C(25)	1.296(3)
N(3)-C(27)	1.470(3)
C(1)-C(2)	1.375(3)
C(1)-C(6)	1.494(3)
C(2)-C(3)	1.413(3)
C(2)-H(2A)	0.9500
C(4)-C(5)	1.528(3)
C(4)-H(4A)	0.9900
C(4)-H(4B)	0.9900
C(5)-H(5A)	0.9900
C(5)-H(5B)	0.9900
C(6)-C(7)	1.385(4)
C(6)-C(11)	1.387(4)
C(7)-C(8)	1.387(4)
C(7)-H(7A)	0.9500

C(8)-C(9)	1.388(4)
C(8)-H(8A)	0.9500
C(9)-C(10)	1.384(4)
C(9)-H(9A)	0.9500
C(10)-C(11)	1.386(4)
C(10)-H(10A)	0.9500
C(11)-H(11A)	0.9500
C(12)-C(13)	1.374(3)
C(12)-C(17)	1.484(3)
C(13)-C(14)	1.399(4)
C(13)-H(13A)	0.9500
C(15)-C(16)	1.513(4)
C(15)-H(15A)	0.9900
C(15)-H(15B)	0.9900
C(16)-H(16A)	0.9900
C(16)-H(16B)	0.9900
C(17)-C(22)	1.389(3)
C(17)-C(18)	1.402(3)
C(18)-C(19)	1.372(4)
C(18)-H(18A)	0.9500
C(19)-C(20)	1.380(4)
C(19)-H(19A)	0.9500
C(20)-C(21)	1.384(4)
C(20)-H(20A)	0.9500
C(21)-C(22)	1.380(4)
C(21)-H(21A)	0.9500
C(22)-H(22A)	0.9500
C(23)-C(24)	1.375(3)
C(23)-C(28)	1.485(3)
C(24)-C(25)	1.405(4)
C(24)-H(24A)	0.9500
C(26)-C(27)	1.528(3)
C(26)-H(26A)	0.9900
C(26)-H(26B)	0.9900
C(27)-H(27A)	0.9900
C(27)-H(27B)	0.9900

C(28)-C(33)	1.392(3)
C(28)-C(29)	1.403(3)
C(29)-C(30)	1.374(4)
C(29)-H(29A)	0.9500
C(30)-C(31)	1.392(4)
C(30)-H(30A)	0.9500
C(31)-C(32)	1.380(4)
C(31)-H(31A)	0.9500
C(32)-C(33)	1.386(4)
C(32)-H(32A)	0.9500
C(33)-H(33A)	0.9500
O(5)-Co(1)-O(1)	88.61(7)
O(5)-Co(1)-O(3)	174.68(7)
O(1)-Co(1)-O(3)	89.04(7)
O(5)-Co(1)-N(3)	94.33(7)
O(1)-Co(1)-N(3)	84.09(8)
O(3)-Co(1)-N(3)	90.18(8)
O(5)-Co(1)-N(2)	89.78(8)
O(1)-Co(1)-N(2)	174.84(7)
O(3)-Co(1)-N(2)	92.96(8)
N(3)-Co(1)-N(2)	91.15(8)
O(5)-Co(1)-N(1)	88.15(7)
O(1)-Co(1)-N(1)	92.11(7)
O(3)-Co(1)-N(1)	87.16(7)
N(3)-Co(1)-N(1)	175.40(8)
N(2)-Co(1)-N(1)	92.73(8)
C(1)-O(1)-Co(1)	125.40(15)
C(3)-O(2)-C(4)	107.09(18)
C(12)-O(3)-Co(1)	126.63(15)
C(14)-O(4)-C(15)	107.23(19)
C(23)-O(5)-Co(1)	126.15(15)
C(25)-O(6)-C(26)	107.56(18)
C(3)-N(1)-C(5)	109.44(19)
C(3)-N(1)-Co(1)	123.46(16)
C(5)-N(1)-Co(1)	126.38(15)

C(14)-N(2)-C(16)	108.9(2)
C(14)-N(2)-Co(1)	124.12(17)
C(16)-N(2)-Co(1)	126.54(16)
C(25)-N(3)-C(27)	109.74(19)
C(25)-N(3)-Co(1)	123.18(17)
C(27)-N(3)-Co(1)	124.44(15)
O(1)-C(1)-C(2)	125.7(2)
O(1)-C(1)-C(6)	113.6(2)
C(2)-C(1)-C(6)	120.6(2)
C(1)-C(2)-C(3)	120.2(2)
C(1)-C(2)-H(2A)	119.9
C(3)-C(2)-H(2A)	119.9
N(1)-C(3)-O(2)	115.1(2)
N(1)-C(3)-C(2)	127.7(2)
O(2)-C(3)-C(2)	117.2(2)
O(2)-C(4)-C(5)	104.92(18)
O(2)-C(4)-H(4A)	110.8
C(5)-C(4)-H(4A)	110.8
O(2)-C(4)-H(4B)	110.8
C(5)-C(4)-H(4B)	110.8
H(4A)-C(4)-H(4B)	108.8
N(1)-C(5)-C(4)	103.03(19)
N(1)-C(5)-H(5A)	111.2
C(4)-C(5)-H(5A)	111.2
N(1)-C(5)-H(5B)	111.2
C(4)-C(5)-H(5B)	111.2
H(5A)-C(5)-H(5B)	109.1
C(7)-C(6)-C(11)	119.5(2)
C(7)-C(6)-C(1)	121.3(2)
C(11)-C(6)-C(1)	119.2(2)
C(6)-C(7)-C(8)	119.8(3)
C(6)-C(7)-H(7A)	120.1
C(8)-C(7)-H(7A)	120.1
C(7)-C(8)-C(9)	120.3(3)
C(7)-C(8)-H(8A)	119.8
C(9)-C(8)-H(8A)	119.8

C(10)-C(9)-C(8)	119.9(3)
C(10)-C(9)-H(9A)	120.0
C(8)-C(9)-H(9A)	120.0
C(9)-C(10)-C(11)	119.5(3)
C(9)-C(10)-H(10A)	120.3
C(11)-C(10)-H(10A)	120.3
C(10)-C(11)-C(6)	120.8(3)
C(10)-C(11)-H(11A)	119.6
C(6)-C(11)-H(11A)	119.6
O(3)-C(12)-C(13)	125.8(2)
O(3)-C(12)-C(17)	114.5(2)
C(13)-C(12)-C(17)	119.6(2)
C(12)-C(13)-C(14)	122.2(2)
C(12)-C(13)-H(13A)	118.9
C(14)-C(13)-H(13A)	118.9
N(2)-C(14)-O(4)	114.7(2)
N(2)-C(14)-C(13)	127.1(2)
O(4)-C(14)-C(13)	118.2(2)
O(4)-C(15)-C(16)	105.2(2)
O(4)-C(15)-H(15A)	110.7
C(16)-C(15)-H(15A)	110.7
O(4)-C(15)-H(15B)	110.7
C(16)-C(15)-H(15B)	110.7
H(15A)-C(15)-H(15B)	108.8
N(2)-C(16)-C(15)	103.6(2)
N(2)-C(16)-H(16A)	111.0
C(15)-C(16)-H(16A)	111.0
N(2)-C(16)-H(16B)	111.0
C(15)-C(16)-H(16B)	111.0
H(16A)-C(16)-H(16B)	109.0
C(22)-C(17)-C(18)	118.0(2)
C(22)-C(17)-C(12)	120.0(2)
C(18)-C(17)-C(12)	122.0(2)
C(19)-C(18)-C(17)	121.0(3)
C(19)-C(18)-H(18A)	119.5
C(17)-C(18)-H(18A)	119.5

C(18)-C(19)-C(20)	120.3(2)
C(18)-C(19)-H(19A)	119.9
C(20)-C(19)-H(19A)	119.9
C(19)-C(20)-C(21)	119.6(3)
C(19)-C(20)-H(20A)	120.2
C(21)-C(20)-H(20A)	120.2
C(22)-C(21)-C(20)	120.2(3)
C(22)-C(21)-H(21A)	119.9
C(20)-C(21)-H(21A)	119.9
C(21)-C(22)-C(17)	120.9(2)
C(21)-C(22)-H(22A)	119.6
C(17)-C(22)-H(22A)	119.6
O(5)-C(23)-C(24)	125.4(2)
O(5)-C(23)-C(28)	113.9(2)
C(24)-C(23)-C(28)	120.6(2)
C(23)-C(24)-C(25)	122.5(2)
C(23)-C(24)-H(24A)	118.8
C(25)-C(24)-H(24A)	118.8
N(3)-C(25)-O(6)	114.5(2)
N(3)-C(25)-C(24)	127.7(2)
O(6)-C(25)-C(24)	117.8(2)
O(6)-C(26)-C(27)	104.79(19)
O(6)-C(26)-H(26A)	110.8
C(27)-C(26)-H(26A)	110.8
O(6)-C(26)-H(26B)	110.8
C(27)-C(26)-H(26B)	110.8
H(26A)-C(26)-H(26B)	108.9
N(3)-C(27)-C(26)	103.1(2)
N(3)-C(27)-H(27A)	111.1
C(26)-C(27)-H(27A)	111.1
N(3)-C(27)-H(27B)	111.1
C(26)-C(27)-H(27B)	111.1
H(27A)-C(27)-H(27B)	109.1
C(33)-C(28)-C(29)	117.6(2)
C(33)-C(28)-C(23)	120.7(2)
C(29)-C(28)-C(23)	121.5(2)

C(30)-C(29)-C(28)	121.5(2)
C(30)-C(29)-H(29A)	119.2
C(28)-C(29)-H(29A)	119.2
C(29)-C(30)-C(31)	119.7(3)
C(29)-C(30)-H(30A)	120.2
C(31)-C(30)-H(30A)	120.2
C(32)-C(31)-C(30)	119.9(3)
C(32)-C(31)-H(31A)	120.1
C(30)-C(31)-H(31A)	120.1
C(31)-C(32)-C(33)	120.1(2)
C(31)-C(32)-H(32A)	119.9
C(33)-C(32)-H(32A)	119.9
C(32)-C(33)-C(28)	121.1(2)
C(32)-C(33)-H(33A)	119.5
C(28)-C(33)-H(33A)	119.5

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\approx 10^3$) for d1747_a. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Co(1)	23(1)	20(1)	24(1)	-9(1)	-3(1)	-4(1)
O(1)	23(1)	26(1)	27(1)	-9(1)	-3(1)	-3(1)
O(2)	34(1)	62(1)	30(1)	-20(1)	-3(1)	-17(1)
O(3)	23(1)	25(1)	30(1)	-10(1)	-1(1)	-5(1)
O(4)	30(1)	34(1)	45(1)	-14(1)	5(1)	-14(1)
O(5)	28(1)	23(1)	28(1)	-11(1)	-4(1)	-4(1)
O(6)	46(1)	29(1)	23(1)	-9(1)	-7(1)	-5(1)
N(1)	25(1)	22(1)	25(1)	-9(1)	-1(1)	-5(1)
N(2)	29(1)	22(1)	29(1)	-9(1)	-3(1)	-9(1)
N(3)	26(1)	21(1)	27(1)	-9(1)	-4(1)	-3(1)
C(1)	26(1)	17(1)	32(1)	-9(1)	-1(1)	-2(1)
C(2)	22(1)	29(1)	34(1)	-11(1)	-3(1)	-5(1)
C(3)	29(1)	20(1)	26(1)	-7(1)	-6(1)	-4(1)
C(4)	36(1)	35(2)	31(1)	-16(1)	1(1)	-10(1)
C(5)	31(1)	40(2)	26(1)	-16(1)	1(1)	-7(1)
C(6)	24(1)	24(1)	32(1)	-7(1)	-3(1)	-1(1)
C(7)	38(2)	35(2)	45(2)	-15(1)	4(1)	-12(1)
C(8)	43(2)	48(2)	63(2)	-18(2)	12(2)	-23(2)
C(9)	45(2)	59(2)	44(2)	-11(2)	10(1)	-18(2)
C(10)	36(2)	62(2)	33(1)	-18(1)	-1(1)	-10(1)
C(11)	28(1)	45(2)	30(1)	-10(1)	-3(1)	-7(1)
C(12)	24(1)	25(1)	23(1)	-10(1)	-4(1)	-3(1)
C(13)	23(1)	28(2)	36(1)	-12(1)	1(1)	-5(1)
C(14)	25(1)	31(2)	29(1)	-13(1)	0(1)	-10(1)
C(15)	42(2)	32(2)	44(2)	-16(1)	5(1)	-18(1)
C(16)	39(2)	26(2)	54(2)	-10(1)	2(1)	-12(1)
C(17)	26(1)	23(1)	25(1)	-9(1)	-7(1)	-2(1)
C(18)	33(1)	30(2)	35(1)	-14(1)	2(1)	-6(1)
C(19)	36(1)	31(2)	47(2)	-20(1)	1(1)	-1(1)
C(20)	42(2)	24(2)	46(2)	-11(1)	-7(1)	-4(1)
C(21)	37(1)	29(2)	38(1)	-8(1)	-2(1)	-10(1)

C(22)	28(1)	26(1)	31(1)	-11(1)	-2(1)	-4(1)
C(23)	25(1)	25(1)	29(1)	-12(1)	2(1)	-8(1)
C(24)	34(1)	25(1)	25(1)	-12(1)	1(1)	-10(1)
C(25)	25(1)	30(1)	23(1)	-8(1)	-1(1)	-9(1)
C(26)	33(1)	28(2)	31(1)	-9(1)	-6(1)	-2(1)
C(27)	36(1)	27(2)	28(1)	-7(1)	-6(1)	-3(1)
C(28)	28(1)	25(1)	25(1)	-12(1)	3(1)	-5(1)
C(29)	41(2)	28(2)	29(1)	-11(1)	-2(1)	-7(1)
C(30)	58(2)	25(2)	33(1)	-12(1)	-1(1)	-7(1)
C(31)	44(2)	24(2)	35(1)	-8(1)	0(1)	3(1)
C(32)	27(1)	35(2)	37(1)	-11(1)	-1(1)	-1(1)
C(33)	29(1)	32(2)	34(1)	-12(1)	0(1)	-6(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\approx 2 \times 10^{-3}$) for d1747_a.

	x	y	z	U(eq)
H(2A)	1207	7971	6463	35
H(4A)	3937	7930	3708	40
H(4B)	4258	6295	4281	40
H(5A)	5820	6300	5259	39
H(5B)	5439	7955	4732	39
H(7A)	416	6395	7914	47
H(8A)	-1130	6274	9249	60
H(9A)	-1097	7137	10636	61
H(10A)	485	8133	10687	54
H(11A)	2099	8141	9399	43
H(13A)	8988	4824	6862	36
H(15A)	8244	9329	5182	45
H(15B)	8528	9198	6370	45
H(16A)	6362	9713	6661	48
H(16B)	6105	9677	5511	48
H(18A)	9117	2900	6564	40
H(19A)	9744	502	7029	46
H(20A)	8572	-637	8295	46
H(21A)	6762	652	9100	42
H(22A)	6139	3057	8651	35
H(24A)	4447	10144	9049	32
H(26A)	6028	5689	10971	39
H(26B)	7399	5952	10594	39
H(27A)	7324	5316	9174	39
H(27B)	5897	5163	9510	39
H(29A)	4280	12262	7965	40
H(30A)	3211	14615	7158	48
H(31A)	1479	15346	5943	46
H(32A)	891	13701	5508	42
H(33A)	1989	11335	6299	39

Table 6. Torsion angles [$^{\circ}$] for d1747_a.

O(5)-Co(1)-O(1)-C(1)	63.96(18)
O(3)-Co(1)-O(1)-C(1)	-111.26(18)
N(3)-Co(1)-O(1)-C(1)	158.46(19)
N(1)-Co(1)-O(1)-C(1)	-24.14(19)
O(1)-Co(1)-O(5)-C(23)	82.72(18)
N(3)-Co(1)-O(5)-C(23)	-1.25(19)
N(2)-Co(1)-O(5)-C(23)	-92.38(18)
N(1)-Co(1)-O(5)-C(23)	174.88(18)
Co(1)-O(1)-C(1)-C(2)	18.5(3)
Co(1)-O(1)-C(1)-C(6)	-163.39(15)
O(1)-C(1)-C(2)-C(3)	4.4(4)
C(6)-C(1)-C(2)-C(3)	-173.5(2)
C(5)-N(1)-C(3)-O(2)	-5.2(3)
Co(1)-N(1)-C(3)-O(2)	-176.00(16)
C(5)-N(1)-C(3)-C(2)	172.8(2)
Co(1)-N(1)-C(3)-C(2)	2.1(4)
C(4)-O(2)-C(3)-N(1)	7.3(3)
C(4)-O(2)-C(3)-C(2)	-171.0(2)
C(1)-C(2)-C(3)-N(1)	-15.6(4)
C(1)-C(2)-C(3)-O(2)	162.5(2)
C(3)-O(2)-C(4)-C(5)	-5.9(3)
C(3)-N(1)-C(5)-C(4)	1.0(3)
Co(1)-N(1)-C(5)-C(4)	171.43(17)
O(2)-C(4)-C(5)-N(1)	3.0(3)
O(1)-C(1)-C(6)-C(7)	-137.5(2)
C(2)-C(1)-C(6)-C(7)	40.6(4)
O(1)-C(1)-C(6)-C(11)	41.0(3)
C(2)-C(1)-C(6)-C(11)	-140.8(3)
C(11)-C(6)-C(7)-C(8)	-0.4(4)
C(1)-C(6)-C(7)-C(8)	178.1(3)
C(6)-C(7)-C(8)-C(9)	1.1(5)
C(7)-C(8)-C(9)-C(10)	-0.1(5)
C(8)-C(9)-C(10)-C(11)	-1.8(5)
C(9)-C(10)-C(11)-C(6)	2.5(4)

C(7)-C(6)-C(11)-C(10)	-1.4(4)
C(1)-C(6)-C(11)-C(10)	-180.0(2)
Co(1)-O(3)-C(12)-C(13)	-1.7(3)
Co(1)-O(3)-C(12)-C(17)	177.36(14)
O(3)-C(12)-C(13)-C(14)	-5.4(4)
C(17)-C(12)-C(13)-C(14)	175.6(2)
C(16)-N(2)-C(14)-O(4)	-0.1(3)
Co(1)-N(2)-C(14)-O(4)	-172.97(14)
C(16)-N(2)-C(14)-C(13)	-178.4(2)
Co(1)-N(2)-C(14)-C(13)	8.8(3)
C(15)-O(4)-C(14)-N(2)	4.1(3)
C(15)-O(4)-C(14)-C(13)	-177.5(2)
C(12)-C(13)-C(14)-N(2)	1.5(4)
C(12)-C(13)-C(14)-O(4)	-176.7(2)
C(14)-O(4)-C(15)-C(16)	-6.1(3)
C(14)-N(2)-C(16)-C(15)	-3.6(3)
Co(1)-N(2)-C(16)-C(15)	168.96(16)
O(4)-C(15)-C(16)-N(2)	5.8(3)
O(3)-C(12)-C(17)-C(22)	-24.8(3)
C(13)-C(12)-C(17)-C(22)	154.3(2)
O(3)-C(12)-C(17)-C(18)	155.3(2)
C(13)-C(12)-C(17)-C(18)	-25.6(3)
C(22)-C(17)-C(18)-C(19)	-1.0(4)
C(12)-C(17)-C(18)-C(19)	178.9(2)
C(17)-C(18)-C(19)-C(20)	0.7(4)
C(18)-C(19)-C(20)-C(21)	-0.1(4)
C(19)-C(20)-C(21)-C(22)	-0.3(4)
C(20)-C(21)-C(22)-C(17)	0.0(4)
C(18)-C(17)-C(22)-C(21)	0.7(3)
C(12)-C(17)-C(22)-C(21)	-179.2(2)
Co(1)-O(5)-C(23)-C(24)	-2.7(3)
Co(1)-O(5)-C(23)-C(28)	175.61(14)
O(5)-C(23)-C(24)-C(25)	1.2(4)
C(28)-C(23)-C(24)-C(25)	-177.0(2)
C(27)-N(3)-C(25)-O(6)	5.6(3)
Co(1)-N(3)-C(25)-O(6)	167.83(15)

C(27)-N(3)-C(25)-C(24)	-173.8(2)
Co(1)-N(3)-C(25)-C(24)	-11.5(4)
C(26)-O(6)-C(25)-N(3)	-2.7(3)
C(26)-O(6)-C(25)-C(24)	176.7(2)
C(23)-C(24)-C(25)-N(3)	6.7(4)
C(23)-C(24)-C(25)-O(6)	-172.6(2)
C(25)-O(6)-C(26)-C(27)	-1.1(3)
C(25)-N(3)-C(27)-C(26)	-5.7(3)
Co(1)-N(3)-C(27)-C(26)	-167.71(16)
O(6)-C(26)-C(27)-N(3)	3.9(3)
O(5)-C(23)-C(28)-C(33)	29.1(3)
C(24)-C(23)-C(28)-C(33)	-152.5(2)
O(5)-C(23)-C(28)-C(29)	-146.4(2)
C(24)-C(23)-C(28)-C(29)	32.0(3)
C(33)-C(28)-C(29)-C(30)	-0.4(4)
C(23)-C(28)-C(29)-C(30)	175.3(2)
C(28)-C(29)-C(30)-C(31)	1.1(4)
C(29)-C(30)-C(31)-C(32)	-1.1(4)
C(30)-C(31)-C(32)-C(33)	0.6(4)
C(31)-C(32)-C(33)-C(28)	0.1(4)
C(29)-C(28)-C(33)-C(32)	-0.1(4)
C(23)-C(28)-C(33)-C(32)	-175.9(2)

Symmetry transformations used to generate equivalent atoms:

F3: Crystal data and structure refinement for d1842_a (**11c**).

Identification code	d1842_a (11c)	
Empirical formula	C ₃₆ H ₃₈ Co N ₃ O ₁₀	
Formula weight	731.62	
Temperature	150(2) K	
Wavelength	1.54178 Å	
Crystal system	Monoclinic	
Space group	P2 ₁ /c	
Unit cell dimensions	a = 8.778(4) Å	α = 90°.
	b = 24.578(11) Å	β = 95.956(9)°.
	c = 16.544(7) Å	γ = 90°.
Volume	3550(3) Å ³	
Z	4	
Density (calculated)	1.369 Mg/m ³	
Absorption coefficient	4.306 mm ⁻¹	
F(000)	1528	
Crystal size	0.180 x 0.160 x 0.020 mm ³	
Theta range for data collection	3.232 to 66.669°.	
Index ranges	-7 ≤ h ≤ 10, -28 ≤ k ≤ 29, -19 ≤ l ≤ 19	
Reflections collected	17977	
Independent reflections	6112 [R(int) = 0.0888]	
Completeness to theta = 66.669°	97.2 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.7528 and 0.4703	

Refinement method	Full-matrix least-squares on F^2
Data / restraints / parameters	6112 / 0 / 454
Goodness-of-fit on F^2	1.031
Final R indices [$I > 2\sigma(I)$]	$R_1 = 0.0723$, $wR_2 = 0.1654$
R indices (all data)	$R_1 = 0.1186$, $wR_2 = 0.1896$
Extinction coefficient	n/a
Largest diff. peak and hole	0.591 and -0.590 $e.\text{\AA}^{-3}$

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\approx^2 \times 10^3$) for d1842_a. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	U(eq)
Co(1)	7223(1)	7132(1)	6801(1)	27(1)
O(1)	5813(4)	6646(1)	6213(2)	31(1)
O(2)	8318(4)	7420(2)	4501(2)	41(1)
O(3)	-118(4)	5565(2)	4434(2)	40(1)
O(4)	5778(3)	7693(1)	6527(2)	30(1)
O(5)	4645(4)	6946(1)	8697(2)	36(1)
O(6)	569(5)	9337(2)	5243(2)	52(1)
O(7)	8620(3)	6541(1)	6960(2)	29(1)
O(8)	10565(4)	7903(1)	8256(2)	31(1)
O(9)	12778(5)	4478(2)	7441(3)	62(1)
N(1)	8147(4)	7329(2)	5845(2)	28(1)
N(2)	6316(4)	6928(2)	7755(3)	31(1)
N(3)	8520(4)	7619(2)	7436(2)	28(1)
C(1)	5393(5)	6674(2)	5439(3)	29(1)
C(2)	6146(5)	6946(2)	4881(3)	30(1)
C(3)	7532(5)	7228(2)	5109(3)	30(1)
C(4)	9704(6)	7665(3)	4890(3)	48(2)
C(5)	9593(6)	7622(2)	5795(3)	40(1)
C(6)	3936(5)	6379(2)	5182(3)	31(1)
C(7)	2719(6)	6422(2)	5636(3)	36(1)
C(8)	1328(6)	6168(2)	5400(3)	37(1)
C(9)	1193(5)	5841(2)	4718(3)	32(1)
C(10)	2416(6)	5779(2)	4259(3)	40(1)
C(11)	3775(6)	6054(2)	4484(3)	40(1)
C(12)	-1395(6)	5602(2)	4901(4)	43(2)
C(13)	4579(5)	7804(2)	6899(3)	28(1)
C(14)	4234(5)	7582(2)	7613(3)	31(1)
C(15)	5118(5)	7158(2)	8005(3)	29(1)
C(16)	5750(6)	6529(2)	8985(3)	40(1)
C(17)	6836(6)	6488(2)	8325(3)	37(1)
C(18)	3532(5)	8211(2)	6458(3)	29(1)

C(19)	3482(5)	8257(2)	5621(3)	34(1)
C(20)	2514(5)	8630(2)	5192(3)	34(1)
C(21)	1591(6)	8963(2)	5615(3)	37(1)
C(22)	1642(6)	8925(2)	6443(3)	40(1)
C(23)	2588(5)	8554(2)	6872(3)	34(1)
C(24)	466(8)	9387(3)	4391(4)	56(2)
C(25)	9949(5)	6546(2)	7403(3)	28(1)
C(26)	10584(5)	6979(2)	7831(3)	30(1)
C(27)	9848(5)	7489(2)	7818(3)	28(1)
C(28)	9533(5)	8369(2)	8190(3)	37(1)
C(29)	8159(6)	8189(2)	7615(3)	35(1)
C(30)	10744(5)	6011(2)	7429(3)	29(1)
C(31)	12333(6)	5964(2)	7595(3)	37(1)
C(32)	13059(6)	5464(2)	7611(4)	45(2)
C(33)	12191(7)	4996(2)	7466(4)	42(1)
C(34)	10607(6)	5031(2)	7284(3)	37(1)
C(35)	9910(6)	5532(2)	7266(3)	34(1)
C(36)	14396(7)	4416(3)	7647(5)	80(3)
O(1W)	6101(6)	5506(2)	6513(4)	101(2)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for d1842_a.

Co(1)-N(3)	1.892(4)
Co(1)-O(4)	1.896(3)
Co(1)-O(7)	1.902(3)
Co(1)-N(2)	1.906(4)
Co(1)-O(1)	1.912(3)
Co(1)-N(1)	1.913(4)
O(1)-C(1)	1.296(5)
O(2)-C(3)	1.361(6)
O(2)-C(4)	1.448(6)
O(3)-C(9)	1.376(6)
O(3)-C(12)	1.429(6)
O(4)-C(13)	1.302(6)
O(5)-C(15)	1.362(6)
O(5)-C(16)	1.456(6)
O(6)-C(21)	1.383(6)
O(6)-C(24)	1.408(7)
O(7)-C(25)	1.312(5)
O(8)-C(27)	1.365(6)
O(8)-C(28)	1.457(6)
O(9)-C(33)	1.377(6)
O(9)-C(36)	1.434(7)
N(1)-C(3)	1.305(6)
N(1)-C(5)	1.469(6)
N(2)-C(15)	1.298(6)
N(2)-C(17)	1.477(6)
N(3)-C(27)	1.307(6)
N(3)-C(29)	1.473(6)
C(1)-C(2)	1.364(7)
C(1)-C(6)	1.494(7)
C(2)-C(3)	1.417(7)
C(2)-H(2A)	0.9500
C(4)-C(5)	1.513(7)
C(4)-H(4A)	0.9900
C(4)-H(4B)	0.9900

C(5)-H(5A)	0.9900
C(5)-H(5B)	0.9900
C(6)-C(7)	1.372(7)
C(6)-C(11)	1.399(7)
C(7)-C(8)	1.391(7)
C(7)-H(7A)	0.9500
C(8)-C(9)	1.380(7)
C(8)-H(8A)	0.9500
C(9)-C(10)	1.387(7)
C(10)-C(11)	1.388(7)
C(10)-H(10A)	0.9500
C(11)-H(11A)	0.9500
C(12)-H(12A)	0.9800
C(12)-H(12B)	0.9800
C(12)-H(12C)	0.9800
C(13)-C(14)	1.363(7)
C(13)-C(18)	1.496(7)
C(14)-C(15)	1.416(7)
C(14)-H(14A)	0.9500
C(16)-C(17)	1.525(7)
C(16)-H(16A)	0.9900
C(16)-H(16B)	0.9900
C(17)-H(17A)	0.9900
C(17)-H(17B)	0.9900
C(18)-C(19)	1.385(7)
C(18)-C(23)	1.409(7)
C(19)-C(20)	1.393(7)
C(19)-H(19A)	0.9500
C(20)-C(21)	1.392(7)
C(20)-H(20A)	0.9500
C(21)-C(22)	1.369(7)
C(22)-C(23)	1.379(7)
C(22)-H(22A)	0.9500
C(23)-H(23A)	0.9500
C(24)-H(24A)	0.9800
C(24)-H(24B)	0.9800

C(24)-H(24C)	0.9800
C(25)-C(26)	1.364(7)
C(25)-C(30)	1.488(7)
C(26)-C(27)	1.409(7)
C(26)-H(26A)	0.9500
C(28)-C(29)	1.522(7)
C(28)-H(28A)	0.9900
C(28)-H(28B)	0.9900
C(29)-H(29A)	0.9900
C(29)-H(29B)	0.9900
C(30)-C(35)	1.397(7)
C(30)-C(31)	1.399(7)
C(31)-C(32)	1.382(7)
C(31)-H(31A)	0.9500
C(32)-C(33)	1.386(8)
C(32)-H(32A)	0.9500
C(33)-C(34)	1.395(7)
C(34)-C(35)	1.374(7)
C(34)-H(34A)	0.9500
C(35)-H(35A)	0.9500
C(36)-H(36A)	0.9800
C(36)-H(36B)	0.9800
C(36)-H(36C)	0.9800
O(1W)-H(1WA)	0.8420
O(1W)-H(1WB)	0.8477

N(3)-Co(1)-O(4)	91.27(16)
N(3)-Co(1)-O(7)	93.73(16)
O(4)-Co(1)-O(7)	173.77(14)
N(3)-Co(1)-N(2)	89.22(17)
O(4)-Co(1)-N(2)	93.58(16)
O(7)-Co(1)-N(2)	90.21(16)
N(3)-Co(1)-O(1)	176.07(16)
O(4)-Co(1)-O(1)	87.13(15)
O(7)-Co(1)-O(1)	88.11(14)
N(2)-Co(1)-O(1)	87.30(16)

N(3)-Co(1)-N(1)	90.92(17)
O(4)-Co(1)-N(1)	87.06(16)
O(7)-Co(1)-N(1)	89.14(16)
N(2)-Co(1)-N(1)	179.35(18)
O(1)-Co(1)-N(1)	92.58(16)
C(1)-O(1)-Co(1)	124.9(3)
C(3)-O(2)-C(4)	106.4(4)
C(9)-O(3)-C(12)	117.3(4)
C(13)-O(4)-Co(1)	126.4(3)
C(15)-O(5)-C(16)	107.0(4)
C(21)-O(6)-C(24)	118.4(4)
C(25)-O(7)-Co(1)	126.3(3)
C(27)-O(8)-C(28)	107.3(4)
C(33)-O(9)-C(36)	117.1(5)
C(3)-N(1)-C(5)	108.4(4)
C(3)-N(1)-Co(1)	123.6(3)
C(5)-N(1)-Co(1)	127.9(3)
C(15)-N(2)-C(17)	108.8(4)
C(15)-N(2)-Co(1)	124.8(4)
C(17)-N(2)-Co(1)	126.4(3)
C(27)-N(3)-C(29)	109.6(4)
C(27)-N(3)-Co(1)	124.6(4)
C(29)-N(3)-Co(1)	125.7(3)
O(1)-C(1)-C(2)	125.9(5)
O(1)-C(1)-C(6)	113.8(4)
C(2)-C(1)-C(6)	120.3(4)
C(1)-C(2)-C(3)	121.5(5)
C(1)-C(2)-H(2A)	119.2
C(3)-C(2)-H(2A)	119.2
N(1)-C(3)-O(2)	115.6(4)
N(1)-C(3)-C(2)	127.0(5)
O(2)-C(3)-C(2)	117.4(4)
O(2)-C(4)-C(5)	105.9(4)
O(2)-C(4)-H(4A)	110.6
C(5)-C(4)-H(4A)	110.6
O(2)-C(4)-H(4B)	110.6

C(5)-C(4)-H(4B)	110.6
H(4A)-C(4)-H(4B)	108.7
N(1)-C(5)-C(4)	103.6(4)
N(1)-C(5)-H(5A)	111.0
C(4)-C(5)-H(5A)	111.0
N(1)-C(5)-H(5B)	111.0
C(4)-C(5)-H(5B)	111.0
H(5A)-C(5)-H(5B)	109.0
C(7)-C(6)-C(11)	118.6(5)
C(7)-C(6)-C(1)	119.9(5)
C(11)-C(6)-C(1)	121.5(5)
C(6)-C(7)-C(8)	121.5(5)
C(6)-C(7)-H(7A)	119.2
C(8)-C(7)-H(7A)	119.2
C(9)-C(8)-C(7)	119.2(5)
C(9)-C(8)-H(8A)	120.4
C(7)-C(8)-H(8A)	120.4
O(3)-C(9)-C(8)	124.3(5)
O(3)-C(9)-C(10)	115.2(5)
C(8)-C(9)-C(10)	120.5(5)
C(9)-C(10)-C(11)	119.5(5)
C(9)-C(10)-H(10A)	120.3
C(11)-C(10)-H(10A)	120.3
C(10)-C(11)-C(6)	120.6(5)
C(10)-C(11)-H(11A)	119.7
C(6)-C(11)-H(11A)	119.7
O(3)-C(12)-H(12A)	109.5
O(3)-C(12)-H(12B)	109.5
H(12A)-C(12)-H(12B)	109.5
O(3)-C(12)-H(12C)	109.5
H(12A)-C(12)-H(12C)	109.5
H(12B)-C(12)-H(12C)	109.5
O(4)-C(13)-C(14)	126.0(5)
O(4)-C(13)-C(18)	113.3(4)
C(14)-C(13)-C(18)	120.7(4)
C(13)-C(14)-C(15)	122.1(5)

C(13)-C(14)-H(14A)	118.9
C(15)-C(14)-H(14A)	118.9
N(2)-C(15)-O(5)	115.4(4)
N(2)-C(15)-C(14)	126.7(5)
O(5)-C(15)-C(14)	117.8(4)
O(5)-C(16)-C(17)	104.9(4)
O(5)-C(16)-H(16A)	110.8
C(17)-C(16)-H(16A)	110.8
O(5)-C(16)-H(16B)	110.8
C(17)-C(16)-H(16B)	110.8
H(16A)-C(16)-H(16B)	108.8
N(2)-C(17)-C(16)	103.5(4)
N(2)-C(17)-H(17A)	111.1
C(16)-C(17)-H(17A)	111.1
N(2)-C(17)-H(17B)	111.1
C(16)-C(17)-H(17B)	111.1
H(17A)-C(17)-H(17B)	109.0
C(19)-C(18)-C(23)	118.6(5)
C(19)-C(18)-C(13)	119.8(5)
C(23)-C(18)-C(13)	121.7(5)
C(18)-C(19)-C(20)	121.1(5)
C(18)-C(19)-H(19A)	119.4
C(20)-C(19)-H(19A)	119.4
C(21)-C(20)-C(19)	119.2(5)
C(21)-C(20)-H(20A)	120.4
C(19)-C(20)-H(20A)	120.4
C(22)-C(21)-O(6)	116.3(5)
C(22)-C(21)-C(20)	120.2(5)
O(6)-C(21)-C(20)	123.4(5)
C(21)-C(22)-C(23)	121.0(5)
C(21)-C(22)-H(22A)	119.5
C(23)-C(22)-H(22A)	119.5
C(22)-C(23)-C(18)	119.9(5)
C(22)-C(23)-H(23A)	120.0
C(18)-C(23)-H(23A)	120.0
O(6)-C(24)-H(24A)	109.5

O(6)-C(24)-H(24B)	109.5
H(24A)-C(24)-H(24B)	109.5
O(6)-C(24)-H(24C)	109.5
H(24A)-C(24)-H(24C)	109.5
H(24B)-C(24)-H(24C)	109.5
O(7)-C(25)-C(26)	126.1(5)
O(7)-C(25)-C(30)	113.2(4)
C(26)-C(25)-C(30)	120.6(4)
C(25)-C(26)-C(27)	121.5(5)
C(25)-C(26)-H(26A)	119.2
C(27)-C(26)-H(26A)	119.2
N(3)-C(27)-O(8)	114.4(4)
N(3)-C(27)-C(26)	127.6(5)
O(8)-C(27)-C(26)	118.0(4)
O(8)-C(28)-C(29)	105.3(4)
O(8)-C(28)-H(28A)	110.7
C(29)-C(28)-H(28A)	110.7
O(8)-C(28)-H(28B)	110.7
C(29)-C(28)-H(28B)	110.7
H(28A)-C(28)-H(28B)	108.8
N(3)-C(29)-C(28)	103.3(4)
N(3)-C(29)-H(29A)	111.1
C(28)-C(29)-H(29A)	111.1
N(3)-C(29)-H(29B)	111.1
C(28)-C(29)-H(29B)	111.1
H(29A)-C(29)-H(29B)	109.1
C(35)-C(30)-C(31)	117.4(5)
C(35)-C(30)-C(25)	120.3(4)
C(31)-C(30)-C(25)	122.3(5)
C(32)-C(31)-C(30)	121.7(5)
C(32)-C(31)-H(31A)	119.2
C(30)-C(31)-H(31A)	119.2
C(31)-C(32)-C(33)	119.3(5)
C(31)-C(32)-H(32A)	120.3
C(33)-C(32)-H(32A)	120.3
O(9)-C(33)-C(32)	124.9(5)

O(9)-C(33)-C(34)	114.6(5)
C(32)-C(33)-C(34)	120.3(5)
C(35)-C(34)-C(33)	119.4(5)
C(35)-C(34)-H(34A)	120.3
C(33)-C(34)-H(34A)	120.3
C(34)-C(35)-C(30)	121.8(5)
C(34)-C(35)-H(35A)	119.1
C(30)-C(35)-H(35A)	119.1
O(9)-C(36)-H(36A)	109.5
O(9)-C(36)-H(36B)	109.5
H(36A)-C(36)-H(36B)	109.5
O(9)-C(36)-H(36C)	109.5
H(36A)-C(36)-H(36C)	109.5
H(36B)-C(36)-H(36C)	109.5
H(1WA)-O(1W)-H(1WB)	106.5

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\approx \times 10^3$) for d1842_a. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Co(1)	17(1)	37(1)	27(1)	-1(1)	4(1)	-1(1)
O(1)	24(2)	42(2)	27(2)	2(2)	1(2)	-2(2)
O(2)	32(2)	61(3)	30(2)	-1(2)	9(2)	-14(2)
O(3)	24(2)	51(2)	45(2)	-2(2)	-2(2)	-7(2)
O(4)	19(2)	44(2)	28(2)	1(2)	8(2)	1(2)
O(5)	29(2)	47(2)	34(2)	7(2)	11(2)	4(2)
O(6)	56(3)	61(3)	38(2)	7(2)	2(2)	26(2)
O(7)	19(2)	40(2)	29(2)	0(2)	2(1)	-1(2)
O(8)	23(2)	38(2)	32(2)	-7(2)	2(2)	-1(2)
O(9)	41(2)	44(3)	97(4)	-2(2)	-8(2)	13(2)
N(1)	20(2)	33(2)	32(2)	-5(2)	4(2)	-2(2)
N(2)	21(2)	36(2)	36(3)	1(2)	1(2)	-1(2)
N(3)	23(2)	33(2)	28(2)	0(2)	6(2)	2(2)
C(1)	19(2)	46(3)	23(3)	-4(2)	1(2)	3(2)
C(2)	17(2)	50(3)	23(3)	-5(2)	1(2)	-4(2)
C(3)	24(3)	41(3)	26(3)	-5(2)	8(2)	2(2)
C(4)	38(3)	70(4)	36(3)	-3(3)	4(3)	-22(3)
C(5)	30(3)	57(4)	36(3)	-2(3)	11(2)	-10(3)
C(6)	21(3)	38(3)	33(3)	2(2)	-1(2)	-2(2)
C(7)	30(3)	46(3)	32(3)	-7(3)	4(2)	-6(3)
C(8)	23(3)	50(3)	40(3)	0(3)	6(2)	-1(2)
C(9)	21(3)	39(3)	35(3)	3(2)	-6(2)	-8(2)
C(10)	33(3)	50(4)	35(3)	-9(3)	0(3)	-2(3)
C(11)	25(3)	57(4)	40(3)	-8(3)	5(2)	-4(3)
C(12)	23(3)	52(4)	53(4)	2(3)	2(3)	-9(3)
C(13)	18(2)	40(3)	25(3)	-4(2)	-1(2)	-2(2)
C(14)	19(2)	44(3)	31(3)	2(2)	9(2)	3(2)
C(15)	22(2)	36(3)	27(3)	-2(2)	1(2)	-6(2)
C(16)	31(3)	53(4)	37(3)	10(3)	5(2)	4(3)
C(17)	33(3)	50(3)	30(3)	6(3)	6(2)	-3(3)
C(18)	14(2)	38(3)	34(3)	2(2)	3(2)	-2(2)

C(19)	22(3)	49(3)	31(3)	-1(2)	7(2)	3(2)
C(20)	30(3)	43(3)	29(3)	2(2)	-1(2)	3(2)
C(21)	32(3)	39(3)	41(3)	2(3)	4(3)	7(2)
C(22)	36(3)	49(4)	34(3)	2(3)	7(3)	9(3)
C(23)	26(3)	47(3)	30(3)	-1(2)	5(2)	2(2)
C(24)	64(4)	63(4)	41(4)	9(3)	0(3)	18(3)
C(25)	19(2)	34(3)	32(3)	-1(2)	7(2)	0(2)
C(26)	21(3)	36(3)	33(3)	-4(2)	-2(2)	1(2)
C(27)	22(3)	38(3)	25(3)	-1(2)	3(2)	-4(2)
C(28)	27(3)	34(3)	49(4)	-6(3)	5(3)	3(2)
C(29)	27(3)	34(3)	43(3)	-6(2)	3(2)	0(2)
C(30)	24(3)	35(3)	28(3)	-2(2)	1(2)	1(2)
C(31)	24(3)	38(3)	47(3)	-2(3)	-1(2)	0(2)
C(32)	30(3)	44(3)	62(4)	-2(3)	-1(3)	9(3)
C(33)	42(3)	37(3)	48(4)	-1(3)	1(3)	9(3)
C(34)	39(3)	36(3)	35(3)	1(2)	1(3)	1(2)
C(35)	29(3)	42(3)	31(3)	1(2)	6(2)	0(2)
C(36)	43(4)	54(4)	140(8)	-3(5)	-7(4)	20(3)
O(1W)	77(4)	78(4)	148(6)	18(4)	8(4)	0(3)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\approx^2 \times 10^{-3}$) for d1842_a.

	x	y	z	U(eq)
H(2A)	5733	6945	4327	36
H(4A)	9777	8051	4726	58
H(4B)	10618	7469	4740	58
H(5A)	10470	7415	6067	48
H(5B)	9564	7987	6046	48
H(7A)	2829	6631	6122	43
H(8A)	480	6218	5706	45
H(10A)	2324	5551	3793	47
H(11A)	4604	6021	4161	49
H(12A)	-2247	5386	4640	64
H(12B)	-1713	5983	4932	64
H(12C)	-1102	5463	5450	64
H(14A)	3372	7715	7854	37
H(16A)	5232	6177	9055	48
H(16B)	6312	6636	9510	48
H(17A)	7911	6545	8554	45
H(17B)	6749	6129	8054	45
H(19A)	4118	8030	5334	40
H(20A)	2484	8656	4618	41
H(22A)	1017	9158	6727	47
H(23A)	2602	8530	7446	41
H(24A)	-325	9654	4210	85
H(24B)	198	9034	4140	85
H(24C)	1454	9508	4229	85
H(26A)	11544	6935	8146	36
H(28A)	10035	8688	7967	44
H(28B)	9215	8466	8728	44
H(29A)	7201	8214	7880	42
H(29B)	8055	8412	7114	42
H(31A)	12930	6283	7700	44

H(32A)	14141	5441	7720	54
H(34A)	10014	4711	7174	44
H(35A)	8831	5554	7139	40
H(36A)	14678	4033	7589	120
H(36B)	14670	4532	8209	120
H(36C)	14946	4640	7282	120
H(1WA)	5934	5808	6281	152
H(1WB)	5620	5270	6213	152

Table 6. Torsion angles [$^{\circ}$] for d1842_a.

N(3)-Co(1)-O(4)-C(13)	-94.3(4)
N(2)-Co(1)-O(4)-C(13)	-5.0(4)
O(1)-Co(1)-O(4)-C(13)	82.1(4)
N(1)-Co(1)-O(4)-C(13)	174.8(4)
O(4)-Co(1)-N(3)-C(27)	-178.4(4)
O(7)-Co(1)-N(3)-C(27)	-2.1(4)
N(2)-Co(1)-N(3)-C(27)	88.0(4)
N(1)-Co(1)-N(3)-C(27)	-91.3(4)
O(4)-Co(1)-N(3)-C(29)	5.4(4)
O(7)-Co(1)-N(3)-C(29)	-178.3(4)
N(2)-Co(1)-N(3)-C(29)	-88.2(4)
N(1)-Co(1)-N(3)-C(29)	92.5(4)
Co(1)-O(1)-C(1)-C(2)	18.6(7)
Co(1)-O(1)-C(1)-C(6)	-160.6(3)
O(1)-C(1)-C(2)-C(3)	0.9(8)
C(6)-C(1)-C(2)-C(3)	-179.9(4)
C(5)-N(1)-C(3)-O(2)	-1.5(6)
Co(1)-N(1)-C(3)-O(2)	177.1(3)
C(5)-N(1)-C(3)-C(2)	178.8(5)
Co(1)-N(1)-C(3)-C(2)	-2.6(7)
C(4)-O(2)-C(3)-N(1)	3.0(6)
C(4)-O(2)-C(3)-C(2)	-177.3(5)
C(1)-C(2)-C(3)-N(1)	-9.5(8)
C(1)-C(2)-C(3)-O(2)	170.7(5)
C(3)-O(2)-C(4)-C(5)	-3.1(6)
C(3)-N(1)-C(5)-C(4)	-0.6(6)
Co(1)-N(1)-C(5)-C(4)	-179.1(4)
O(2)-C(4)-C(5)-N(1)	2.2(6)
O(1)-C(1)-C(6)-C(7)	43.1(7)
C(2)-C(1)-C(6)-C(7)	-136.2(5)
O(1)-C(1)-C(6)-C(11)	-136.7(5)
C(2)-C(1)-C(6)-C(11)	44.0(7)
C(11)-C(6)-C(7)-C(8)	-2.3(8)
C(1)-C(6)-C(7)-C(8)	177.9(5)

C(6)-C(7)-C(8)-C(9)	3.5(8)
C(12)-O(3)-C(9)-C(8)	-2.9(7)
C(12)-O(3)-C(9)-C(10)	178.1(5)
C(7)-C(8)-C(9)-O(3)	179.2(5)
C(7)-C(8)-C(9)-C(10)	-1.8(8)
O(3)-C(9)-C(10)-C(11)	178.2(5)
C(8)-C(9)-C(10)-C(11)	-0.8(8)
C(9)-C(10)-C(11)-C(6)	1.9(8)
C(7)-C(6)-C(11)-C(10)	-0.4(8)
C(1)-C(6)-C(11)-C(10)	179.4(5)
Co(1)-O(4)-C(13)-C(14)	8.3(7)
Co(1)-O(4)-C(13)-C(18)	-170.8(3)
O(4)-C(13)-C(14)-C(15)	-5.3(8)
C(18)-C(13)-C(14)-C(15)	173.7(5)
C(17)-N(2)-C(15)-O(5)	-0.3(6)
Co(1)-N(2)-C(15)-O(5)	179.4(3)
C(17)-N(2)-C(15)-C(14)	-177.2(5)
Co(1)-N(2)-C(15)-C(14)	2.4(7)
C(16)-O(5)-C(15)-N(2)	3.8(6)
C(16)-O(5)-C(15)-C(14)	-179.0(4)
C(13)-C(14)-C(15)-N(2)	-0.4(8)
C(13)-C(14)-C(15)-O(5)	-177.3(5)
C(15)-O(5)-C(16)-C(17)	-5.4(5)
C(15)-N(2)-C(17)-C(16)	-3.1(6)
Co(1)-N(2)-C(17)-C(16)	177.2(3)
O(5)-C(16)-C(17)-N(2)	5.1(5)
O(4)-C(13)-C(18)-C(19)	27.4(6)
C(14)-C(13)-C(18)-C(19)	-151.7(5)
O(4)-C(13)-C(18)-C(23)	-152.5(4)
C(14)-C(13)-C(18)-C(23)	28.4(7)
C(23)-C(18)-C(19)-C(20)	-0.4(7)
C(13)-C(18)-C(19)-C(20)	179.7(4)
C(18)-C(19)-C(20)-C(21)	0.3(8)
C(24)-O(6)-C(21)-C(22)	-179.5(5)
C(24)-O(6)-C(21)-C(20)	-0.7(8)
C(19)-C(20)-C(21)-C(22)	0.3(8)

C(19)-C(20)-C(21)-O(6)	-178.5(5)
O(6)-C(21)-C(22)-C(23)	178.0(5)
C(20)-C(21)-C(22)-C(23)	-0.8(8)
C(21)-C(22)-C(23)-C(18)	0.7(8)
C(19)-C(18)-C(23)-C(22)	-0.1(8)
C(13)-C(18)-C(23)-C(22)	179.8(5)
Co(1)-O(7)-C(25)-C(26)	-0.4(7)
Co(1)-O(7)-C(25)-C(30)	177.7(3)
O(7)-C(25)-C(26)-C(27)	-1.7(8)
C(30)-C(25)-C(26)-C(27)	-179.7(5)
C(29)-N(3)-C(27)-O(8)	-1.6(6)
Co(1)-N(3)-C(27)-O(8)	-178.3(3)
C(29)-N(3)-C(27)-C(26)	177.6(5)
Co(1)-N(3)-C(27)-C(26)	0.9(7)
C(28)-O(8)-C(27)-N(3)	3.3(5)
C(28)-O(8)-C(27)-C(26)	-176.0(4)
C(25)-C(26)-C(27)-N(3)	1.5(8)
C(25)-C(26)-C(27)-O(8)	-179.3(5)
C(27)-O(8)-C(28)-C(29)	-3.4(5)
C(27)-N(3)-C(29)-C(28)	-0.7(5)
Co(1)-N(3)-C(29)-C(28)	176.0(3)
O(8)-C(28)-C(29)-N(3)	2.5(5)
O(7)-C(25)-C(30)-C(35)	-22.5(7)
C(26)-C(25)-C(30)-C(35)	155.8(5)
O(7)-C(25)-C(30)-C(31)	155.9(5)
C(26)-C(25)-C(30)-C(31)	-25.8(8)
C(35)-C(30)-C(31)-C(32)	-1.0(8)
C(25)-C(30)-C(31)-C(32)	-179.4(5)
C(30)-C(31)-C(32)-C(33)	-0.6(9)
C(36)-O(9)-C(33)-C(32)	5.3(9)
C(36)-O(9)-C(33)-C(34)	-178.2(6)
C(31)-C(32)-C(33)-O(9)	178.1(6)
C(31)-C(32)-C(33)-C(34)	1.7(9)
O(9)-C(33)-C(34)-C(35)	-177.9(5)
C(32)-C(33)-C(34)-C(35)	-1.2(9)
C(33)-C(34)-C(35)-C(30)	-0.4(8)

C(31)-C(30)-C(35)-C(34)	1.5(8)
C(25)-C(30)-C(35)-C(34)	180.0(5)

Symmetry transformations used to generate equivalent atoms:

Table 7. Hydrogen bonds for d1842_a [$\approx$ and ∞].

D-H...A	d(D-H)	d(H...A)	d(D...A)	$\angle$ (DHA)
O(1W)-H(1WA)...O(1)	0.84	2.07	2.853(6)	155.6

Symmetry transformations used to generate equivalent atoms:

F4: Crystal data and structure refinement for d1870ra_b (**9j**).

Identification code	d1870ra_b (9j)	
Empirical formula	C ₁₆ H ₂₄ Co N ₂ O ₄	
Formula weight	367.30	
Temperature	150(2) K	
Wavelength	0.71073 Å	
Crystal system	Orthorhombic	
Space group	Pca2 ₁	
Unit cell dimensions	a = 10.6455(5) Å	a = 90°.
	b = 14.6865(9) Å	b = 90°.
	c = 11.3187(6) Å	g = 90°.
Volume	1769.62(17) Å ³	
Z	4	
Density (calculated)	1.379 Mg/m ³	
Absorption coefficient	0.990 mm ⁻¹	
F(000)	772	
Crystal size	0.190 x 0.190 x 0.100 mm ³	
Theta range for data collection	2.363 to 27.547°.	
Index ranges	-13 ≤ h ≤ 12, -19 ≤ k ≤ 18, -14 ≤ l ≤ 14	
Reflections collected	18782	
Independent reflections	4071 [R(int) = 0.0458]	
Completeness to theta = 25.242°	99.9 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.7456 and 0.6801	
Refinement method	Full-matrix least-squares on F ²	

Data / restraints / parameters	4071 / 1 / 214
Goodness-of-fit on F^2	1.035
Final R indices [$I > 2\sigma(I)$]	$R_1 = 0.0323$, $wR_2 = 0.0637$
R indices (all data)	$R_1 = 0.0701$, $wR_2 = 0.0763$
Absolute structure parameter	0.023(11)
Extinction coefficient	n/a
Largest diff. peak and hole	0.297 and -0.405 e. $\text{\AA}^{-3}$

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d1870ra_b. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	$U(\text{eq})$
Co(1)	4978(1)	2528(1)	4394(1)	17(1)
O(1)	5491(2)	1478(2)	3465(2)	23(1)
O(2)	2345(3)	724(2)	5598(3)	42(1)
O(3)	4592(2)	3653(2)	3580(2)	25(1)
O(4)	7672(3)	4134(2)	5934(2)	38(1)
N(1)	3510(3)	1932(2)	5092(2)	23(1)
N(2)	6347(3)	3034(2)	5342(2)	19(1)
C(1)	3353(3)	1059(3)	4995(3)	27(1)
C(2)	4077(3)	422(2)	4347(4)	27(1)
C(3)	5070(4)	660(3)	3626(3)	23(1)
C(4)	1886(4)	1472(4)	6329(4)	50(2)
C(5)	2440(4)	2322(3)	5751(4)	31(1)
C(6)	5743(4)	-66(3)	2936(4)	35(1)
C(7)	2900(4)	3030(4)	6646(4)	56(2)
C(8)	1536(4)	2767(3)	4892(4)	35(1)
C(9)	5153(4)	4424(3)	3755(4)	25(1)
C(10)	6134(3)	4566(3)	4528(4)	30(1)
C(11)	6671(3)	3889(3)	5245(3)	24(1)
C(12)	7953(4)	3363(3)	6689(4)	40(1)
C(13)	7221(3)	2563(3)	6154(4)	25(1)
C(14)	4656(5)	5197(3)	3032(4)	45(1)
C(15)	6501(4)	2023(4)	7075(4)	49(1)
C(16)	8066(4)	1933(3)	5460(4)	35(1)

Table 3. Bond lengths [Å] and angles [°] for d1870ra_b.

Co(1)-O(3)	1.935(3)
Co(1)-O(1)	1.944(3)
Co(1)-N(2)	1.957(3)
Co(1)-N(1)	1.958(3)
O(1)-C(3)	1.295(4)
O(2)-C(1)	1.364(4)
O(2)-C(4)	1.459(6)
O(3)-C(9)	1.296(5)
O(4)-C(11)	1.368(4)
O(4)-C(12)	1.450(5)
N(1)-C(1)	1.297(5)
N(1)-C(5)	1.478(5)
N(2)-C(11)	1.307(5)
N(2)-C(13)	1.480(5)
C(1)-C(2)	1.417(6)
C(2)-C(3)	1.380(5)
C(2)-H(2A)	0.9500
C(3)-C(6)	1.503(5)
C(4)-C(5)	1.529(6)
C(4)-H(4A)	0.9900
C(4)-H(4B)	0.9900
C(5)-C(8)	1.516(5)
C(5)-C(7)	1.532(6)
C(6)-H(6A)	0.9800
C(6)-H(6B)	0.9800
C(6)-H(6C)	0.9800
C(7)-H(7A)	0.9800
C(7)-H(7B)	0.9800
C(7)-H(7C)	0.9800
C(8)-H(8A)	0.9800
C(8)-H(8B)	0.9800
C(8)-H(8C)	0.9800
C(9)-C(10)	1.378(5)
C(9)-C(14)	1.497(6)

C(10)-C(11)	1.405(6)
C(10)-H(10A)	0.9500
C(12)-C(13)	1.535(6)
C(12)-H(12A)	0.9900
C(12)-H(12B)	0.9900
C(13)-C(16)	1.510(6)
C(13)-C(15)	1.517(6)
C(14)-H(14A)	0.9800
C(14)-H(14B)	0.9800
C(14)-H(14C)	0.9800
C(15)-H(15A)	0.9800
C(15)-H(15B)	0.9800
C(15)-H(15C)	0.9800
C(16)-H(16A)	0.9800
C(16)-H(16B)	0.9800
C(16)-H(16C)	0.9800

O(3)-Co(1)-O(1)	118.64(13)
O(3)-Co(1)-N(2)	95.42(13)
O(1)-Co(1)-N(2)	112.90(11)
O(3)-Co(1)-N(1)	113.84(12)
O(1)-Co(1)-N(1)	95.03(13)
N(2)-Co(1)-N(1)	122.89(16)
C(3)-O(1)-Co(1)	124.2(2)
C(1)-O(2)-C(4)	106.0(3)
C(9)-O(3)-Co(1)	125.2(2)
C(11)-O(4)-C(12)	107.0(3)
C(1)-N(1)-C(5)	109.0(3)
C(1)-N(1)-Co(1)	120.7(3)
C(5)-N(1)-Co(1)	130.2(3)
C(11)-N(2)-C(13)	109.7(3)
C(11)-N(2)-Co(1)	121.0(3)
C(13)-N(2)-Co(1)	129.1(3)
N(1)-C(1)-O(2)	114.6(4)
N(1)-C(1)-C(2)	128.7(3)
O(2)-C(1)-C(2)	116.7(4)

C(3)-C(2)-C(1)	123.8(3)
C(3)-C(2)-H(2A)	118.1
C(1)-C(2)-H(2A)	118.1
O(1)-C(3)-C(2)	125.7(4)
O(1)-C(3)-C(6)	114.8(4)
C(2)-C(3)-C(6)	119.5(4)
O(2)-C(4)-C(5)	104.1(3)
O(2)-C(4)-H(4A)	110.9
C(5)-C(4)-H(4A)	110.9
O(2)-C(4)-H(4B)	110.9
C(5)-C(4)-H(4B)	110.9
H(4A)-C(4)-H(4B)	109.0
N(1)-C(5)-C(8)	109.5(3)
N(1)-C(5)-C(4)	101.4(3)
C(8)-C(5)-C(4)	112.4(3)
N(1)-C(5)-C(7)	110.5(3)
C(8)-C(5)-C(7)	109.5(4)
C(4)-C(5)-C(7)	113.3(4)
C(3)-C(6)-H(6A)	109.5
C(3)-C(6)-H(6B)	109.5
H(6A)-C(6)-H(6B)	109.5
C(3)-C(6)-H(6C)	109.5
H(6A)-C(6)-H(6C)	109.5
H(6B)-C(6)-H(6C)	109.5
C(5)-C(7)-H(7A)	109.5
C(5)-C(7)-H(7B)	109.5
H(7A)-C(7)-H(7B)	109.5
C(5)-C(7)-H(7C)	109.5
H(7A)-C(7)-H(7C)	109.5
H(7B)-C(7)-H(7C)	109.5
C(5)-C(8)-H(8A)	109.5
C(5)-C(8)-H(8B)	109.5
H(8A)-C(8)-H(8B)	109.5
C(5)-C(8)-H(8C)	109.5
H(8A)-C(8)-H(8C)	109.5
H(8B)-C(8)-H(8C)	109.5

O(3)-C(9)-C(10)	125.4(4)
O(3)-C(9)-C(14)	114.6(4)
C(10)-C(9)-C(14)	120.0(4)
C(9)-C(10)-C(11)	124.6(4)
C(9)-C(10)-H(10A)	117.7
C(11)-C(10)-H(10A)	117.7
N(2)-C(11)-O(4)	114.2(4)
N(2)-C(11)-C(10)	128.4(4)
O(4)-C(11)-C(10)	117.4(4)
O(4)-C(12)-C(13)	105.1(3)
O(4)-C(12)-H(12A)	110.7
C(13)-C(12)-H(12A)	110.7
O(4)-C(12)-H(12B)	110.7
C(13)-C(12)-H(12B)	110.7
H(12A)-C(12)-H(12B)	108.8
N(2)-C(13)-C(16)	109.7(3)
N(2)-C(13)-C(15)	110.7(3)
C(16)-C(13)-C(15)	109.8(4)
N(2)-C(13)-C(12)	101.9(3)
C(16)-C(13)-C(12)	111.8(3)
C(15)-C(13)-C(12)	112.7(4)
C(9)-C(14)-H(14A)	109.5
C(9)-C(14)-H(14B)	109.5
H(14A)-C(14)-H(14B)	109.5
C(9)-C(14)-H(14C)	109.5
H(14A)-C(14)-H(14C)	109.5
H(14B)-C(14)-H(14C)	109.5
C(13)-C(15)-H(15A)	109.5
C(13)-C(15)-H(15B)	109.5
H(15A)-C(15)-H(15B)	109.5
C(13)-C(15)-H(15C)	109.5
H(15A)-C(15)-H(15C)	109.5
H(15B)-C(15)-H(15C)	109.5
C(13)-C(16)-H(16A)	109.5
C(13)-C(16)-H(16B)	109.5
H(16A)-C(16)-H(16B)	109.5

C(13)-C(16)-H(16C)	109.5
H(16A)-C(16)-H(16C)	109.5
H(16B)-C(16)-H(16C)	109.5

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d1870ra_b. The anisotropic displacement factor exponent takes the form: $-2p^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Co(1)	15(1)	19(1)	16(1)	-1(1)	0(1)	-1(1)
O(1)	22(1)	22(2)	24(2)	-4(1)	2(1)	-4(1)
O(2)	27(2)	46(2)	52(2)	25(2)	9(2)	-7(1)
O(3)	24(1)	23(2)	27(2)	4(1)	-2(1)	0(1)
O(4)	34(2)	37(2)	42(2)	-11(2)	-11(2)	-11(1)
N(1)	19(2)	32(2)	18(2)	5(2)	2(1)	0(2)
N(2)	19(2)	25(2)	14(2)	-4(2)	-1(1)	3(1)
C(1)	18(2)	38(3)	25(2)	17(2)	-2(2)	-5(2)
C(2)	28(2)	19(2)	35(2)	5(2)	-5(2)	-7(2)
C(3)	27(2)	18(2)	25(2)	-3(2)	-12(2)	-2(2)
C(4)	29(2)	69(4)	52(3)	32(3)	21(2)	15(2)
C(5)	22(2)	49(3)	22(2)	7(2)	10(2)	4(2)
C(6)	41(3)	25(3)	40(3)	-6(2)	-7(2)	4(2)
C(7)	39(3)	96(5)	33(3)	-23(3)	5(2)	9(3)
C(8)	26(2)	38(3)	41(3)	6(2)	4(2)	5(2)
C(9)	31(2)	19(2)	25(2)	1(2)	8(2)	6(2)
C(10)	37(2)	16(2)	36(2)	-5(2)	1(2)	-3(2)
C(11)	27(2)	25(2)	22(2)	-11(2)	2(2)	-2(2)
C(12)	35(2)	46(3)	38(3)	-19(2)	-16(2)	9(2)
C(13)	22(2)	34(2)	19(2)	-1(2)	-7(2)	3(2)
C(14)	50(3)	29(3)	55(3)	14(2)	-9(3)	2(2)
C(15)	38(3)	80(4)	31(3)	20(3)	-6(2)	1(3)
C(16)	30(2)	38(3)	37(3)	-7(2)	-10(2)	13(2)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for d1870ra_b.

	x	y	z	U(eq)
H(2A)	3866	-204	4411	33
H(4A)	2182	1408	7154	60
H(4B)	956	1494	6326	60
H(6A)	5480	-667	3218	53
H(6B)	6651	2	3043	53
H(6C)	5535	-5	2096	53
H(7A)	2199	3217	7152	84
H(7B)	3229	3562	6224	84
H(7C)	3566	2764	7133	84
H(8A)	861	3068	5333	53
H(8B)	1175	2302	4372	53
H(8C)	1986	3218	4416	53
H(10A)	6470	5164	4581	35
H(12A)	7677	3482	7510	48
H(12B)	8866	3235	6692	48
H(14A)	5095	5760	3249	67
H(14B)	3754	5269	3181	67
H(14C)	4793	5070	2191	67
H(15A)	7095	1738	7624	74
H(15B)	6003	1550	6683	74
H(15C)	5941	2432	7511	74
H(16A)	8638	1618	6002	53
H(16B)	8555	2288	4889	53
H(16C)	7553	1484	5039	53

Table 6. Torsion angles [°] for d1870ra_b.

C(5)-N(1)-C(1)-O(2)	-5.8(4)
Co(1)-N(1)-C(1)-O(2)	175.2(2)
C(5)-N(1)-C(1)-C(2)	173.7(4)
Co(1)-N(1)-C(1)-C(2)	-5.3(5)
C(4)-O(2)-C(1)-N(1)	-9.2(4)
C(4)-O(2)-C(1)-C(2)	171.3(3)
N(1)-C(1)-C(2)-C(3)	-4.2(6)
O(2)-C(1)-C(2)-C(3)	175.3(4)
Co(1)-O(1)-C(3)-C(2)	10.0(5)
Co(1)-O(1)-C(3)-C(6)	-170.9(2)
C(1)-C(2)-C(3)-O(1)	1.6(6)
C(1)-C(2)-C(3)-C(6)	-177.4(4)
C(1)-O(2)-C(4)-C(5)	19.3(4)
C(1)-N(1)-C(5)-C(8)	-101.7(4)
Co(1)-N(1)-C(5)-C(8)	77.2(4)
C(1)-N(1)-C(5)-C(4)	17.2(4)
Co(1)-N(1)-C(5)-C(4)	-163.9(3)
C(1)-N(1)-C(5)-C(7)	137.6(4)
Co(1)-N(1)-C(5)-C(7)	-43.6(5)
O(2)-C(4)-C(5)-N(1)	-21.6(4)
O(2)-C(4)-C(5)-C(8)	95.2(4)
O(2)-C(4)-C(5)-C(7)	-140.0(4)
Co(1)-O(3)-C(9)-C(10)	1.5(6)
Co(1)-O(3)-C(9)-C(14)	-178.6(3)
O(3)-C(9)-C(10)-C(11)	0.0(7)
C(14)-C(9)-C(10)-C(11)	-179.9(4)
C(13)-N(2)-C(11)-O(4)	-2.5(4)
Co(1)-N(2)-C(11)-O(4)	-177.8(2)
C(13)-N(2)-C(11)-C(10)	176.9(4)
Co(1)-N(2)-C(11)-C(10)	1.6(5)
C(12)-O(4)-C(11)-N(2)	-7.5(4)
C(12)-O(4)-C(11)-C(10)	173.0(3)
C(9)-C(10)-C(11)-N(2)	-1.8(7)
C(9)-C(10)-C(11)-O(4)	177.6(4)

C(11)-O(4)-C(12)-C(13)	13.6(4)
C(11)-N(2)-C(13)-C(16)	-108.0(4)
Co(1)-N(2)-C(13)-C(16)	66.8(4)
C(11)-N(2)-C(13)-C(15)	130.6(4)
Co(1)-N(2)-C(13)-C(15)	-54.5(5)
C(11)-N(2)-C(13)-C(12)	10.6(4)
Co(1)-N(2)-C(13)-C(12)	-174.6(3)
O(4)-C(12)-C(13)-N(2)	-14.4(4)
O(4)-C(12)-C(13)-C(16)	102.8(4)
O(4)-C(12)-C(13)-C(15)	-133.0(4)

Symmetry transformations used to generate equivalent atoms:

F5: Crystal data and structure refinement for d1867_a (**9k**).

Identification code	d1867_a (9k)	
Empirical formula	C ₁₆ H ₁₈ Co F ₆ N ₂ O ₄	
Formula weight	475.25	
Temperature	150(2) K	
Wavelength	0.71073 Å	
Crystal system	Orthorhombic	
Space group	Pbca	
Unit cell dimensions	a = 11.3622(5) Å	a = 90°.
	b = 18.1752(10) Å	b = 90°.
	c = 18.9609(9) Å	g = 90°.
Volume	3915.6(3) Å ³	
Z	8	
Density (calculated)	1.612 Mg/m ³	
Absorption coefficient	0.957 mm ⁻¹	
F(000)	1928	
Crystal size	0.270 x 0.190 x 0.060 mm ³	
Theta range for data collection	2.148 to 27.520°.	
Index ranges	-14 ≤ h ≤ 8, -17 ≤ k ≤ 23, -24 ≤ l ≤ 17	
Reflections collected	18516	
Independent reflections	4500 [R(int) = 0.0536]	
Completeness to theta = 25.242°	100.0 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.7456 and 0.6827	

Refinement method	Full-matrix least-squares on F^2
Data / restraints / parameters	4500 / 0 / 266
Goodness-of-fit on F^2	1.015
Final R indices [$I > 2\sigma(I)$]	$R_1 = 0.0415$, $wR_2 = 0.0833$
R indices (all data)	$R_1 = 0.0795$, $wR_2 = 0.0981$
Extinction coefficient	n/a
Largest diff. peak and hole	0.557 and -0.406 e. $\text{\AA}^{-3}$

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d1867_a. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	$U(\text{eq})$
Co(1)	3872(1)	3396(1)	6266(1)	20(1)
F(1)	2850(2)	1072(1)	6924(1)	59(1)
F(2)	1374(2)	1696(1)	7303(1)	60(1)
F(3)	2673(2)	1297(1)	8021(1)	46(1)
F(4)	2658(2)	5652(1)	5310(2)	110(1)
F(5)	2049(3)	5169(2)	4391(1)	99(1)
F(6)	1170(2)	4987(1)	5331(1)	72(1)
O(1)	3063(2)	2543(1)	6632(1)	31(1)
O(2)	4921(2)	3547(1)	8328(1)	41(1)
O(3)	2897(2)	4166(1)	5858(1)	30(1)
O(4)	5094(2)	3306(1)	4232(1)	36(1)
N(1)	4576(2)	3673(1)	7174(1)	23(1)
N(2)	4715(2)	3181(1)	5385(1)	24(1)
C(1)	3198(2)	2297(2)	7261(2)	25(1)
C(2)	3815(2)	2596(2)	7802(2)	28(1)
C(3)	4433(2)	3277(2)	7738(1)	26(1)
C(4)	5338(3)	4280(2)	8161(2)	48(1)
C(5)	5319(2)	4314(2)	7357(2)	26(1)
C(6)	2523(3)	1589(2)	7383(2)	32(1)
C(7)	6533(3)	4206(2)	7041(2)	52(1)
C(8)	4758(3)	5025(2)	7091(2)	43(1)
C(9)	2997(2)	4392(2)	5225(2)	26(1)
C(10)	3706(2)	4131(2)	4704(2)	30(1)
C(11)	4505(2)	3533(2)	4802(1)	26(1)
C(12)	5950(3)	2770(2)	4479(2)	36(1)
C(13)	5525(2)	2557(2)	5215(2)	27(1)
C(14)	2220(3)	5046(2)	5063(2)	35(1)
C(15)	6513(3)	2524(2)	5750(2)	46(1)
C(16)	4809(3)	1844(2)	5205(2)	46(1)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for d1867_a.

Co(1)-O(1)	1.9315(19)
Co(1)-O(3)	1.9449(19)
Co(1)-N(1)	1.964(2)
Co(1)-N(2)	1.964(2)
F(1)-C(6)	1.334(3)
F(2)-C(6)	1.329(3)
F(3)-C(6)	1.331(3)
F(4)-C(14)	1.296(4)
F(5)-C(14)	1.308(4)
F(6)-C(14)	1.301(4)
O(1)-C(1)	1.283(3)
O(2)-C(3)	1.342(3)
O(2)-C(4)	1.449(4)
O(3)-C(9)	1.275(3)
O(4)-C(11)	1.337(3)
O(4)-C(12)	1.455(4)
N(1)-C(3)	1.300(3)
N(1)-C(5)	1.480(3)
N(2)-C(11)	1.299(3)
N(2)-C(13)	1.494(3)
C(1)-C(2)	1.357(4)
C(1)-C(6)	1.516(4)
C(2)-C(3)	1.427(4)
C(2)-H(2A)	0.9500
C(4)-C(5)	1.525(4)
C(4)-H(4A)	0.9900
C(4)-H(4B)	0.9900
C(5)-C(7)	1.517(4)
C(5)-C(8)	1.528(4)
C(7)-H(7A)	0.9800
C(7)-H(7B)	0.9800
C(7)-H(7C)	0.9800
C(8)-H(8A)	0.9800
C(8)-H(8B)	0.9800

C(8)-H(8C)	0.9800
C(9)-C(10)	1.360(4)
C(9)-C(14)	1.512(4)
C(10)-C(11)	1.428(4)
C(10)-H(10A)	0.9500
C(12)-C(13)	1.528(4)
C(12)-H(12A)	0.9900
C(12)-H(12B)	0.9900
C(13)-C(15)	1.513(4)
C(13)-C(16)	1.531(4)
C(15)-H(15A)	0.9800
C(15)-H(15B)	0.9800
C(15)-H(15C)	0.9800
C(16)-H(16A)	0.9800
C(16)-H(16B)	0.9800
C(16)-H(16C)	0.9800

O(1)-Co(1)-O(3)	116.69(8)
O(1)-Co(1)-N(1)	94.87(9)
O(3)-Co(1)-N(1)	113.29(9)
O(1)-Co(1)-N(2)	112.19(9)
O(3)-Co(1)-N(2)	94.78(9)
N(1)-Co(1)-N(2)	126.71(9)
C(1)-O(1)-Co(1)	123.83(18)
C(3)-O(2)-C(4)	106.8(2)
C(9)-O(3)-Co(1)	123.78(18)
C(11)-O(4)-C(12)	106.3(2)
C(3)-N(1)-C(5)	108.3(2)
C(3)-N(1)-Co(1)	121.86(19)
C(5)-N(1)-Co(1)	129.84(18)
C(11)-N(2)-C(13)	107.7(2)
C(11)-N(2)-Co(1)	122.36(19)
C(13)-N(2)-Co(1)	129.33(18)
O(1)-C(1)-C(2)	128.7(3)
O(1)-C(1)-C(6)	112.2(2)
C(2)-C(1)-C(6)	119.1(3)

C(1)-C(2)-C(3)	122.5(3)
C(1)-C(2)-H(2A)	118.8
C(3)-C(2)-H(2A)	118.8
N(1)-C(3)-O(2)	115.6(3)
N(1)-C(3)-C(2)	127.8(3)
O(2)-C(3)-C(2)	116.6(2)
O(2)-C(4)-C(5)	104.5(2)
O(2)-C(4)-H(4A)	110.8
C(5)-C(4)-H(4A)	110.8
O(2)-C(4)-H(4B)	110.8
C(5)-C(4)-H(4B)	110.8
H(4A)-C(4)-H(4B)	108.9
N(1)-C(5)-C(7)	108.9(2)
N(1)-C(5)-C(4)	102.2(2)
C(7)-C(5)-C(4)	112.2(3)
N(1)-C(5)-C(8)	110.5(2)
C(7)-C(5)-C(8)	111.0(3)
C(4)-C(5)-C(8)	111.7(3)
F(2)-C(6)-F(3)	106.7(2)
F(2)-C(6)-F(1)	107.6(3)
F(3)-C(6)-F(1)	106.1(2)
F(2)-C(6)-C(1)	110.8(2)
F(3)-C(6)-C(1)	114.4(2)
F(1)-C(6)-C(1)	111.0(2)
C(5)-C(7)-H(7A)	109.5
C(5)-C(7)-H(7B)	109.5
H(7A)-C(7)-H(7B)	109.5
C(5)-C(7)-H(7C)	109.5
H(7A)-C(7)-H(7C)	109.5
H(7B)-C(7)-H(7C)	109.5
C(5)-C(8)-H(8A)	109.5
C(5)-C(8)-H(8B)	109.5
H(8A)-C(8)-H(8B)	109.5
C(5)-C(8)-H(8C)	109.5
H(8A)-C(8)-H(8C)	109.5
H(8B)-C(8)-H(8C)	109.5

O(3)-C(9)-C(10)	128.6(3)
O(3)-C(9)-C(14)	113.2(3)
C(10)-C(9)-C(14)	118.3(3)
C(9)-C(10)-C(11)	123.3(3)
C(9)-C(10)-H(10A)	118.4
C(11)-C(10)-H(10A)	118.4
N(2)-C(11)-O(4)	116.4(3)
N(2)-C(11)-C(10)	127.0(3)
O(4)-C(11)-C(10)	116.6(2)
O(4)-C(12)-C(13)	104.6(2)
O(4)-C(12)-H(12A)	110.8
C(13)-C(12)-H(12A)	110.8
O(4)-C(12)-H(12B)	110.8
C(13)-C(12)-H(12B)	110.8
H(12A)-C(12)-H(12B)	108.9
N(2)-C(13)-C(15)	110.0(2)
N(2)-C(13)-C(12)	101.5(2)
C(15)-C(13)-C(12)	112.8(2)
N(2)-C(13)-C(16)	108.6(2)
C(15)-C(13)-C(16)	111.7(3)
C(12)-C(13)-C(16)	111.7(3)
F(4)-C(14)-F(6)	106.3(3)
F(4)-C(14)-F(5)	105.4(3)
F(6)-C(14)-F(5)	105.0(3)
F(4)-C(14)-C(9)	111.7(3)
F(6)-C(14)-C(9)	113.0(3)
F(5)-C(14)-C(9)	114.7(3)
C(13)-C(15)-H(15A)	109.5
C(13)-C(15)-H(15B)	109.5
H(15A)-C(15)-H(15B)	109.5
C(13)-C(15)-H(15C)	109.5
H(15A)-C(15)-H(15C)	109.5
H(15B)-C(15)-H(15C)	109.5
C(13)-C(16)-H(16A)	109.5
C(13)-C(16)-H(16B)	109.5
H(16A)-C(16)-H(16B)	109.5

C(13)-C(16)-H(16C)	109.5
H(16A)-C(16)-H(16C)	109.5
H(16B)-C(16)-H(16C)	109.5

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d1867_a. The anisotropic displacement factor exponent takes the form: $-2p^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Co(1)	19(1)	25(1)	15(1)	2(1)	-1(1)	1(1)
F(1)	88(2)	36(1)	52(1)	-11(1)	16(1)	-16(1)
F(2)	31(1)	54(1)	94(2)	29(1)	-7(1)	-14(1)
F(3)	60(1)	39(1)	40(1)	20(1)	2(1)	-5(1)
F(4)	82(2)	31(1)	216(4)	-13(2)	-59(2)	10(1)
F(5)	124(2)	127(2)	47(1)	40(2)	8(1)	78(2)
F(6)	41(1)	76(2)	98(2)	45(1)	15(1)	28(1)
O(1)	32(1)	36(1)	23(1)	9(1)	-8(1)	-10(1)
O(2)	60(1)	39(1)	24(1)	-1(1)	-16(1)	-7(1)
O(3)	28(1)	39(1)	21(1)	8(1)	1(1)	8(1)
O(4)	39(1)	49(1)	20(1)	-4(1)	6(1)	3(1)
N(1)	22(1)	26(1)	22(1)	0(1)	0(1)	0(1)
N(2)	24(1)	27(1)	22(1)	-2(1)	0(1)	-1(1)
C(1)	21(1)	29(2)	27(2)	4(1)	2(1)	2(1)
C(2)	35(2)	27(2)	21(1)	7(1)	-1(1)	3(1)
C(3)	26(1)	32(2)	19(1)	-2(1)	-4(1)	5(1)
C(4)	64(2)	43(2)	38(2)	-5(2)	-12(2)	-11(2)
C(5)	22(1)	27(2)	30(2)	-6(1)	-3(1)	-2(1)
C(6)	34(2)	31(2)	33(2)	3(2)	1(1)	-3(1)
C(7)	23(2)	52(2)	81(3)	-23(2)	3(2)	-9(2)
C(8)	32(2)	31(2)	66(2)	2(2)	-6(2)	1(1)
C(9)	25(1)	28(2)	25(2)	6(1)	-5(1)	-4(1)
C(10)	37(2)	34(2)	20(1)	8(1)	-1(1)	-5(1)
C(11)	30(2)	33(2)	16(1)	-3(1)	3(1)	-8(1)
C(12)	35(2)	41(2)	33(2)	-11(2)	6(1)	0(2)
C(13)	25(1)	30(2)	27(2)	-6(1)	4(1)	3(1)
C(14)	38(2)	33(2)	34(2)	12(1)	-5(1)	2(1)
C(15)	40(2)	58(2)	41(2)	-9(2)	-4(2)	22(2)
C(16)	53(2)	30(2)	56(2)	-6(2)	12(2)	-1(2)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for d1867_a.

	x	y	z	U(eq)
H(2A)	3836	2344	8241	33
H(4A)	6145	4357	8343	58
H(4B)	4812	4658	8366	58
H(7A)	6888	3758	7235	78
H(7B)	7029	4631	7156	78
H(7C)	6466	4160	6528	78
H(8A)	4708	5012	6575	64
H(8B)	5241	5446	7237	64
H(8C)	3966	5075	7290	64
H(10A)	3669	4357	4253	36
H(12A)	6747	2988	4499	44
H(12B)	5968	2335	4164	44
H(15A)	6937	2994	5752	69
H(15B)	7056	2127	5624	69
H(15C)	6185	2432	6219	69
H(16A)	4456	1762	5670	69
H(16B)	5327	1430	5088	69
H(16C)	4185	1882	4849	69

Table 6. Torsion angles [°] for d1867_a.

Co(1)-O(1)-C(1)-C(2)	6.3(4)
Co(1)-O(1)-C(1)-C(6)	-175.66(17)
O(1)-C(1)-C(2)-C(3)	0.5(5)
C(6)-C(1)-C(2)-C(3)	-177.4(3)
C(5)-N(1)-C(3)-O(2)	2.9(3)
Co(1)-N(1)-C(3)-O(2)	-177.28(18)
C(5)-N(1)-C(3)-C(2)	-177.7(3)
Co(1)-N(1)-C(3)-C(2)	2.2(4)
C(4)-O(2)-C(3)-N(1)	7.9(3)
C(4)-O(2)-C(3)-C(2)	-171.6(3)
C(1)-C(2)-C(3)-N(1)	-5.3(5)
C(1)-C(2)-C(3)-O(2)	174.2(3)
C(3)-O(2)-C(4)-C(5)	-14.6(3)
C(3)-N(1)-C(5)-C(7)	107.2(3)
Co(1)-N(1)-C(5)-C(7)	-72.6(3)
C(3)-N(1)-C(5)-C(4)	-11.6(3)
Co(1)-N(1)-C(5)-C(4)	168.6(2)
C(3)-N(1)-C(5)-C(8)	-130.7(3)
Co(1)-N(1)-C(5)-C(8)	49.5(3)
O(2)-C(4)-C(5)-N(1)	15.6(3)
O(2)-C(4)-C(5)-C(7)	-100.9(3)
O(2)-C(4)-C(5)-C(8)	133.7(3)
O(1)-C(1)-C(6)-F(2)	-61.9(3)
C(2)-C(1)-C(6)-F(2)	116.4(3)
O(1)-C(1)-C(6)-F(3)	177.5(2)
C(2)-C(1)-C(6)-F(3)	-4.2(4)
O(1)-C(1)-C(6)-F(1)	57.6(3)
C(2)-C(1)-C(6)-F(1)	-124.2(3)
Co(1)-O(3)-C(9)-C(10)	4.7(4)
Co(1)-O(3)-C(9)-C(14)	-174.11(18)
O(3)-C(9)-C(10)-C(11)	-1.0(5)
C(14)-C(9)-C(10)-C(11)	177.8(3)
C(13)-N(2)-C(11)-O(4)	-4.7(3)
Co(1)-N(2)-C(11)-O(4)	-176.49(18)

C(13)-N(2)-C(11)-C(10)	174.7(3)
Co(1)-N(2)-C(11)-C(10)	2.9(4)
C(12)-O(4)-C(11)-N(2)	-7.9(3)
C(12)-O(4)-C(11)-C(10)	172.7(2)
C(9)-C(10)-C(11)-N(2)	-3.4(5)
C(9)-C(10)-C(11)-O(4)	176.0(3)
C(11)-O(4)-C(12)-C(13)	16.4(3)
C(11)-N(2)-C(13)-C(15)	133.9(3)
Co(1)-N(2)-C(13)-C(15)	-55.0(3)
C(11)-N(2)-C(13)-C(12)	14.3(3)
Co(1)-N(2)-C(13)-C(12)	-174.70(18)
C(11)-N(2)-C(13)-C(16)	-103.6(3)
Co(1)-N(2)-C(13)-C(16)	67.5(3)
O(4)-C(12)-C(13)-N(2)	-18.2(3)
O(4)-C(12)-C(13)-C(15)	-135.8(3)
O(4)-C(12)-C(13)-C(16)	97.4(3)
O(3)-C(9)-C(14)-F(4)	78.0(4)
C(10)-C(9)-C(14)-F(4)	-101.0(4)
O(3)-C(9)-C(14)-F(6)	-41.9(4)
C(10)-C(9)-C(14)-F(6)	139.1(3)
O(3)-C(9)-C(14)-F(5)	-162.2(3)
C(10)-C(9)-C(14)-F(5)	18.8(4)

Symmetry transformations used to generate equivalent atoms:

F6: Crystal data and structure refinement for d1889a_a (**91**).

Identification code	d1889a_a (91)	
Empirical formula	C ₁₆ H ₁₈ Co Cl ₆ N ₂ O ₄	
Formula weight	573.95	
Temperature	150(2) K	
Wavelength	0.71073 Å	
Crystal system	Orthorhombic	
Space group	Pna2 ₁	
Unit cell dimensions	a = 32.2380(14) Å	a = 90°.
	b = 8.4769(4) Å	b = 90°.
	c = 8.3149(3) Å	g = 90°.
Volume	2272.28(17) Å ³	
Z	4	
Density (calculated)	1.678 Mg/m ³	
Absorption coefficient	1.486 mm ⁻¹	
F(000)	1156	
Crystal size	0.21 x 0.060 x 0.020 mm ³	
Theta range for data collection	2.484 to 27.551°.	
Index ranges	-30 ≤ h ≤ 41, -10 ≤ k ≤ 11, -7 ≤ l ≤ 10	
Reflections collected	11915	
Independent reflections	4617 [R(int) = 0.0419]	
Completeness to theta = 25.242°	99.8 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.7456 and 0.6777	
Refinement method	Full-matrix least-squares on F ²	

Data / restraints / parameters	4617 / 1 / 266
Goodness-of-fit on F^2	1.021
Final R indices [$I > 2\sigma(I)$]	$R_1 = 0.0378$, $wR_2 = 0.0677$
R indices (all data)	$R_1 = 0.0532$, $wR_2 = 0.0716$
Extinction coefficient	n/a
Largest diff. peak and hole	0.445 and -0.482 e. $\text{\AA}^{-3}$

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d1867_a. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	$U(\text{eq})$
Co(1)	6271(1)	5501(1)	4559(1)	20(1)
Cl(1)	5248(1)	7932(2)	1365(2)	58(1)
Cl(2)	4752(1)	5513(2)	3129(3)	69(1)
Cl(3)	7709(1)	8768(2)	4032(2)	51(1)
Cl(4)	7791(1)	5629(2)	786(2)	34(1)
Cl(5)	7309(1)	6452(2)	4142(1)	32(1)
Cl(6)	5759(1)	8400(2)	1987(2)	33(1)
O(1)	5759(1)	6270(4)	3612(4)	27(1)
O(2)	5812(1)	7957(5)	8414(4)	36(1)
O(3)	6785(1)	6312(4)	3684(4)	27(1)
O(4)	6726(1)	1408(4)	2606(4)	25(1)
N(1)	6156(1)	6409(5)	6684(4)	19(1)
N(2)	6381(1)	3309(5)	3909(5)	20(1)
C(1)	5482(1)	7066(5)	4342(6)	18(1)
C(2)	5496(1)	7615(6)	5890(6)	21(1)
C(3)	5834(2)	7276(6)	6943(5)	20(1)
C(4)	6188(2)	7519(10)	9241(7)	72(3)
C(5)	6415(2)	6373(6)	8169(6)	24(1)
C(6)	5097(1)	7336(6)	3311(6)	25(1)
C(7)	6849(2)	6919(11)	7773(8)	70(3)
C(8)	6413(3)	4755(9)	8849(9)	84(3)
C(9)	7058(1)	5481(5)	2954(5)	17(1)
C(10)	7044(1)	3920(6)	2632(6)	19(1)
C(11)	6708(1)	2925(5)	3091(5)	18(1)
C(12)	6368(2)	616(7)	3258(8)	40(1)
C(13)	6103(1)	1909(6)	4066(6)	26(1)
C(14)	7444(1)	6444(6)	2458(6)	21(1)
C(15)	5703(1)	2218(8)	3182(10)	64(2)
C(16)	6030(1)	1559(8)	5811(8)	55(2)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for d1867_a.

Co(1)-O(3)	1.936(3)
Co(1)-O(1)	1.940(3)
Co(1)-N(1)	1.963(4)
Co(1)-N(2)	1.968(4)
Cl(1)-C(6)	1.764(5)
Cl(2)-C(6)	1.767(5)
Cl(3)-C(6)	1.752(5)
C;(4)-C(14)	1.772(5)
Cl(5)-C(14)	1.791(5)
Cl(6)-C(14)	1.758(5)
O(1)-C(1)	1.274(5)
O(2)-C(3)	1.354(6)
O(2)-C(4)	1.443(6)
O(3)-C(9)	1.280(5)
O(4)-C(11)	1.349(5)
O(4)-C(12)	1.441(6)
N(1)-C(3)	1.290(6)
N(1)-C(5)	1.490(6)
N(2)-C(11)	1.294(6)
N(2)-C(13)	1.493(6)
C(1)-C(2)	1.370(6)
C(1)-C(6)	1.526(6)
C(2)-C(3)	1.427(6)
C(2)-H(2A)	0.9500
C(4)-C(5)	1.508(7)
C(4)-H(4A)	0.9900
C(4)-H(4B)	0.9900
C(5)-C(7)	1.483(8)
C(5)-C(8)	1.510(8)
C(7)-H(7A)	0.9800
C(7)-H(7B)	0.9800
C(7)-H(7C)	0.9800
C(8)-H(8A)	0.9800
C(8)-H(8B)	0.9800

C(8)-H(8C)	0.9800
C(9)-C(10)	1.351(6)
C(9)-C(14)	1.546(6)
C(10)-C(11)	1.427(6)
C(10)-H(10A)	0.9500
C(12)-C(13)	1.542(7)
C(12)-H(12A)	0.9900
C(12)-H(12B)	0.9900
C(13)-C(15)	1.508(8)
C(13)-C(16)	1.500(8)
C(15)-H(15A)	0.9800
C(15)-H(15B)	0.9800
C(15)-H(15C)	0.9800
C(16)-H(16A)	0.9800
C(16)-H(16B)	0.9800
C(16)-H(16C)	0.9800

O(1)-Co(1)-O(3)	117.13(15)
O(3)-Co(1)-N(1)	111.12(15)
O(1)-Co(1)-N(1)	94.21(14)
O(3)-Co(1)-N(2)	94.42(15)
O(1)-Co(1)-N(2)	111.09(15)
N(1)-Co(1)-N(2)	130.67(17)
C(1)-O(1)-Co(1)	125.6(3)
C(3)-O(2)-C(4)	160.0(4)
C(9)-O(3)-Co(1)	124.8(3)
C(11)-O(4)-C(12)	107.3(4)
C(3)-N(1)-C(5)	108.9(4)
C(3)-N(1)-Co(1)	121.7(3)
C(5)-N(1)-Co(1)	129.2(3)
C(11)-N(2)-C(13)	109.5(4)
C(11)-N(2)-Co(1)	121.9(3)
C(13)-N(2)-Co(1)	1128.2(3)
O(1)-C(1)-C(2)	127.1(4)
O(1)-C(1)-C(6)	112.5(4)
C(2)-C(1)-C(6)	120.3(4)

C(1)-C(2)-C(3)	122.3(4)
C(1)-C(2)-H(2A)	118.8
C(3)-C(2)-H(2A)	118.8
N(1)-C(3)-O(2)	115.8(4)
N(1)-C(3)-C(2)	128.8(4)
O(2)-C(3)-C(2)	115.3(4)
O(2)-C(4)-C(5)	107.0(4)
O(2)-C(4)-H(4A)	110.3
C(5)-C(4)-H(4A)	110.3
O(2)-C(4)-H(4B)	110.3
C(5)-C(4)-H(4B)	110.3
H(4A)-C(4)-H(4B)	108.6
C(8)-C(5)-N(1)	109.4(4)
C(8)-C(5)-C(4)	111.7(6)
N(1)-C(5)-C(4)	101.8(4)
C(8)-C(5)-C(7)	111.7(6)
N(1)-C(5)-C(7)	109.3(4)
C(4)-C(5)-C(7)	112.4(6)
C(1)-C(6)-Cl(3)	115.3(3)
C(1)-C(6)-Cl(1)	109.4(3)
Cl(1)-C(6)-Cl(3)	106.9(3)
Cl(2)-C(6)-C(1)	107.8(3)
Cl(3)-C(6)-C(2)	109.5(3)
Cl(1)-C(6)-Cl(2)	107.7(3)
C(5)-C(7)-H(7A)	109.5
C(5)-C(7)-H(7B)	109.5
H(7A)-C(7)-H(7B)	109.5
C(5)-C(7)-H(7C)	109.5
H(7A)-C(7)-H(7C)	109.5
H(7B)-C(7)-H(7C)	109.5
C(5)-C(8)-H(8A)	109.5
C(5)-C(8)-H(8B)	109.5
H(8A)-C(8)-H(8B)	109.5
C(5)-C(8)-H(8C)	109.5
H(8A)-C(8)-H(8C)	109.5
H(8B)-C(8)-H(8C)	109.5

O(3)-C(9)-C(10)	127.7(4)
O(3)-C(9)-C(14)	112.9(4)
C(10)-C(9)-C(14)	119.3(4)
C(9)-C(10)-C(11)	123.4(4)
C(9)-C(10)-H(10A)	118.3
C(11)-C(10)-H(10A)	118.3
N(2)-C(11)-O(4)	115.6(4)
N(2)-C(11)-C(10)	127.6(4)
O(4)-C(11)-C(10)	116.8(4)
O(4)-C(12)-C(13)	106.0(4)
O(4)-C(12)-H(12A)	110.5
C(13)-C(12)-H(12A)	110.5
O(4)-C(12)-H(12B)	110.5
C(13)-C(12)-H(12B)	110.5
H(12A)-C(12)-H(12B)	108.7
N(2)-C(13)-C(16)	109.4(4)
N(2)-C(13)-C(15)	109.4(4)
C(16)-C(13)-C(15)	111.8(5)
C(12)-C(13)-C(16)	101.2(4)
C(15)-C(13)-C(12)	111.6(5)
C(9)-C(14)-Cl(6)	112.6(5)
C(9)-C(14)-Cl(4)	111.0(3)
Cl(6)-C(14)-Cl(4)	108.2(3)
C(9)-C(14)-Cl(5)	107.2(3)
Cl(6)-C(14)-Cl(5)	108.9(3)
Cl(4)-C(14)-Cl(5)	108.3(3)
C(13)-C(15)-H(15A)	109.5
C(13)-C(15)-H(15B)	109.5
H(15A)-C(15)-H(15B)	109.5
C(13)-C(15)-H(15C)	109.5
H(15A)-C(15)-H(15C)	109.5
H(15B)-C(15)-H(15C)	109.5
C(13)-C(16)-H(16A)	109.5
C(13)-C(16)-H(16B)	109.5
H(16A)-C(16)-H(16B)	109.5
C(13)-C(16)-H(16C)	109.5

H(16A)-C(16)-H(16C)	109.5
H(16B)-C(16)-H(16C)	109.5

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d1867_a. The anisotropic displacement factor exponent takes the form: $-2p^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Co(1)	16(1)	19(1)	24(1)	-5(1)	2(1)	2(1)
Cl(1)	47(1)	96(2)	29(1)	18(1)	-8(1)	8(1)
Cl(2)	40(1)	32(1)	135(2)	8(1)	-41(1)	-13(1)
Cl(3)	42(1)	64(1)	46(1)	-18(1)	-18(1)	34(1)
Cl(4)	34(1)	37(1)	31(1)	-5(1)	15(1)	-5(1)
Cl(5)	27(1)	38(1)	30(1)	2(1)	-8(1)	-8(2)
Cl(6)	38(1)	22(1)	38(1)	8(1)	3(1)	-1(1)
O(1)	25(2)	31(2)	24(2)	-6(2)	-3(1)	9(2)
O(2)	34(2)	55(3)	19(2)	-10(2)	-4(2)	21(2)
O(3)	22(2)	21(2)	38(2)	-7(2)	10(1)	-2(1)
O(4)	26(2)	14(2)	36(2)	-9(2)	6(2)	0(1)
N(1)	16(2)	22(2)	20(2)	-2(2)	-2(2)	4(2)
N(2)	15(2)	15(2)	29(2)	-3(2)	4(2)	0(2)
C(1)	14(2)	17(2)	24(3)	2(2)	4(2)	0(2)
C(2)	20(2)	21(3)	23(3)	0(2)	2(2)	6(2)
C(3)	26(3)	17(2)	15(2)	4(2)	4(2)	2(2)
C(4)	57(4)	125(7)	34(4)	-33(4)	-24(3)	61(4)
C(5)	24(2)	32(3)	18(3)	-3(2)	-7(2)	5(2)
C(6)	24(3)	21(3)	31(3)	-2(2)	-2(2)	3(2)
C(7)	42(4)	126(8)	42(4)	-20(5)	-11(3)	-31(4)
C(8)	129(7)	45(5)	78(6)	26(4)	-78(5)	-10(4)
C(9)	20(2)	18(2)	14(2)	-1(2)	-2(2)	0(2)
C(10)	18(2)	19(2)	19(3)	-3(2)	5(2)	1(2)
C(11)	20(2)	16(3)	19(3)	-4(2)	-9(2)	5(2)
C(12)	36(3)	25(3)	60(4)	-16(3)	13(3)	0(2)
C(13)	17(2)	18(3)	44(3)	-4(2)	3(2)	3(1)
C(14)	20(2)	22(3)	22(3)	-3(2)	5(2)	0(2)
C(15)	37(4)	35(4)	120(7)	-15(4)	-28(4)	-11(3)
C(16)	76(5)	33(4)	58(4)	-4(3)	26(4)	-25(3)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for d1867_a.

	x	y	z	U(eq)
H(2A)	5274	8244	6277	25
H(4A)	6122	7020	10286	86
H(4B)	6362	8463	9443	86
H(7A)	7023	6854	8738	105
H(7B)	6840	8013	7393	105
H(7C)	6966	9244	6930	105
H(8A)	6576	4739	9845	126
H(8B)	6536	4022	8071	126
H(8C)	6127	4436	9081	126
H(10A)	7271	3461	2071	23
H(12A)	6452	-188	4055	48
H(12B)	6209	91	2392	48
H(15A)	5522	1295	3274	96
H(15B)	5763	2422	2045	96
H(15C)	5565	3139	3651	96
H(16A)	5871	578	5913	83
H(16B)	5874	2428	6300	83
H(16C)	6297	1443	6361	83

F7: Crystal data and structure refinement for d1825_a (**7g**).

Identification code	d1825_a (7g)	
Empirical formula	C ₃₄ H ₃₂ Cu N ₂ O ₄	
Formula weight	596.15	
Temperature	150(2) K	
Wavelength	0.71073 $\approx$	
Crystal system	Triclinic	
Space group	P-1	
Unit cell dimensions	$a = 5.6548(4) \approx$	$\alpha = 86.961(2)^\circ$
	$b = 10.2976(8) \approx$	$\beta = 83.937(2)^\circ$
	$c = 12.5388(10) \approx$	$\gamma = 79.616(2)^\circ$
Volume	$713.74(9) \approx^3$	
Z	1	
Density (calculated)	1.387 Mg/m ³	
Absorption coefficient	0.807 mm ⁻¹	
F(000)	311	
Crystal size	0.230 x 0.200 x 0.040 mm ³	
Theta range for data collection	1.634 to 27.595 $^\circ$.	
Index ranges	$-7 \leq h \leq 7$, $-13 \leq k \leq 13$, $-16 \leq l \leq 16$	
Reflections collected	21074	
Independent reflections	3316 [R(int) = 0.0263]	
Completeness to theta = 25.242 $^\circ$	100.0 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.7456 and 0.6988	
Refinement method	Full-matrix least-squares on F ²	
Data / restraints / parameters	3316 / 0 / 189	

Goodness-of-fit on F^2	1.070
Final R indices [$I > 2\sigma(I)$]	$R_1 = 0.0266$, $wR_2 = 0.0648$
R indices (all data)	$R_1 = 0.0312$, $wR_2 = 0.0669$
Extinction coefficient	n/a
Largest diff. peak and hole	0.350 and -0.260 e. $\approx^3$

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\approx \times 10^3$) for d1825_a. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	U(eq)
Cu(1)	5000	5000	5000	15(1)
O(1)	3401(2)	6165(1)	6093(1)	24(1)
O(2)	7068(2)	8552(1)	3929(1)	26(1)
N(1)	6552(2)	6439(1)	4247(1)	16(1)
C(1)	8365(3)	6416(1)	3276(1)	18(1)
C(2)	9076(3)	7778(2)	3311(1)	27(1)
C(3)	5889(3)	7668(1)	4512(1)	18(1)
C(4)	4117(3)	8218(1)	5321(1)	22(1)
C(5)	3028(3)	7445(1)	6063(1)	18(1)
C(6)	1260(3)	8096(1)	6933(1)	19(1)
C(7)	-790(3)	8925(2)	6661(1)	24(1)
C(8)	-2554(3)	9522(2)	7455(1)	29(1)
C(9)	-2220(3)	9289(2)	8512(1)	31(1)
C(10)	-115(3)	8457(2)	8836(1)	25(1)
C(11)	275(4)	8216(2)	9932(1)	37(1)
C(12)	2320(4)	7431(2)	10224(1)	40(1)
C(13)	4105(3)	6851(2)	9441(1)	34(1)
C(14)	3802(3)	7051(2)	8373(1)	25(1)
C(15)	1662(3)	7846(1)	8039(1)	21(1)
C(16)	10577(3)	5342(2)	3347(2)	32(1)
C(17)	7101(3)	6302(2)	2281(1)	27(1)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for d1825_a.

Cu(1)-O(1)#1	1.9122(10)
Cu(1)-O(1)	1.9122(10)
Cu(1)-N(1)	2.0007(11)
Cu(1)-N(1)#1	2.0007(11)
O(1)-C(5)	1.2959(17)
O(2)-C(3)	1.3607(17)
O(2)-C(2)	1.4416(18)
N(1)-C(3)	1.3045(18)
N(1)-C(1)	1.5044(17)
C(1)-C(16)	1.520(2)
C(1)-C(17)	1.522(2)
C(1)-C(2)	1.531(2)
C(2)-H(2A)	0.9900
C(2)-H(2B)	0.9900
C(3)-C(4)	1.413(2)
C(4)-C(5)	1.363(2)
C(4)-H(4A)	0.9500
C(5)-C(6)	1.4984(19)
C(6)-C(7)	1.373(2)
C(6)-C(15)	1.431(2)
C(7)-C(8)	1.413(2)
C(7)-H(7A)	0.9500
C(8)-C(9)	1.362(2)
C(8)-H(8A)	0.9500
C(9)-C(10)	1.419(2)
C(9)-H(9A)	0.9500
C(10)-C(11)	1.416(2)
C(10)-C(15)	1.425(2)
C(11)-C(12)	1.358(3)
C(11)-H(11A)	0.9500
C(12)-C(13)	1.405(3)
C(12)-H(12A)	0.9500
C(13)-C(14)	1.368(2)
C(13)-H(13A)	0.9500

C(14)-C(15)	1.423(2)
C(14)-H(14A)	0.9500
C(16)-H(16A)	0.9800
C(16)-H(16B)	0.9800
C(16)-H(16C)	0.9800
C(17)-H(17A)	0.9800
C(17)-H(17B)	0.9800
C(17)-H(17C)	0.9800

O(1)#1-Cu(1)-O(1)	180.0
O(1)#1-Cu(1)-N(1)	88.54(5)
O(1)-Cu(1)-N(1)	91.46(5)
O(1)#1-Cu(1)-N(1)#1	91.46(5)
O(1)-Cu(1)-N(1)#1	88.54(5)
N(1)-Cu(1)-N(1)#1	180.0
C(5)-O(1)-Cu(1)	128.02(9)
C(3)-O(2)-C(2)	105.98(11)
C(3)-N(1)-C(1)	107.21(11)
C(3)-N(1)-Cu(1)	121.52(10)
C(1)-N(1)-Cu(1)	131.00(9)
N(1)-C(1)-C(16)	113.46(12)
N(1)-C(1)-C(17)	108.77(12)
C(16)-C(1)-C(17)	111.78(13)
N(1)-C(1)-C(2)	101.17(11)
C(16)-C(1)-C(2)	109.96(13)
C(17)-C(1)-C(2)	111.25(13)
O(2)-C(2)-C(1)	104.51(12)
O(2)-C(2)-H(2A)	110.8
C(1)-C(2)-H(2A)	110.8
O(2)-C(2)-H(2B)	110.8
C(1)-C(2)-H(2B)	110.8
H(2A)-C(2)-H(2B)	108.9
N(1)-C(3)-O(2)	115.67(13)
N(1)-C(3)-C(4)	129.24(13)
O(2)-C(3)-C(4)	115.08(12)
C(5)-C(4)-C(3)	121.69(13)

C(5)-C(4)-H(4A)	119.2
C(3)-C(4)-H(4A)	119.2
O(1)-C(5)-C(4)	125.40(13)
O(1)-C(5)-C(6)	115.83(12)
C(4)-C(5)-C(6)	118.77(12)
C(7)-C(6)-C(15)	119.89(14)
C(7)-C(6)-C(5)	119.24(13)
C(15)-C(6)-C(5)	120.86(13)
C(6)-C(7)-C(8)	121.24(15)
C(6)-C(7)-H(7A)	119.4
C(8)-C(7)-H(7A)	119.4
C(9)-C(8)-C(7)	119.91(15)
C(9)-C(8)-H(8A)	120.0
C(7)-C(8)-H(8A)	120.0
C(8)-C(9)-C(10)	121.08(15)
C(8)-C(9)-H(9A)	119.5
C(10)-C(9)-H(9A)	119.5
C(11)-C(10)-C(9)	121.85(15)
C(11)-C(10)-C(15)	118.89(15)
C(9)-C(10)-C(15)	119.26(14)
C(12)-C(11)-C(10)	120.90(17)
C(12)-C(11)-H(11A)	119.5
C(10)-C(11)-H(11A)	119.5
C(11)-C(12)-C(13)	120.48(16)
C(11)-C(12)-H(12A)	119.8
C(13)-C(12)-H(12A)	119.8
C(14)-C(13)-C(12)	120.70(17)
C(14)-C(13)-H(13A)	119.6
C(12)-C(13)-H(13A)	119.6
C(13)-C(14)-C(15)	120.34(15)
C(13)-C(14)-H(14A)	119.8
C(15)-C(14)-H(14A)	119.8
C(14)-C(15)-C(10)	118.66(14)
C(14)-C(15)-C(6)	122.69(14)
C(10)-C(15)-C(6)	118.61(14)
C(1)-C(16)-H(16A)	109.5

C(1)-C(16)-H(16B)	109.5
H(16A)-C(16)-H(16B)	109.5
C(1)-C(16)-H(16C)	109.5
H(16A)-C(16)-H(16C)	109.5
H(16B)-C(16)-H(16C)	109.5
C(1)-C(17)-H(17A)	109.5
C(1)-C(17)-H(17B)	109.5
H(17A)-C(17)-H(17B)	109.5
C(1)-C(17)-H(17C)	109.5
H(17A)-C(17)-H(17C)	109.5
H(17B)-C(17)-H(17C)	109.5

Symmetry transformations used to generate equivalent atoms:

#1 -x+1,-y+1,-z+1

Table 4. Anisotropic displacement parameters ($\approx \times 10^3$) for d1825_a. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Cu(1)	18(1)	12(1)	16(1)	-1(1)	1(1)	-3(1)
O(1)	34(1)	13(1)	23(1)	-2(1)	8(1)	-6(1)
O(2)	35(1)	16(1)	28(1)	-1(1)	8(1)	-10(1)
N(1)	17(1)	15(1)	16(1)	-1(1)	0(1)	-4(1)
C(1)	18(1)	18(1)	17(1)	0(1)	1(1)	-6(1)
C(2)	31(1)	24(1)	28(1)	-5(1)	8(1)	-14(1)
C(3)	22(1)	15(1)	19(1)	2(1)	-2(1)	-7(1)
C(4)	28(1)	12(1)	24(1)	-3(1)	2(1)	-4(1)
C(5)	20(1)	16(1)	18(1)	-2(1)	-2(1)	-4(1)
C(6)	23(1)	14(1)	21(1)	-2(1)	1(1)	-6(1)
C(7)	27(1)	20(1)	26(1)	-1(1)	-2(1)	-4(1)
C(8)	25(1)	22(1)	38(1)	-2(1)	2(1)	0(1)
C(9)	30(1)	27(1)	32(1)	-8(1)	11(1)	-5(1)
C(10)	31(1)	23(1)	23(1)	-4(1)	5(1)	-10(1)
C(11)	49(1)	41(1)	21(1)	-6(1)	5(1)	-13(1)
C(12)	54(1)	49(1)	21(1)	1(1)	-6(1)	-18(1)
C(13)	39(1)	36(1)	31(1)	4(1)	-12(1)	-11(1)
C(14)	29(1)	22(1)	25(1)	-2(1)	-2(1)	-8(1)
C(15)	26(1)	17(1)	21(1)	-2(1)	0(1)	-10(1)
C(16)	19(1)	34(1)	39(1)	6(1)	5(1)	-1(1)
C(17)	33(1)	33(1)	18(1)	2(1)	-2(1)	-14(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\approx^2 \times 10^{-3}$) for d1825_a.

	x	y	z	U(eq)
H(2A)	10582	7717	3662	33
H(2B)	9315	8171	2578	33
H(4A)	3673	9152	5348	26
H(7A)	-1028	9101	5925	29
H(8A)	-3971	10084	7251	35
H(9A)	-3415	9691	9042	37
H(11A)	-909	8608	10470	44
H(12A)	2544	7273	10963	48
H(13A)	5537	6315	9656	41
H(14A)	5027	6656	7852	30
H(16A)	10157	4484	3224	48
H(16B)	11140	5327	4061	48
H(16C)	11862	5524	2802	48
H(17A)	6444	5480	2334	41
H(17B)	8262	6296	1642	41
H(17C)	5782	7057	2225	41

Table 6. Torsion angles [$^{\circ}$] for d1825_a.

C(3)-N(1)-C(1)-C(16)	134.57(14)
Cu(1)-N(1)-C(1)-C(16)	-51.41(17)
C(3)-N(1)-C(1)-C(17)	-100.34(14)
Cu(1)-N(1)-C(1)-C(17)	73.67(15)
C(3)-N(1)-C(1)-C(2)	16.86(15)
Cu(1)-N(1)-C(1)-C(2)	-169.12(10)
C(3)-O(2)-C(2)-C(1)	21.01(16)
N(1)-C(1)-C(2)-O(2)	-22.63(15)
C(16)-C(1)-C(2)-O(2)	-142.86(13)
C(17)-C(1)-C(2)-O(2)	92.75(15)
C(1)-N(1)-C(3)-O(2)	-4.54(17)
Cu(1)-N(1)-C(3)-O(2)	-179.24(9)
C(1)-N(1)-C(3)-C(4)	175.53(15)
Cu(1)-N(1)-C(3)-C(4)	0.8(2)
C(2)-O(2)-C(3)-N(1)	-11.01(17)
C(2)-O(2)-C(3)-C(4)	168.93(13)
N(1)-C(3)-C(4)-C(5)	9.2(3)
O(2)-C(3)-C(4)-C(5)	-170.76(14)
Cu(1)-O(1)-C(5)-C(4)	-13.7(2)
Cu(1)-O(1)-C(5)-C(6)	166.92(10)
C(3)-C(4)-C(5)-O(1)	-2.6(2)
C(3)-C(4)-C(5)-C(6)	176.79(13)
O(1)-C(5)-C(6)-C(7)	-119.23(15)
C(4)-C(5)-C(6)-C(7)	61.31(19)
O(1)-C(5)-C(6)-C(15)	59.56(19)
C(4)-C(5)-C(6)-C(15)	-119.89(16)
C(15)-C(6)-C(7)-C(8)	-1.3(2)
C(5)-C(6)-C(7)-C(8)	177.51(14)
C(6)-C(7)-C(8)-C(9)	0.8(2)
C(7)-C(8)-C(9)-C(10)	0.2(3)
C(8)-C(9)-C(10)-C(11)	179.29(16)
C(8)-C(9)-C(10)-C(15)	-0.6(2)
C(9)-C(10)-C(11)-C(12)	-179.02(17)
C(15)-C(10)-C(11)-C(12)	0.9(3)

C(10)-C(11)-C(12)-C(13)	0.6(3)
C(11)-C(12)-C(13)-C(14)	-0.9(3)
C(12)-C(13)-C(14)-C(15)	-0.3(3)
C(13)-C(14)-C(15)-C(10)	1.8(2)
C(13)-C(14)-C(15)-C(6)	179.45(15)
C(11)-C(10)-C(15)-C(14)	-2.1(2)
C(9)-C(10)-C(15)-C(14)	177.82(14)
C(11)-C(10)-C(15)-C(6)	-179.80(14)
C(9)-C(10)-C(15)-C(6)	0.1(2)
C(7)-C(6)-C(15)-C(14)	-176.78(14)
C(5)-C(6)-C(15)-C(14)	4.4(2)
C(7)-C(6)-C(15)-C(10)	0.8(2)
C(5)-C(6)-C(15)-C(10)	-177.95(13)

Symmetry transformations used to generate equivalent atoms:

#1 -x+1,-y+1,-z+1

F8: Crystal data and structure refinement for d1871_a (**7h**).

Identification code	d1871_a (7h)	
Empirical formula	C ₂₂ H ₃₆ Cu N ₂ O ₄	
Formula weight	456.07	
Temperature	150(2) K	
Wavelength	0.71073 Å	
Crystal system	Triclinic	
Space group	P-1	
Unit cell dimensions	a = 11.4940(9) Å	a = 92.155(3)°.
	b = 13.6089(12) Å	b = 101.732(2)°.
	c = 15.8322(14) Å	g = 101.890(2)°.
Volume	2364.5(3) Å ³	
Z	4	
Density (calculated)	1.281 Mg/m ³	
Absorption coefficient	0.952 mm ⁻¹	
F(000)	972	
Crystal size	0.260 x 0.100 x 0.030 mm ³	
Theta range for data collection	1.534 to 27.568°.	
Index ranges	-14 ≤ h ≤ 14, -17 ≤ k ≤ 17, -20 ≤ l ≤ 20	
Reflections collected	77128	
Independent reflections	10909 [R(int) = 0.1093]	
Completeness to theta = 25.242°	100.0 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.7456 and 0.6549	

Refinement method	Full-matrix least-squares on F^2
Data / restraints / parameters	10909 / 38 / 567
Goodness-of-fit on F^2	0.995
Final R indices [$I > 2\sigma(I)$]	$R_1 = 0.0468$, $wR_2 = 0.0874$
R indices (all data)	$R_1 = 0.1087$, $wR_2 = 0.1078$
Extinction coefficient	n/a
Largest diff. peak and hole	0.496 and -0.696 e. $\text{\AA}^{-3}$

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d1871_a. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	$U(\text{eq})$
Cu(1A)	4074(1)	7392(1)	7488(1)	22(1)
O(1A)	3971(2)	8787(2)	7552(1)	25(1)
O(2A)	4423(2)	8099(2)	5046(1)	38(1)
O(3A)	3163(2)	6025(2)	7439(1)	28(1)
O(4A)	5402(2)	6673(2)	9937(1)	32(1)
N(1A)	4220(2)	7404(2)	6286(2)	26(1)
N(2A)	4888(2)	7356(2)	8681(2)	20(1)
C(1A)	3868(3)	9372(2)	6929(2)	23(1)
C(2A)	3947(3)	9121(2)	6101(2)	27(1)
C(3A)	4180(3)	8192(2)	5853(2)	27(1)
C(6A)	3690(3)	10425(2)	7191(2)	28(1)
C(7A)	4901(3)	11030(3)	7739(3)	51(1)
C(8A)	3273(4)	10988(3)	6405(3)	57(1)
C(9A)	2737(3)	10310(2)	7749(2)	36(1)
C(4C)	4460(12)	7043(6)	4878(5)	45(4)
C(5C)	4555(3)	6627(2)	5760(2)	27(1)
C(10C)	3854(8)	5528(5)	5762(6)	37(2)
C(11C)	5883(6)	6666(7)	6195(7)	39(3)
C(4A)	4977(9)	7227(6)	5041(5)	38(3)
C(5A)	4555(3)	6627(2)	5760(2)	27(1)
C(10A)	3336(5)	5899(5)	5368(5)	38(2)
C(11A)	5533(8)	6161(7)	6252(6)	44(2)
C(12A)	3114(3)	5440(2)	8063(2)	23(1)
C(13A)	3797(3)	5675(2)	8887(2)	25(1)
C(14A)	4673(3)	6584(2)	9129(2)	24(1)
C(15A)	6373(3)	7549(2)	9951(2)	34(1)
C(16A)	5862(3)	8149(2)	9229(2)	23(1)
C(17A)	2253(3)	4394(2)	7810(2)	30(1)
C(18A)	3003(5)	3613(5)	7697(5)	44(1)
C(19A)	1622(7)	4085(5)	8582(4)	44(1)
C(20A)	1258(6)	4371(6)	7019(4)	44(1)

C(18C)	2880(6)	3830(6)	7218(5)	44(1)
C(19C)	1991(8)	3797(6)	8539(4)	44(1)
C(20C)	1100(7)	4593(7)	7223(5)	44(1)
C(21A)	6796(3)	8607(2)	8724(2)	32(1)
C(22A)	5272(3)	8938(2)	9580(2)	32(1)
Cu(1B)	892(1)	2416(1)	2423(1)	21(1)
O(1B)	1260(2)	1244(2)	2968(1)	26(1)
O(2B)	-1069(2)	199(2)	551(1)	29(1)
O(3B)	1561(2)	3603(1)	1904(1)	24(1)
O(4B)	894(2)	4587(2)	4277(1)	33(1)
N(1B)	-73(2)	1618(2)	1377(2)	19(1)
N(2B)	824(2)	3192(2)	3452(2)	21(1)
C(1B)	1009(3)	328(2)	2634(2)	24(1)
C(2B)	300(3)	-7(2)	1828(2)	23(1)
C(3B)	-248(3)	644(2)	1279(2)	22(1)
C(4B)	-1649(3)	984(2)	189(2)	33(1)
C(5B)	-791(3)	1973(2)	606(2)	24(1)
C(6B)	1532(3)	-412(2)	3237(2)	31(1)
C(7B)	807(4)	-555(3)	3946(3)	67(1)
C(8B)	1466(4)	-1429(3)	2771(3)	62(1)
C(9B)	2862(3)	55(3)	3645(2)	42(1)
C(10B)	58(3)	2393(2)	17(2)	34(1)
C(11B)	-1478(3)	2727(3)	872(2)	36(1)
C(12B)	1892(3)	4526(2)	2243(2)	22(1)
C(13B)	1721(3)	4832(2)	3033(2)	27(1)
C(14B)	1150(3)	4167(2)	3561(2)	25(1)
C(15B)	76(3)	3784(3)	4580(2)	35(1)
C(16B)	332(3)	2822(2)	4212(2)	25(1)
C(17B)	2476(3)	5279(2)	1674(2)	31(1)
C(18B)	3507(4)	4897(3)	1404(3)	66(1)
C(19B)	2915(5)	6329(3)	2089(3)	91(2)
C(20B)	1528(4)	5286(4)	847(3)	82(2)
C(21B)	-812(3)	2006(3)	3925(2)	36(1)
C(22B)	1319(3)	2463(3)	4842(2)	35(1)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for d1871_a.

Cu(1A)-O(1A)	1.927(2)
Cu(1A)-O(3A)	1.928(2)
Cu(1A)-N(2A)	1.937(2)
Cu(1A)-N(1A)	1.945(2)
O(1A)-C(1A)	1.293(3)
O(2A)-C(3A)	1.370(3)
O(2A)-C(4A)	1.458(6)
O(2A)-C(4C)	1.462(7)
O(3A)-C(12A)	1.294(4)
O(4A)-C(14A)	1.365(3)
O(4A)-C(15A)	1.452(4)
N(1A)-C(3A)	1.299(4)
N(1A)-C(5A)	1.481(4)
N(1A)-C(5C)	1.481(4)
N(2A)-C(14A)	1.300(4)
N(2A)-C(16A)	1.485(3)
C(1A)-C(2A)	1.367(4)
C(1A)-C(6A)	1.542(4)
C(2A)-C(3A)	1.403(4)
C(2A)-H(2AA)	0.9500
C(6A)-C(9A)	1.530(4)
C(6A)-C(8A)	1.530(5)
C(6A)-C(7A)	1.533(4)
C(7A)-H(7AA)	0.9800
C(7A)-H(7AB)	0.9800
C(7A)-H(7AC)	0.9800
C(8A)-H(8AA)	0.9800
C(8A)-H(8AB)	0.9800
C(8A)-H(8AC)	0.9800
C(9A)-H(9AA)	0.9800
C(9A)-H(9AB)	0.9800
C(9A)-H(9AC)	0.9800
C(4C)-C(5C)	1.519(7)
C(4C)-H(4AA)	0.9900

C(4C)-H(4AB)	0.9900
C(5C)-C(11C)	1.531(7)
C(5C)-C(10C)	1.545(6)
C(10C)-H(10A)	0.9800
C(10C)-H(10B)	0.9800
C(10C)-H(10C)	0.9800
C(11C)-H(11A)	0.9800
C(11C)-H(11B)	0.9800
C(11C)-H(11C)	0.9800
C(4A)-C(5A)	1.525(7)
C(4A)-H(4CA)	0.9900
C(4A)-H(4CB)	0.9900
C(5A)-C(11A)	1.499(6)
C(5A)-C(10A)	1.535(5)
C(10A)-H(10D)	0.9800
C(10A)-H(10E)	0.9800
C(10A)-H(10F)	0.9800
C(11A)-H(11D)	0.9800
C(11A)-H(11E)	0.9800
C(11A)-H(11F)	0.9800
C(12A)-C(13A)	1.367(4)
C(12A)-C(17A)	1.544(4)
C(13A)-C(14A)	1.409(4)
C(13A)-H(13A)	0.9500
C(15A)-C(16A)	1.521(4)
C(15A)-H(15A)	0.9900
C(15A)-H(15B)	0.9900
C(16A)-C(21A)	1.514(4)
C(16A)-C(22A)	1.527(4)
C(17A)-C(19C)	1.482(5)
C(17A)-C(20A)	1.509(5)
C(17A)-C(18A)	1.526(5)
C(17A)-C(20C)	1.536(5)
C(17A)-C(18C)	1.559(5)
C(17A)-C(19A)	1.571(5)
C(18A)-H(18A)	0.9800

C(18A)-H(18B)	0.9800
C(18A)-H(18C)	0.9800
C(19A)-H(19A)	0.9800
C(19A)-H(19B)	0.9800
C(19A)-H(19C)	0.9800
C(20A)-H(20A)	0.9800
C(20A)-H(20B)	0.9800
C(20A)-H(20C)	0.9800
C(18C)-H(18G)	0.9800
C(18C)-H(18H)	0.9800
C(18C)-H(18I)	0.9800
C(19C)-H(19G)	0.9800
C(19C)-H(19H)	0.9800
C(19C)-H(19I)	0.9800
C(20C)-H(20G)	0.9800
C(20C)-H(20H)	0.9800
C(20C)-H(20I)	0.9800
C(21A)-H(21A)	0.9800
C(21A)-H(21B)	0.9800
C(21A)-H(21C)	0.9800
C(22A)-H(22A)	0.9800
C(22A)-H(22B)	0.9800
C(22A)-H(22C)	0.9800
Cu(1B)-O(1B)	1.923(2)
Cu(1B)-O(3B)	1.928(2)
Cu(1B)-N(2B)	1.933(2)
Cu(1B)-N(1B)	1.942(2)
O(1B)-C(1B)	1.288(3)
O(2B)-C(3B)	1.363(3)
O(2B)-C(4B)	1.448(4)
O(3B)-C(12B)	1.297(3)
O(4B)-C(14B)	1.362(3)
O(4B)-C(15B)	1.451(4)
N(1B)-C(3B)	1.297(3)
N(1B)-C(5B)	1.484(4)
N(2B)-C(14B)	1.296(4)

N(2B)-C(16B)	1.492(4)
C(1B)-C(2B)	1.373(4)
C(1B)-C(6B)	1.540(4)
C(2B)-C(3B)	1.414(4)
C(2B)-H(2BA)	0.9500
C(4B)-C(5B)	1.531(4)
C(4B)-H(4BA)	0.9900
C(4B)-H(4BB)	0.9900
C(5B)-C(11B)	1.513(4)
C(5B)-C(10B)	1.526(4)
C(6B)-C(7B)	1.524(5)
C(6B)-C(8B)	1.525(5)
C(6B)-C(9B)	1.525(4)
C(7B)-H(7BA)	0.9800
C(7B)-H(7BB)	0.9800
C(7B)-H(7BC)	0.9800
C(8B)-H(8BA)	0.9800
C(8B)-H(8BB)	0.9800
C(8B)-H(8BC)	0.9800
C(9B)-H(9BA)	0.9800
C(9B)-H(9BB)	0.9800
C(9B)-H(9BC)	0.9800
C(10B)-H(10G)	0.9800
C(10B)-H(10H)	0.9800
C(10B)-H(10I)	0.9800
C(11B)-H(11G)	0.9800
C(11B)-H(11H)	0.9800
C(11B)-H(11I)	0.9800
C(12B)-C(13B)	1.368(4)
C(12B)-C(17B)	1.531(4)
C(13B)-C(14B)	1.409(4)
C(13B)-H(13B)	0.9500
C(15B)-C(16B)	1.520(4)
C(15B)-H(15C)	0.9900
C(15B)-H(15D)	0.9900
C(16B)-C(21B)	1.513(4)

C(16B)-C(22B)	1.525(4)
C(17B)-C(19B)	1.492(5)
C(17B)-C(18B)	1.517(5)
C(17B)-C(20B)	1.526(5)
C(18B)-H(18D)	0.9800
C(18B)-H(18E)	0.9800
C(18B)-H(18F)	0.9800
C(19B)-H(19D)	0.9800
C(19B)-H(19E)	0.9800
C(19B)-H(19F)	0.9800
C(20B)-H(20D)	0.9800
C(20B)-H(20E)	0.9800
C(20B)-H(20F)	0.9800
C(21B)-H(21D)	0.9800
C(21B)-H(21E)	0.9800
C(21B)-H(21F)	0.9800
C(22B)-H(22D)	0.9800
C(22B)-H(22E)	0.9800
C(22B)-H(22F)	0.9800

O(1A)-Cu(1A)-O(3A)	144.37(9)
O(1A)-Cu(1A)-N(2A)	97.49(9)
O(3A)-Cu(1A)-N(2A)	92.15(9)
O(1A)-Cu(1A)-N(1A)	92.45(10)
O(3A)-Cu(1A)-N(1A)	98.05(10)
N(2A)-Cu(1A)-N(1A)	146.69(10)
C(1A)-O(1A)-Cu(1A)	128.12(19)
C(3A)-O(2A)-C(4A)	105.4(3)
C(3A)-O(2A)-C(4C)	106.6(4)
C(12A)-O(3A)-Cu(1A)	128.24(19)
C(14A)-O(4A)-C(15A)	105.3(2)
C(3A)-N(1A)-C(5A)	109.0(2)
C(3A)-N(1A)-C(5C)	109.0(2)
C(3A)-N(1A)-Cu(1A)	122.6(2)
C(5A)-N(1A)-Cu(1A)	127.9(2)
C(5C)-N(1A)-Cu(1A)	127.9(2)

C(14A)-N(2A)-C(16A)	108.4(2)
C(14A)-N(2A)-Cu(1A)	123.7(2)
C(16A)-N(2A)-Cu(1A)	127.88(19)
O(1A)-C(1A)-C(2A)	124.5(3)
O(1A)-C(1A)-C(6A)	114.8(3)
C(2A)-C(1A)-C(6A)	120.7(3)
C(1A)-C(2A)-C(3A)	122.3(3)
C(1A)-C(2A)-H(2AA)	118.8
C(3A)-C(2A)-H(2AA)	118.8
N(1A)-C(3A)-O(2A)	114.6(3)
N(1A)-C(3A)-C(2A)	129.3(3)
O(2A)-C(3A)-C(2A)	116.2(3)
C(9A)-C(6A)-C(8A)	109.0(3)
C(9A)-C(6A)-C(7A)	108.3(3)
C(8A)-C(6A)-C(7A)	110.1(3)
C(9A)-C(6A)-C(1A)	108.9(3)
C(8A)-C(6A)-C(1A)	112.4(3)
C(7A)-C(6A)-C(1A)	108.2(3)
C(6A)-C(7A)-H(7AA)	109.5
C(6A)-C(7A)-H(7AB)	109.5
H(7AA)-C(7A)-H(7AB)	109.5
C(6A)-C(7A)-H(7AC)	109.5
H(7AA)-C(7A)-H(7AC)	109.5
H(7AB)-C(7A)-H(7AC)	109.5
C(6A)-C(8A)-H(8AA)	109.5
C(6A)-C(8A)-H(8AB)	109.5
H(8AA)-C(8A)-H(8AB)	109.5
C(6A)-C(8A)-H(8AC)	109.5
H(8AA)-C(8A)-H(8AC)	109.5
H(8AB)-C(8A)-H(8AC)	109.5
C(6A)-C(9A)-H(9AA)	109.5
C(6A)-C(9A)-H(9AB)	109.5
H(9AA)-C(9A)-H(9AB)	109.5
C(6A)-C(9A)-H(9AC)	109.5
H(9AA)-C(9A)-H(9AC)	109.5
H(9AB)-C(9A)-H(9AC)	109.5

O(2A)-C(4C)-C(5C)	104.3(5)
O(2A)-C(4C)-H(4AA)	110.9
C(5C)-C(4C)-H(4AA)	110.9
O(2A)-C(4C)-H(4AB)	110.9
C(5C)-C(4C)-H(4AB)	110.9
H(4AA)-C(4C)-H(4AB)	108.9
N(1A)-C(5C)-C(4C)	103.0(4)
N(1A)-C(5C)-C(11C)	103.3(5)
C(4C)-C(5C)-C(11C)	111.1(7)
N(1A)-C(5C)-C(10C)	116.7(4)
C(4C)-C(5C)-C(10C)	115.5(5)
C(11C)-C(5C)-C(10C)	106.6(5)
C(5C)-C(10C)-H(10A)	109.5
C(5C)-C(10C)-H(10B)	109.5
H(10A)-C(10C)-H(10B)	109.5
C(5C)-C(10C)-H(10C)	109.5
H(10A)-C(10C)-H(10C)	109.5
H(10B)-C(10C)-H(10C)	109.5
C(5C)-C(11C)-H(11A)	109.5
C(5C)-C(11C)-H(11B)	109.5
H(11A)-C(11C)-H(11B)	109.5
C(5C)-C(11C)-H(11C)	109.5
H(11A)-C(11C)-H(11C)	109.5
H(11B)-C(11C)-H(11C)	109.5
O(2A)-C(4A)-C(5A)	104.2(4)
O(2A)-C(4A)-H(4CA)	110.9
C(5A)-C(4A)-H(4CA)	110.9
O(2A)-C(4A)-H(4CB)	110.9
C(5A)-C(4A)-H(4CB)	110.9
H(4CA)-C(4A)-H(4CB)	108.9
N(1A)-C(5A)-C(11A)	113.6(4)
N(1A)-C(5A)-C(4A)	101.8(3)
C(11A)-C(5A)-C(4A)	111.9(5)
N(1A)-C(5A)-C(10A)	104.1(3)
C(11A)-C(5A)-C(10A)	115.9(5)
C(4A)-C(5A)-C(10A)	108.3(5)

C(5A)-C(10A)-H(10D)	109.5
C(5A)-C(10A)-H(10E)	109.5
H(10D)-C(10A)-H(10E)	109.5
C(5A)-C(10A)-H(10F)	109.5
H(10D)-C(10A)-H(10F)	109.5
H(10E)-C(10A)-H(10F)	109.5
C(5A)-C(11A)-H(11D)	109.5
C(5A)-C(11A)-H(11E)	109.5
H(11D)-C(11A)-H(11E)	109.5
C(5A)-C(11A)-H(11F)	109.5
H(11D)-C(11A)-H(11F)	109.5
H(11E)-C(11A)-H(11F)	109.5
O(3A)-C(12A)-C(13A)	124.7(3)
O(3A)-C(12A)-C(17A)	115.1(3)
C(13A)-C(12A)-C(17A)	120.1(3)
C(12A)-C(13A)-C(14A)	122.2(3)
C(12A)-C(13A)-H(13A)	118.9
C(14A)-C(13A)-H(13A)	118.9
N(2A)-C(14A)-O(4A)	115.1(3)
N(2A)-C(14A)-C(13A)	128.5(3)
O(4A)-C(14A)-C(13A)	116.4(3)
O(4A)-C(15A)-C(16A)	104.7(2)
O(4A)-C(15A)-H(15A)	110.8
C(16A)-C(15A)-H(15A)	110.8
O(4A)-C(15A)-H(15B)	110.8
C(16A)-C(15A)-H(15B)	110.8
H(15A)-C(15A)-H(15B)	108.9
N(2A)-C(16A)-C(21A)	110.9(2)
N(2A)-C(16A)-C(15A)	101.3(2)
C(21A)-C(16A)-C(15A)	112.8(3)
N(2A)-C(16A)-C(22A)	108.5(2)
C(21A)-C(16A)-C(22A)	112.0(3)
C(15A)-C(16A)-C(22A)	110.7(3)
C(20A)-C(17A)-C(18A)	111.6(4)
C(19C)-C(17A)-C(20C)	112.8(4)
C(19C)-C(17A)-C(12A)	115.8(4)

C(20A)-C(17A)-C(12A)	113.3(4)
C(18A)-C(17A)-C(12A)	109.2(3)
C(20C)-C(17A)-C(12A)	105.2(4)
C(19C)-C(17A)-C(18C)	110.9(4)
C(20C)-C(17A)-C(18C)	106.4(4)
C(12A)-C(17A)-C(18C)	104.9(4)
C(20A)-C(17A)-C(19A)	107.4(4)
C(18A)-C(17A)-C(19A)	106.4(3)
C(12A)-C(17A)-C(19A)	108.6(3)
C(17A)-C(18A)-H(18A)	109.5
C(17A)-C(18A)-H(18B)	109.5
H(18A)-C(18A)-H(18B)	109.5
C(17A)-C(18A)-H(18C)	109.5
H(18A)-C(18A)-H(18C)	109.5
H(18B)-C(18A)-H(18C)	109.5
C(17A)-C(19A)-H(19A)	109.5
C(17A)-C(19A)-H(19B)	109.5
H(19A)-C(19A)-H(19B)	109.5
C(17A)-C(19A)-H(19C)	109.5
H(19A)-C(19A)-H(19C)	109.5
H(19B)-C(19A)-H(19C)	109.5
C(17A)-C(20A)-H(20A)	109.5
C(17A)-C(20A)-H(20B)	109.5
H(20A)-C(20A)-H(20B)	109.5
C(17A)-C(20A)-H(20C)	109.5
H(20A)-C(20A)-H(20C)	109.5
H(20B)-C(20A)-H(20C)	109.5
C(17A)-C(18C)-H(18G)	109.5
C(17A)-C(18C)-H(18H)	109.5
H(18G)-C(18C)-H(18H)	109.5
C(17A)-C(18C)-H(18I)	109.5
H(18G)-C(18C)-H(18I)	109.5
H(18H)-C(18C)-H(18I)	109.5
C(17A)-C(19C)-H(19G)	109.5
C(17A)-C(19C)-H(19H)	109.5
H(19G)-C(19C)-H(19H)	109.5

C(17A)-C(19C)-H(19I)	109.5
H(19G)-C(19C)-H(19I)	109.5
H(19H)-C(19C)-H(19I)	109.5
C(17A)-C(20C)-H(20G)	109.5
C(17A)-C(20C)-H(20H)	109.5
H(20G)-C(20C)-H(20H)	109.5
C(17A)-C(20C)-H(20I)	109.5
H(20G)-C(20C)-H(20I)	109.5
H(20H)-C(20C)-H(20I)	109.5
C(16A)-C(21A)-H(21A)	109.5
C(16A)-C(21A)-H(21B)	109.5
H(21A)-C(21A)-H(21B)	109.5
C(16A)-C(21A)-H(21C)	109.5
H(21A)-C(21A)-H(21C)	109.5
H(21B)-C(21A)-H(21C)	109.5
C(16A)-C(22A)-H(22A)	109.5
C(16A)-C(22A)-H(22B)	109.5
H(22A)-C(22A)-H(22B)	109.5
C(16A)-C(22A)-H(22C)	109.5
H(22A)-C(22A)-H(22C)	109.5
H(22B)-C(22A)-H(22C)	109.5
O(1B)-Cu(1B)-O(3B)	144.32(9)
O(1B)-Cu(1B)-N(2B)	97.60(10)
O(3B)-Cu(1B)-N(2B)	92.58(9)
O(1B)-Cu(1B)-N(1B)	92.50(9)
O(3B)-Cu(1B)-N(1B)	98.69(9)
N(2B)-Cu(1B)-N(1B)	144.66(10)
C(1B)-O(1B)-Cu(1B)	128.2(2)
C(3B)-O(2B)-C(4B)	105.7(2)
C(12B)-O(3B)-Cu(1B)	128.1(2)
C(14B)-O(4B)-C(15B)	105.3(2)
C(3B)-N(1B)-C(5B)	109.0(2)
C(3B)-N(1B)-Cu(1B)	123.2(2)
C(5B)-N(1B)-Cu(1B)	127.69(18)
C(14B)-N(2B)-C(16B)	108.3(2)
C(14B)-N(2B)-Cu(1B)	123.5(2)

C(16B)-N(2B)-Cu(1B)	128.08(19)
O(1B)-C(1B)-C(2B)	124.8(3)
O(1B)-C(1B)-C(6B)	114.4(3)
C(2B)-C(1B)-C(6B)	120.7(3)
C(1B)-C(2B)-C(3B)	122.0(3)
C(1B)-C(2B)-H(2BA)	119.0
C(3B)-C(2B)-H(2BA)	119.0
N(1B)-C(3B)-O(2B)	115.1(3)
N(1B)-C(3B)-C(2B)	128.5(3)
O(2B)-C(3B)-C(2B)	116.4(3)
O(2B)-C(4B)-C(5B)	105.1(2)
O(2B)-C(4B)-H(4BA)	110.7
C(5B)-C(4B)-H(4BA)	110.7
O(2B)-C(4B)-H(4BB)	110.7
C(5B)-C(4B)-H(4BB)	110.7
H(4BA)-C(4B)-H(4BB)	108.8
N(1B)-C(5B)-C(11B)	110.9(2)
N(1B)-C(5B)-C(10B)	109.0(2)
C(11B)-C(5B)-C(10B)	112.9(3)
N(1B)-C(5B)-C(4B)	101.0(2)
C(11B)-C(5B)-C(4B)	111.9(3)
C(10B)-C(5B)-C(4B)	110.5(3)
C(7B)-C(6B)-C(8B)	110.0(3)
C(7B)-C(6B)-C(9B)	109.2(3)
C(8B)-C(6B)-C(9B)	108.4(3)
C(7B)-C(6B)-C(1B)	107.2(3)
C(8B)-C(6B)-C(1B)	113.2(3)
C(9B)-C(6B)-C(1B)	108.8(3)
C(6B)-C(7B)-H(7BA)	109.5
C(6B)-C(7B)-H(7BB)	109.5
H(7BA)-C(7B)-H(7BB)	109.5
C(6B)-C(7B)-H(7BC)	109.5
H(7BA)-C(7B)-H(7BC)	109.5
H(7BB)-C(7B)-H(7BC)	109.5
C(6B)-C(8B)-H(8BA)	109.5
C(6B)-C(8B)-H(8BB)	109.5

H(8BA)-C(8B)-H(8BB)	109.5
C(6B)-C(8B)-H(8BC)	109.5
H(8BA)-C(8B)-H(8BC)	109.5
H(8BB)-C(8B)-H(8BC)	109.5
C(6B)-C(9B)-H(9BA)	109.5
C(6B)-C(9B)-H(9BB)	109.5
H(9BA)-C(9B)-H(9BB)	109.5
C(6B)-C(9B)-H(9BC)	109.5
H(9BA)-C(9B)-H(9BC)	109.5
H(9BB)-C(9B)-H(9BC)	109.5
C(5B)-C(10B)-H(10G)	109.5
C(5B)-C(10B)-H(10H)	109.5
H(10G)-C(10B)-H(10H)	109.5
C(5B)-C(10B)-H(10I)	109.5
H(10G)-C(10B)-H(10I)	109.5
H(10H)-C(10B)-H(10I)	109.5
C(5B)-C(11B)-H(11G)	109.5
C(5B)-C(11B)-H(11H)	109.5
H(11G)-C(11B)-H(11H)	109.5
C(5B)-C(11B)-H(11I)	109.5
H(11G)-C(11B)-H(11I)	109.5
H(11H)-C(11B)-H(11I)	109.5
O(3B)-C(12B)-C(13B)	124.1(3)
O(3B)-C(12B)-C(17B)	114.6(3)
C(13B)-C(12B)-C(17B)	121.4(3)
C(12B)-C(13B)-C(14B)	123.0(3)
C(12B)-C(13B)-H(13B)	118.5
C(14B)-C(13B)-H(13B)	118.5
N(2B)-C(14B)-O(4B)	114.9(3)
N(2B)-C(14B)-C(13B)	128.2(3)
O(4B)-C(14B)-C(13B)	116.9(3)
O(4B)-C(15B)-C(16B)	104.3(2)
O(4B)-C(15B)-H(15C)	110.9
C(16B)-C(15B)-H(15C)	110.9
O(4B)-C(15B)-H(15D)	110.9
C(16B)-C(15B)-H(15D)	110.9

H(15C)-C(15B)-H(15D)	108.9
N(2B)-C(16B)-C(21B)	110.8(2)
N(2B)-C(16B)-C(15B)	100.7(2)
C(21B)-C(16B)-C(15B)	112.2(3)
N(2B)-C(16B)-C(22B)	109.1(2)
C(21B)-C(16B)-C(22B)	112.1(3)
C(15B)-C(16B)-C(22B)	111.3(3)
C(19B)-C(17B)-C(18B)	110.7(3)
C(19B)-C(17B)-C(20B)	108.6(4)
C(18B)-C(17B)-C(20B)	106.6(4)
C(19B)-C(17B)-C(12B)	114.1(3)
C(18B)-C(17B)-C(12B)	108.7(3)
C(20B)-C(17B)-C(12B)	107.8(3)
C(17B)-C(18B)-H(18D)	109.5
C(17B)-C(18B)-H(18E)	109.5
H(18D)-C(18B)-H(18E)	109.5
C(17B)-C(18B)-H(18F)	109.5
H(18D)-C(18B)-H(18F)	109.5
H(18E)-C(18B)-H(18F)	109.5
C(17B)-C(19B)-H(19D)	109.5
C(17B)-C(19B)-H(19E)	109.5
H(19D)-C(19B)-H(19E)	109.5
C(17B)-C(19B)-H(19F)	109.5
H(19D)-C(19B)-H(19F)	109.5
H(19E)-C(19B)-H(19F)	109.5
C(17B)-C(20B)-H(20D)	109.5
C(17B)-C(20B)-H(20E)	109.5
H(20D)-C(20B)-H(20E)	109.5
C(17B)-C(20B)-H(20F)	109.5
H(20D)-C(20B)-H(20F)	109.5
H(20E)-C(20B)-H(20F)	109.5
C(16B)-C(21B)-H(21D)	109.5
C(16B)-C(21B)-H(21E)	109.5
H(21D)-C(21B)-H(21E)	109.5
C(16B)-C(21B)-H(21F)	109.5
H(21D)-C(21B)-H(21F)	109.5

H(21E)-C(21B)-H(21F)	109.5
C(16B)-C(22B)-H(22D)	109.5
C(16B)-C(22B)-H(22E)	109.5
H(22D)-C(22B)-H(22E)	109.5
C(16B)-C(22B)-H(22F)	109.5
H(22D)-C(22B)-H(22F)	109.5
H(22E)-C(22B)-H(22F)	109.5

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d1871_a. The anisotropic displacement factor exponent takes the form: $-2p^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Cu(1A)	26(1)	21(1)	18(1)	0(1)	3(1)	8(1)
O(1A)	31(1)	23(1)	21(1)	2(1)	6(1)	8(1)
O(2A)	55(2)	47(2)	21(1)	9(1)	16(1)	27(1)
O(3A)	37(1)	22(1)	21(1)	1(1)	-4(1)	6(1)
O(4A)	33(1)	34(1)	21(1)	5(1)	-5(1)	-3(1)
N(1A)	28(2)	29(2)	23(1)	-2(1)	4(1)	15(1)
N(2A)	18(1)	21(1)	18(1)	-2(1)	2(1)	3(1)
C(1A)	17(2)	24(2)	28(2)	3(1)	4(1)	6(1)
C(2A)	30(2)	31(2)	23(2)	8(1)	5(1)	14(1)
C(3A)	26(2)	39(2)	20(2)	2(1)	6(1)	14(2)
C(6A)	32(2)	21(2)	35(2)	2(1)	12(2)	9(1)
C(7A)	38(2)	26(2)	83(3)	-14(2)	14(2)	-1(2)
C(8A)	99(3)	37(2)	53(3)	19(2)	29(2)	42(2)
C(9A)	39(2)	32(2)	41(2)	-4(2)	17(2)	13(2)
C(4C)	66(10)	61(7)	23(5)	3(5)	4(5)	53(6)
C(5C)	29(2)	27(2)	26(2)	-4(1)	10(1)	10(1)
C(10C)	40(5)	29(5)	43(5)	-9(4)	16(4)	4(4)
C(11C)	56(6)	39(6)	42(6)	11(5)	20(5)	42(5)
C(4A)	47(7)	63(6)	15(4)	-1(4)	2(4)	40(4)
C(5A)	29(2)	27(2)	26(2)	-4(1)	10(1)	10(1)
C(10A)	37(4)	37(4)	36(4)	-17(3)	11(3)	2(3)
C(11A)	72(6)	41(6)	35(4)	13(5)	14(4)	45(5)
C(12A)	21(2)	20(2)	30(2)	0(1)	5(1)	9(1)
C(13A)	25(2)	24(2)	24(2)	3(1)	4(1)	3(1)
C(14A)	24(2)	27(2)	19(2)	-1(1)	2(1)	8(1)
C(15A)	32(2)	29(2)	33(2)	-1(2)	-4(2)	0(2)
C(16A)	23(2)	24(2)	19(2)	-5(1)	1(1)	3(1)
C(17A)	28(2)	21(2)	37(2)	-2(1)	-1(2)	6(1)
C(18A)	48(2)	23(2)	52(2)	2(2)	-4(2)	0(1)
C(19A)	48(2)	23(2)	52(2)	2(2)	-4(2)	0(1)
C(20A)	48(2)	23(2)	52(2)	2(2)	-4(2)	0(1)

C(18C)	48(2)	23(2)	52(2)	2(2)	-4(2)	0(1)
C(19C)	48(2)	23(2)	52(2)	2(2)	-4(2)	0(1)
C(20C)	48(2)	23(2)	52(2)	2(2)	-4(2)	0(1)
C(21A)	26(2)	31(2)	37(2)	-3(2)	8(2)	1(1)
C(22A)	32(2)	31(2)	30(2)	-12(2)	4(2)	3(2)
Cu(1B)	25(1)	19(1)	19(1)	-1(1)	4(1)	4(1)
O(1B)	32(1)	20(1)	24(1)	0(1)	3(1)	6(1)
O(2B)	32(1)	22(1)	27(1)	-4(1)	-1(1)	1(1)
O(3B)	30(1)	16(1)	27(1)	-2(1)	8(1)	4(1)
O(4B)	36(1)	34(1)	31(1)	-11(1)	12(1)	8(1)
N(1B)	22(1)	17(1)	18(1)	2(1)	4(1)	4(1)
N(2B)	19(1)	24(1)	21(1)	-1(1)	5(1)	4(1)
C(1B)	23(2)	22(2)	31(2)	7(1)	12(1)	5(1)
C(2B)	28(2)	15(2)	28(2)	0(1)	10(1)	6(1)
C(3B)	18(2)	22(2)	24(2)	-4(1)	8(1)	0(1)
C(4B)	28(2)	30(2)	33(2)	0(2)	-4(2)	-1(1)
C(5B)	23(2)	26(2)	22(2)	3(1)	-2(1)	4(1)
C(6B)	31(2)	25(2)	39(2)	11(2)	6(2)	9(1)
C(7B)	67(3)	83(3)	69(3)	57(3)	31(2)	30(2)
C(8B)	78(3)	26(2)	73(3)	5(2)	-15(2)	22(2)
C(9B)	42(2)	44(2)	42(2)	14(2)	2(2)	18(2)
C(10B)	36(2)	32(2)	27(2)	6(2)	2(2)	-1(2)
C(11B)	32(2)	38(2)	40(2)	4(2)	2(2)	15(2)
C(12B)	16(2)	20(2)	31(2)	-1(1)	4(1)	6(1)
C(13B)	25(2)	18(2)	34(2)	-7(1)	4(1)	3(1)
C(14B)	22(2)	27(2)	26(2)	-7(1)	2(1)	9(1)
C(15B)	35(2)	45(2)	29(2)	-5(2)	12(2)	12(2)
C(16B)	24(2)	34(2)	20(2)	1(1)	6(1)	8(1)
C(17B)	32(2)	22(2)	40(2)	5(2)	12(2)	5(1)
C(18B)	67(3)	63(3)	94(4)	42(3)	54(3)	28(2)
C(19B)	150(5)	32(2)	86(4)	-12(2)	67(4)	-34(3)
C(20B)	84(4)	67(3)	80(4)	50(3)	-5(3)	-3(3)
C(21B)	28(2)	48(2)	29(2)	5(2)	6(2)	0(2)
C(22B)	32(2)	49(2)	24(2)	6(2)	5(2)	11(2)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for d1871_a.

	x	y	z	U(eq)
H(2AA)	3841	9592	5680	32
H(7AA)	5508	11159	7379	76
H(7AB)	5186	10645	8220	76
H(7AC)	4784	11673	7969	76
H(8AA)	3908	11111	6068	85
H(8AB)	3124	11633	6602	85
H(8AC)	2521	10580	6043	85
H(9AA)	1957	9918	7411	53
H(9AB)	2639	10977	7935	53
H(9AC)	3004	9960	8259	53
H(4AA)	5174	6987	4634	54
H(4AB)	3711	6679	4468	54
H(10A)	2981	5484	5548	56
H(10B)	3996	5320	6353	56
H(10C)	4140	5085	5387	56
H(11A)	6366	7350	6201	59
H(11B)	6205	6194	5871	59
H(11C)	5928	6477	6790	59
H(4CA)	5876	7438	5164	46
H(4CB)	4693	6825	4475	46
H(10D)	2781	6269	5029	56
H(10E)	2975	5599	5834	56
H(10F)	3473	5366	4991	56
H(11D)	6257	6690	6495	65
H(11E)	5740	5681	5862	65
H(11F)	5241	5808	6723	65
H(13A)	3676	5210	9311	30
H(15A)	7111	7345	9843	41
H(15B)	6582	7951	10517	41
H(18A)	3640	3648	8220	66

H(18B)	2472	2938	7598	66
H(18C)	3380	3752	7199	66
H(19A)	2246	4093	9107	66
H(19B)	1122	4564	8677	66
H(19C)	1105	3406	8442	66
H(20A)	795	4879	7113	66
H(20B)	1621	4516	6516	66
H(20C)	712	3702	6915	66
H(18G)	3044	4239	6742	66
H(18H)	3648	3718	7559	66
H(18I)	2341	3180	6983	66
H(19G)	2755	3692	8891	66
H(19H)	1588	4161	8897	66
H(19I)	1456	3143	8312	66
H(20G)	1324	4988	6754	66
H(20H)	553	3949	6979	66
H(20I)	686	4967	7564	66
H(21A)	7091	8069	8460	48
H(21B)	7481	9063	9116	48
H(21C)	6422	8984	8269	48
H(22A)	4680	8614	9906	49
H(22B)	4855	9254	9096	49
H(22C)	5900	9453	9961	49
H(2BA)	173	-697	1631	28
H(4BA)	-2460	922	332	40
H(4BB)	-1750	946	-448	40
H(7BA)	-57	-815	3684	101
H(7BB)	909	94	4274	101
H(7BC)	1104	-1035	4337	101
H(8BA)	612	-1769	2552	93
H(8BB)	1869	-1846	3175	93
H(8BC)	1876	-1327	2285	93
H(9BA)	2911	664	4013	63
H(9BB)	3310	231	3188	63
H(9BC)	3220	-430	3996	63
H(10G)	686	2961	330	51

H(10H)	-412	2621	-497	51
H(10I)	446	1864	-159	51
H(11G)	-898	3342	1148	55
H(11H)	-1962	2437	1281	55
H(11I)	-2019	2889	359	55
H(13B)	2001	5523	3235	32
H(15C)	-784	3821	4362	42
H(15D)	248	3823	5220	42
H(18D)	4148	4892	1917	100
H(18E)	3842	5341	998	100
H(18F)	3199	4212	1124	100
H(19D)	3563	6343	2601	137
H(19E)	2238	6560	2262	137
H(19F)	3231	6774	1676	137
H(20D)	860	5561	987	124
H(20E)	1209	4596	579	124
H(20F)	1906	5704	443	124
H(21D)	-1385	2242	3474	54
H(21E)	-1188	1846	4420	54
H(21F)	-606	1401	3695	54
H(22D)	1586	1932	4546	52
H(22E)	992	2196	5333	52
H(22F)	2013	3029	5052	52

Table 6. Torsion angles [°] for d1871_a.

Cu(1A)-O(1A)-C(1A)-C(2A)	-7.6(4)
Cu(1A)-O(1A)-C(1A)-C(6A)	174.64(18)
O(1A)-C(1A)-C(2A)-C(3A)	-1.5(5)
C(6A)-C(1A)-C(2A)-C(3A)	176.2(3)
C(5A)-N(1A)-C(3A)-O(2A)	1.3(4)
C(5C)-N(1A)-C(3A)-O(2A)	1.3(4)
Cu(1A)-N(1A)-C(3A)-O(2A)	173.53(19)
C(5A)-N(1A)-C(3A)-C(2A)	-177.4(3)
C(5C)-N(1A)-C(3A)-C(2A)	-177.4(3)
Cu(1A)-N(1A)-C(3A)-C(2A)	-5.2(5)
C(4A)-O(2A)-C(3A)-N(1A)	-15.3(5)
C(4C)-O(2A)-C(3A)-N(1A)	9.1(6)
C(4A)-O(2A)-C(3A)-C(2A)	163.5(5)
C(4C)-O(2A)-C(3A)-C(2A)	-172.0(6)
C(1A)-C(2A)-C(3A)-N(1A)	8.5(5)
C(1A)-C(2A)-C(3A)-O(2A)	-170.2(3)
O(1A)-C(1A)-C(6A)-C(9A)	-45.5(3)
C(2A)-C(1A)-C(6A)-C(9A)	136.6(3)
O(1A)-C(1A)-C(6A)-C(8A)	-166.3(3)
C(2A)-C(1A)-C(6A)-C(8A)	15.8(4)
O(1A)-C(1A)-C(6A)-C(7A)	72.0(3)
C(2A)-C(1A)-C(6A)-C(7A)	-105.9(3)
C(3A)-O(2A)-C(4C)-C(5C)	-14.9(9)
C(3A)-N(1A)-C(5C)-C(4C)	-10.5(6)
Cu(1A)-N(1A)-C(5C)-C(4C)	177.7(5)
C(3A)-N(1A)-C(5C)-C(11C)	105.2(5)
Cu(1A)-N(1A)-C(5C)-C(11C)	-66.5(5)
C(3A)-N(1A)-C(5C)-C(10C)	-138.2(5)
Cu(1A)-N(1A)-C(5C)-C(10C)	50.1(5)
O(2A)-C(4C)-C(5C)-N(1A)	15.1(8)
O(2A)-C(4C)-C(5C)-C(11C)	-94.9(8)
O(2A)-C(4C)-C(5C)-C(10C)	143.5(7)
C(3A)-O(2A)-C(4A)-C(5A)	21.9(7)
C(3A)-N(1A)-C(5A)-C(11A)	133.0(5)

Cu(1A)-N(1A)-C(5A)-C(11A)	-38.7(5)
C(3A)-N(1A)-C(5A)-C(4A)	12.5(5)
Cu(1A)-N(1A)-C(5A)-C(4A)	-159.2(4)
C(3A)-N(1A)-C(5A)-C(10A)	-100.0(4)
Cu(1A)-N(1A)-C(5A)-C(10A)	88.3(4)
O(2A)-C(4A)-C(5A)-N(1A)	-20.6(6)
O(2A)-C(4A)-C(5A)-C(11A)	-142.3(6)
O(2A)-C(4A)-C(5A)-C(10A)	88.7(7)
Cu(1A)-O(3A)-C(12A)-C(13A)	-5.1(4)
Cu(1A)-O(3A)-C(12A)-C(17A)	177.86(18)
O(3A)-C(12A)-C(13A)-C(14A)	-2.6(5)
C(17A)-C(12A)-C(13A)-C(14A)	174.3(3)
C(16A)-N(2A)-C(14A)-O(4A)	-2.8(3)
Cu(1A)-N(2A)-C(14A)-O(4A)	175.24(18)
C(16A)-N(2A)-C(14A)-C(13A)	177.5(3)
Cu(1A)-N(2A)-C(14A)-C(13A)	-4.5(4)
C(15A)-O(4A)-C(14A)-N(2A)	-12.1(3)
C(15A)-O(4A)-C(14A)-C(13A)	167.7(3)
C(12A)-C(13A)-C(14A)-N(2A)	7.9(5)
C(12A)-C(13A)-C(14A)-O(4A)	-171.8(3)
C(14A)-O(4A)-C(15A)-C(16A)	21.0(3)
C(14A)-N(2A)-C(16A)-C(21A)	135.5(3)
Cu(1A)-N(2A)-C(16A)-C(21A)	-42.5(3)
C(14A)-N(2A)-C(16A)-C(15A)	15.5(3)
Cu(1A)-N(2A)-C(16A)-C(15A)	-162.4(2)
C(14A)-N(2A)-C(16A)-C(22A)	-101.1(3)
Cu(1A)-N(2A)-C(16A)-C(22A)	81.0(3)
O(4A)-C(15A)-C(16A)-N(2A)	-21.8(3)
O(4A)-C(15A)-C(16A)-C(21A)	-140.5(3)
O(4A)-C(15A)-C(16A)-C(22A)	93.1(3)
O(3A)-C(12A)-C(17A)-C(19C)	-168.0(4)
C(13A)-C(12A)-C(17A)-C(19C)	14.9(5)
O(3A)-C(12A)-C(17A)-C(20A)	-23.7(5)
C(13A)-C(12A)-C(17A)-C(20A)	159.1(4)
O(3A)-C(12A)-C(17A)-C(18A)	101.3(4)
C(13A)-C(12A)-C(17A)-C(18A)	-75.8(4)

O(3A)-C(12A)-C(17A)-C(20C)	-42.7(5)
C(13A)-C(12A)-C(17A)-C(20C)	140.2(4)
O(3A)-C(12A)-C(17A)-C(18C)	69.4(4)
C(13A)-C(12A)-C(17A)-C(18C)	-107.8(4)
O(3A)-C(12A)-C(17A)-C(19A)	-143.0(4)
C(13A)-C(12A)-C(17A)-C(19A)	39.8(4)
Cu(1B)-O(1B)-C(1B)-C(2B)	-7.3(4)
Cu(1B)-O(1B)-C(1B)-C(6B)	175.30(19)
O(1B)-C(1B)-C(2B)-C(3B)	-0.9(5)
C(6B)-C(1B)-C(2B)-C(3B)	176.4(3)
C(5B)-N(1B)-C(3B)-O(2B)	-3.7(3)
Cu(1B)-N(1B)-C(3B)-O(2B)	172.17(18)
C(5B)-N(1B)-C(3B)-C(2B)	176.3(3)
Cu(1B)-N(1B)-C(3B)-C(2B)	-7.8(4)
C(4B)-O(2B)-C(3B)-N(1B)	-9.9(3)
C(4B)-O(2B)-C(3B)-C(2B)	170.0(3)
C(1B)-C(2B)-C(3B)-N(1B)	9.1(5)
C(1B)-C(2B)-C(3B)-O(2B)	-170.8(3)
C(3B)-O(2B)-C(4B)-C(5B)	18.5(3)
C(3B)-N(1B)-C(5B)-C(11B)	133.4(3)
Cu(1B)-N(1B)-C(5B)-C(11B)	-42.2(3)
C(3B)-N(1B)-C(5B)-C(10B)	-101.7(3)
Cu(1B)-N(1B)-C(5B)-C(10B)	82.7(3)
C(3B)-N(1B)-C(5B)-C(4B)	14.6(3)
Cu(1B)-N(1B)-C(5B)-C(4B)	-161.0(2)
O(2B)-C(4B)-C(5B)-N(1B)	-19.8(3)
O(2B)-C(4B)-C(5B)-C(11B)	-137.8(3)
O(2B)-C(4B)-C(5B)-C(10B)	95.5(3)
O(1B)-C(1B)-C(6B)-C(7B)	71.2(4)
C(2B)-C(1B)-C(6B)-C(7B)	-106.3(4)
O(1B)-C(1B)-C(6B)-C(8B)	-167.4(3)
C(2B)-C(1B)-C(6B)-C(8B)	15.1(4)
O(1B)-C(1B)-C(6B)-C(9B)	-46.7(4)
C(2B)-C(1B)-C(6B)-C(9B)	135.8(3)
Cu(1B)-O(3B)-C(12B)-C(13B)	-5.7(4)
Cu(1B)-O(3B)-C(12B)-C(17B)	175.68(19)

O(3B)-C(12B)-C(13B)-C(14B)	-0.7(5)
C(17B)-C(12B)-C(13B)-C(14B)	177.8(3)
C(16B)-N(2B)-C(14B)-O(4B)	-4.1(3)
Cu(1B)-N(2B)-C(14B)-O(4B)	172.05(18)
C(16B)-N(2B)-C(14B)-C(13B)	176.3(3)
Cu(1B)-N(2B)-C(14B)-C(13B)	-7.6(4)
C(15B)-O(4B)-C(14B)-N(2B)	-12.6(3)
C(15B)-O(4B)-C(14B)-C(13B)	167.0(3)
C(12B)-C(13B)-C(14B)-N(2B)	8.0(5)
C(12B)-C(13B)-C(14B)-O(4B)	-171.6(3)
C(14B)-O(4B)-C(15B)-C(16B)	23.1(3)
C(14B)-N(2B)-C(16B)-C(21B)	136.8(3)
Cu(1B)-N(2B)-C(16B)-C(21B)	-39.1(3)
C(14B)-N(2B)-C(16B)-C(15B)	18.0(3)
Cu(1B)-N(2B)-C(16B)-C(15B)	-158.0(2)
C(14B)-N(2B)-C(16B)-C(22B)	-99.2(3)
Cu(1B)-N(2B)-C(16B)-C(22B)	84.8(3)
O(4B)-C(15B)-C(16B)-N(2B)	-24.4(3)
O(4B)-C(15B)-C(16B)-C(21B)	-142.3(3)
O(4B)-C(15B)-C(16B)-C(22B)	91.1(3)
O(3B)-C(12B)-C(17B)-C(19B)	-176.8(3)
C(13B)-C(12B)-C(17B)-C(19B)	4.5(5)
O(3B)-C(12B)-C(17B)-C(18B)	-52.8(4)
C(13B)-C(12B)-C(17B)-C(18B)	128.6(3)
O(3B)-C(12B)-C(17B)-C(20B)	62.4(4)
C(13B)-C(12B)-C(17B)-C(20B)	-116.3(4)

Symmetry transformations used to generate equivalent atoms:

F9: Crystal data and structure refinement for d1804_a (**15**).

Identification code	d1804_a (15)	
Empirical formula	C ₂₁ H ₂₆ Ir N O ₂	
Formula weight	516.63	
Temperature	150(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P2 ₁ /n	
Unit cell dimensions	a = 8.4355(6) Å	a = 90°.
	b = 12.5712(9) Å	b = 102.943(2)°.
	c = 17.6529(12) Å	g = 90°.
Volume	1824.4(2) Å ³	
Z	4	
Density (calculated)	1.881 Mg/m ³	
Absorption coefficient	7.332 mm ⁻¹	
F(000)	1008	
Crystal size	0.180 x 0.120 x 0.050 mm ³	
Theta range for data collection	2.006 to 27.539°.	
Index ranges	-10 ≤ h ≤ 10, -16 ≤ k ≤ 16, -22 ≤ l ≤ 19	
Reflections collected	23640	
Independent reflections	4189 [R(int) = 0.0637]	
Completeness to theta = 25.242°	100.0 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.7456 and 0.5347	
Refinement method	Full-matrix least-squares on F ²	

Data / restraints / parameters	4189 / 48 / 228
Goodness-of-fit on F^2	0.991
Final R indices [$I > 2\sigma(I)$]	$R_1 = 0.0232$, $wR_2 = 0.0361$
R indices (all data)	$R_1 = 0.0492$, $wR_2 = 0.0410$
Extinction coefficient	n/a
Largest diff. peak and hole	2.276 and -1.027 e. $\text{\AA}^{-3}$

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d1804_a. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	$U(\text{eq})$
Ir(1)	4308(1)	7066(1)	4099(1)	12(1)
O(1)	5373(3)	6368(2)	3296(1)	16(1)
O(2)	8397(3)	5479(2)	5460(1)	20(1)
N(1)	6155(4)	6472(2)	4988(2)	12(1)
C(1)	2209(4)	7040(3)	3180(2)	16(1)
C(2)	3056(4)	8014(3)	3171(2)	16(1)
C(3)	2427(5)	9070(3)	3414(2)	22(1)
C(4)	3110(5)	9298(3)	4274(2)	20(1)
C(5)	3566(5)	8293(3)	4750(2)	18(1)
C(6)	2504(5)	7419(3)	4731(2)	18(1)
C(7)	770(5)	7412(3)	4255(2)	22(1)
C(8)	649(4)	6927(3)	3450(2)	20(1)
C(9)	6603(4)	5715(3)	3410(2)	13(1)
C(10)	7509(4)	5436(3)	4119(2)	14(1)
C(11)	7259(5)	5822(3)	4832(2)	15(1)
C(12)	7794(5)	5796(3)	6129(2)	20(1)
C(13)	6695(5)	6741(3)	5844(2)	15(1)
C(14)	7682(5)	7767(3)	5917(2)	20(1)
C(15)	5347(4)	6782(3)	6284(2)	18(1)
C(16)	6964(4)	5280(3)	2676(2)	15(1)
C(17)	7623(4)	4272(3)	2647(2)	19(1)
C(18)	7922(4)	3879(3)	1959(2)	23(1)
C(19)	7590(5)	4497(3)	1297(2)	27(1)
C(20)	6930(5)	5501(3)	1314(2)	27(1)
C(21)	6610(5)	5891(3)	2003(2)	21(1)

Table 3. Bond lengths [Å] and angles [°] for d1804 a.

Ir(1)-O(1)	2.039(2)
Ir(1)-N(1)	2.087(3)
Ir(1)-C(5)	2.101(4)
Ir(1)-C(2)	2.108(3)
Ir(1)-C(1)	2.117(3)
Ir(1)-C(6)	2.126(4)
O(1)-C(9)	1.303(4)
O(2)-C(11)	1.363(4)
O(2)-C(12)	1.443(4)
N(1)-C(11)	1.314(4)
N(1)-C(13)	1.516(4)
C(1)-C(2)	1.419(5)
C(1)-C(8)	1.504(5)
C(1)-H(1A)	1.0000
C(2)-C(3)	1.527(5)
C(2)-H(2A)	1.0000
C(3)-C(4)	1.525(5)
C(3)-H(3A)	0.9900
C(3)-H(3B)	0.9900
C(4)-C(5)	1.518(5)
C(4)-H(4A)	0.9900
C(4)-H(4B)	0.9900
C(5)-C(6)	1.413(5)
C(5)-H(5A)	1.0000
C(6)-C(7)	1.515(5)
C(6)-H(6A)	1.0000
C(7)-C(8)	1.529(5)
C(7)-H(7A)	0.9900
C(7)-H(7B)	0.9900
C(8)-H(8A)	0.9900
C(8)-H(8B)	0.9900
C(9)-C(10)	1.358(5)
C(9)-C(16)	1.499(5)
C(10)-C(11)	1.408(5)
C(10)-H(10A)	0.9500

C(12)-C(13)	1.521(5)
C(12)-H(12A)	0.9900
C(12)-H(12B)	0.9900
C(13)-C(15)	1.514(5)
C(13)-C(14)	1.525(5)
C(14)-H(14A)	0.9800
C(14)-H(14B)	0.9800
C(14)-H(14C)	0.9800
C(15)-H(15A)	0.9800
C(15)-H(15B)	0.9800
C(15)-H(15C)	0.9800
C(16)-C(17)	1.390(5)
C(16)-C(21)	1.390(5)
C(17)-C(18)	1.386(5)
C(17)-H(17A)	0.9500
C(18)-C(19)	1.379(5)
C(18)-H(18A)	0.9500
C(19)-C(20)	1.382(6)
C(19)-H(19A)	0.9500
C(20)-C(21)	1.393(5)
C(20)-H(20A)	0.9500
C(21)-H(21A)	0.9500
O(1)-Ir(1)-N(1)	89.87(10)
O(1)-Ir(1)-C(5)	158.12(13)
N(1)-Ir(1)-C(5)	96.61(13)
O(1)-Ir(1)-C(2)	85.69(12)
N(1)-Ir(1)-C(2)	161.21(13)
C(5)-Ir(1)-C(2)	81.46(14)
O(1)-Ir(1)-C(1)	83.31(12)
N(1)-Ir(1)-C(1)	157.88(14)
C(5)-Ir(1)-C(1)	97.48(15)
C(2)-Ir(1)-C(1)	39.24(13)
O(1)-Ir(1)-C(6)	159.01(12)
N(1)-Ir(1)-C(6)	100.44(13)
C(5)-Ir(1)-C(6)	39.06(13)

C(2)-Ir(1)-C(6)	89.85(14)
C(1)-Ir(1)-C(6)	80.32(14)
C(9)-O(1)-Ir(1)	128.5(2)
C(11)-O(2)-C(12)	105.4(3)
C(11)-N(1)-C(13)	106.2(3)
C(11)-N(1)-Ir(1)	120.6(2)
C(13)-N(1)-Ir(1)	132.6(2)
C(2)-C(1)-C(8)	124.3(3)
C(2)-C(1)-Ir(1)	70.05(19)
C(8)-C(1)-Ir(1)	113.5(2)
C(2)-C(1)-H(1A)	113.8
C(8)-C(1)-H(1A)	113.8
Ir(1)-C(1)-H(1A)	113.8
C(1)-C(2)-C(3)	122.7(3)
C(1)-C(2)-Ir(1)	70.71(19)
C(3)-C(2)-Ir(1)	114.7(2)
C(1)-C(2)-H(2A)	113.9
C(3)-C(2)-H(2A)	113.9
Ir(1)-C(2)-H(2A)	113.9
C(4)-C(3)-C(2)	111.4(3)
C(4)-C(3)-H(3A)	109.3
C(2)-C(3)-H(3A)	109.3
C(4)-C(3)-H(3B)	109.3
C(2)-C(3)-H(3B)	109.3
H(3A)-C(3)-H(3B)	108.0
C(5)-C(4)-C(3)	112.8(3)
C(5)-C(4)-H(4A)	109.0
C(3)-C(4)-H(4A)	109.0
C(5)-C(4)-H(4B)	109.0
C(3)-C(4)-H(4B)	109.0
H(4A)-C(4)-H(4B)	107.8
C(6)-C(5)-C(4)	123.6(3)
C(6)-C(5)-Ir(1)	71.4(2)
C(4)-C(5)-Ir(1)	112.2(2)
C(6)-C(5)-H(5A)	114.1
C(4)-C(5)-H(5A)	114.1

Ir(1)-C(5)-H(5A)	114.1
C(5)-C(6)-C(7)	123.1(4)
C(5)-C(6)-Ir(1)	69.5(2)
C(7)-C(6)-Ir(1)	115.0(3)
C(5)-C(6)-H(6A)	113.9
C(7)-C(6)-H(6A)	113.9
Ir(1)-C(6)-H(6A)	113.9
C(6)-C(7)-C(8)	111.9(3)
C(6)-C(7)-H(7A)	109.2
C(8)-C(7)-H(7A)	109.2
C(6)-C(7)-H(7B)	109.2
C(8)-C(7)-H(7B)	109.2
H(7A)-C(7)-H(7B)	107.9
C(1)-C(8)-C(7)	112.1(3)
C(1)-C(8)-H(8A)	109.2
C(7)-C(8)-H(8A)	109.2
C(1)-C(8)-H(8B)	109.2
C(7)-C(8)-H(8B)	109.2
H(8A)-C(8)-H(8B)	107.9
O(1)-C(9)-C(10)	124.8(3)
O(1)-C(9)-C(16)	113.9(3)
C(10)-C(9)-C(16)	121.4(3)
C(9)-C(10)-C(11)	124.6(3)
C(9)-C(10)-H(10A)	117.7
C(11)-C(10)-H(10A)	117.7
N(1)-C(11)-O(2)	115.4(3)
N(1)-C(11)-C(10)	131.1(3)
O(2)-C(11)-C(10)	113.5(3)
O(2)-C(12)-C(13)	104.2(3)
O(2)-C(12)-H(12A)	110.9
C(13)-C(12)-H(12A)	110.9
O(2)-C(12)-H(12B)	110.9
C(13)-C(12)-H(12B)	110.9
H(12A)-C(12)-H(12B)	108.9
C(15)-C(13)-N(1)	115.2(3)
C(15)-C(13)-C(12)	109.3(3)

N(1)-C(13)-C(12)	100.4(3)
C(15)-C(13)-C(14)	112.7(3)
N(1)-C(13)-C(14)	108.0(3)
C(12)-C(13)-C(14)	110.6(3)
C(13)-C(14)-H(14A)	109.5
C(13)-C(14)-H(14B)	109.5
H(14A)-C(14)-H(14B)	109.5
C(13)-C(14)-H(14C)	109.5
H(14A)-C(14)-H(14C)	109.5
H(14B)-C(14)-H(14C)	109.5
C(13)-C(15)-H(15A)	109.5
C(13)-C(15)-H(15B)	109.5
H(15A)-C(15)-H(15B)	109.5
C(13)-C(15)-H(15C)	109.5
H(15A)-C(15)-H(15C)	109.5
H(15B)-C(15)-H(15C)	109.5
C(17)-C(16)-C(21)	118.9(3)
C(17)-C(16)-C(9)	121.7(3)
C(21)-C(16)-C(9)	119.5(3)
C(18)-C(17)-C(16)	120.7(4)
C(18)-C(17)-H(17A)	119.7
C(16)-C(17)-H(17A)	119.7
C(19)-C(18)-C(17)	120.0(4)
C(19)-C(18)-H(18A)	120.0
C(17)-C(18)-H(18A)	120.0
C(18)-C(19)-C(20)	120.2(4)
C(18)-C(19)-H(19A)	119.9
C(20)-C(19)-H(19A)	119.9
C(19)-C(20)-C(21)	119.9(4)
C(19)-C(20)-H(20A)	120.1
C(21)-C(20)-H(20A)	120.1
C(16)-C(21)-C(20)	120.4(4)
C(16)-C(21)-H(21A)	119.8
C(20)-C(21)-H(21A)	119.8

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d1804_a. The anisotropic displacement factor exponent takes the form: $-2p^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Ir(1)	12(1)	12(1)	11(1)	0(1)	2(1)	1(1)
O(1)	17(2)	21(2)	12(1)	-1(1)	4(1)	3(1)
O(2)	19(2)	25(2)	14(1)	0(1)	-2(1)	5(1)
N(1)	13(2)	14(2)	9(2)	-1(1)	1(1)	-2(1)
C(1)	14(2)	21(2)	11(2)	0(2)	-3(1)	3(2)
C(2)	16(2)	21(2)	7(2)	3(2)	-6(1)	3(2)
C(3)	25(2)	19(2)	20(2)	1(2)	0(2)	4(2)
C(4)	23(2)	16(2)	20(2)	-2(2)	4(2)	2(2)
C(5)	19(2)	21(2)	15(2)	-3(1)	5(2)	2(2)
C(6)	20(2)	21(2)	15(2)	-2(2)	5(2)	2(2)
C(7)	15(2)	28(2)	22(2)	-1(2)	5(2)	3(2)
C(8)	13(2)	23(2)	23(2)	2(2)	0(2)	1(2)
C(9)	10(2)	12(2)	19(2)	-2(2)	6(2)	-6(2)
C(10)	11(2)	11(2)	18(2)	-2(2)	0(2)	0(2)
C(11)	17(2)	14(2)	13(2)	2(2)	-2(2)	-5(2)
C(12)	22(2)	23(2)	15(2)	2(2)	1(2)	3(2)
C(13)	16(2)	18(2)	9(2)	0(2)	0(2)	-2(2)
C(14)	22(2)	22(2)	16(2)	-1(2)	1(2)	-5(2)
C(15)	19(2)	24(2)	9(2)	0(2)	1(2)	-1(2)
C(16)	10(2)	20(2)	15(2)	-5(2)	3(2)	-4(2)
C(17)	17(2)	19(2)	21(2)	-4(2)	4(2)	0(2)
C(18)	18(2)	21(2)	31(2)	-10(2)	6(2)	1(2)
C(19)	26(2)	35(3)	25(2)	-11(2)	16(2)	-2(2)
C(20)	34(3)	31(3)	17(2)	1(2)	8(2)	-4(2)
C(21)	21(2)	23(2)	20(2)	1(2)	5(2)	3(2)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for d1804_a.

	x	y	z	U(eq)
H(1A)	2297	6546	2747	20
H(2A)	3620	8076	2731	19
H(3A)	2742	9652	3099	26
H(3B)	1225	9049	3311	26
H(4A)	4086	9754	4328	23
H(4B)	2290	9696	4483	23
H(5A)	4315	8416	5268	22
H(6A)	2654	7032	5237	22
H(7A)	351	8150	4197	26
H(7B)	80	6998	4534	26
H(8A)	373	6163	3465	24
H(8B)	-239	7281	3073	24
H(10A)	8373	4946	4135	17
H(12A)	8699	6004	6564	24
H(12B)	7174	5211	6303	24
H(14A)	8630	7663	5689	30
H(14B)	8048	7956	6467	30
H(14C)	7003	8340	5640	30
H(15A)	5819	6745	6844	26
H(15B)	4609	6180	6129	26
H(15C)	4742	7450	6164	26
H(17A)	7871	3847	3104	23
H(18A)	8356	3184	1943	28
H(19A)	7815	4232	828	32
H(20A)	6696	5924	856	32
H(21A)	6147	6578	2012	25

Table 6. Torsion angles [°] for d1804_a.

C(8)-C(1)-C(2)-C(3)	-2.1(5)
Ir(1)-C(1)-C(2)-C(3)	-107.5(3)
C(8)-C(1)-C(2)-Ir(1)	105.5(3)
C(1)-C(2)-C(3)-C(4)	92.4(4)
Ir(1)-C(2)-C(3)-C(4)	10.2(4)
C(2)-C(3)-C(4)-C(5)	-26.2(5)
C(3)-C(4)-C(5)-C(6)	-51.4(5)
C(3)-C(4)-C(5)-Ir(1)	30.5(4)
C(4)-C(5)-C(6)-C(7)	-2.4(6)
Ir(1)-C(5)-C(6)-C(7)	-107.2(3)
C(4)-C(5)-C(6)-Ir(1)	104.8(3)
C(5)-C(6)-C(7)-C(8)	92.8(4)
Ir(1)-C(6)-C(7)-C(8)	11.8(4)
C(2)-C(1)-C(8)-C(7)	-51.7(5)
Ir(1)-C(1)-C(8)-C(7)	29.5(4)
C(6)-C(7)-C(8)-C(1)	-26.4(5)
Ir(1)-O(1)-C(9)-C(10)	-5.5(5)
Ir(1)-O(1)-C(9)-C(16)	174.7(2)
O(1)-C(9)-C(10)-C(11)	-0.8(6)
C(16)-C(9)-C(10)-C(11)	179.1(3)
C(13)-N(1)-C(11)-O(2)	7.8(4)
Ir(1)-N(1)-C(11)-O(2)	-179.5(2)
C(13)-N(1)-C(11)-C(10)	-170.0(4)
Ir(1)-N(1)-C(11)-C(10)	2.7(5)
C(12)-O(2)-C(11)-N(1)	11.4(4)
C(12)-O(2)-C(11)-C(10)	-170.4(3)
C(9)-C(10)-C(11)-N(1)	2.1(6)
C(9)-C(10)-C(11)-O(2)	-175.7(3)
C(11)-O(2)-C(12)-C(13)	-25.1(4)
C(11)-N(1)-C(13)-C(15)	-139.4(3)
Ir(1)-N(1)-C(13)-C(15)	49.1(4)
C(11)-N(1)-C(13)-C(12)	-22.2(4)
Ir(1)-N(1)-C(13)-C(12)	166.3(2)
C(11)-N(1)-C(13)-C(14)	93.6(3)

Ir(1)-N(1)-C(13)-C(14)	-77.8(4)
O(2)-C(12)-C(13)-C(15)	149.8(3)
O(2)-C(12)-C(13)-N(1)	28.3(3)
O(2)-C(12)-C(13)-C(14)	-85.5(3)
O(1)-C(9)-C(16)-C(17)	-149.1(3)
C(10)-C(9)-C(16)-C(17)	31.1(5)
O(1)-C(9)-C(16)-C(21)	29.9(5)
C(10)-C(9)-C(16)-C(21)	-149.9(4)
C(21)-C(16)-C(17)-C(18)	-0.1(6)
C(9)-C(16)-C(17)-C(18)	178.9(3)
C(16)-C(17)-C(18)-C(19)	1.0(6)
C(17)-C(18)-C(19)-C(20)	-1.2(6)
C(18)-C(19)-C(20)-C(21)	0.4(6)
C(17)-C(16)-C(21)-C(20)	-0.7(6)
C(9)-C(16)-C(21)-C(20)	-179.8(4)
C(19)-C(20)-C(21)-C(16)	0.6(6)

Symmetry transformations used to generate equivalent atoms:

F10: Crystal data and structure refinement for d1872_a (**16**).

Identification code	d1872_a (16)	
Empirical formula	C ₂₂ H ₂₉ I Ir N O ₂	
Formula weight	658.56	
Temperature	150(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P2 ₁ /n	
Unit cell dimensions	a = 14.6277(6) Å	α = 90°.
	b = 9.7486(4) Å	β = 101.6540(10)°.
	c = 14.6525(6) Å	γ = 90°.
Volume	2046.37(15) Å ³	
Z	4	
Density (calculated)	2.138 Mg/m ³	
Absorption coefficient	8.050 mm ⁻¹	
F(000)	1256	
Crystal size	0.180 x 0.170 x 0.040 mm ³	
Theta range for data collection	1.794 to 27.541°.	
Index ranges	-19 ≤ h ≤ 19, -12 ≤ k ≤ 12, -19 ≤ l ≤ 19	
Reflections collected	30365	
Independent reflections	4701 [R(int) = 0.0325]	
Completeness to theta = 25.242°	100.0 %	

Absorption correction	Semi-empirical from equivalents
Max. and min. transmission	0.7456 and 0.5749
Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	4701 / 0 / 247
Goodness-of-fit on F ²	1.069
Final R indices [I>2sigma(I)]	R1 = 0.0178, wR2 = 0.0343
R indices (all data)	R1 = 0.0273, wR2 = 0.0367
Extinction coefficient	n/a
Largest diff. peak and hole	0.741 and -0.905 e. ^{~3}

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\approx 2 \times 10^3$) for d1872_a. U(eq) is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	U(eq)
Ir(1)	2177(1)	3248(1)	4367(1)	12(1)
I(1)	3162(1)	3484(1)	6255(1)	18(1)
O(1)	3401(1)	2494(2)	4104(2)	16(1)
O(2)	3748(1)	6843(2)	4180(2)	17(1)
N(1)	2646(2)	5285(2)	4314(2)	12(1)
C(1)	4109(2)	3199(3)	3951(2)	13(1)
C(2)	4185(2)	4595(3)	3990(2)	15(1)
C(3)	3489(2)	5506(3)	4173(2)	14(1)
C(4)	3037(2)	7619(3)	4499(2)	17(1)
C(5)	2187(2)	6656(3)	4378(2)	16(1)
C(6)	4858(2)	2318(3)	3693(2)	13(1)
C(7)	4866(2)	916(3)	3902(2)	19(1)
C(8)	5518(2)	56(3)	3634(2)	23(1)
C(9)	6165(2)	570(4)	3150(2)	24(1)
C(10)	6173(2)	1953(4)	2960(2)	23(1)
C(11)	5525(2)	2835(3)	3227(2)	17(1)
C(12)	1727(2)	6762(3)	5220(2)	19(1)
C(13)	1508(2)	7005(3)	3474(2)	19(1)
C(14)	676(2)	3674(3)	4101(2)	16(1)
C(15)	137(2)	2639(3)	3429(2)	20(1)
C(16)	594(2)	1211(3)	3488(2)	21(1)
C(17)	1637(2)	1232(3)	3872(2)	19(1)
C(18)	2036(2)	1117(3)	4821(2)	18(1)
C(19)	1476(2)	982(3)	5572(2)	20(1)
C(20)	727(2)	2095(3)	5505(2)	18(1)
C(21)	970(2)	3425(3)	5057(2)	15(1)
C(22)	1882(2)	3577(3)	2861(2)	15(1)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for d1872_a.

Ir(1)-O(1)	2.0426(19)
Ir(1)-N(1)	2.108(2)
Ir(1)-C(22)	2.184(3)
Ir(1)-C(17)	2.187(3)
Ir(1)-C(14)	2.191(3)
Ir(1)-C(18)	2.204(3)
Ir(1)-C(21)	2.211(3)
Ir(1)-I(1)	2.8598(2)
O(1)-C(1)	1.299(3)
O(2)-C(3)	1.357(3)
O(2)-C(4)	1.439(3)
N(1)-C(3)	1.309(4)
N(1)-C(5)	1.507(3)
C(1)-C(2)	1.366(4)
C(1)-C(6)	1.500(4)
C(2)-C(3)	1.417(4)
C(2)-H(2A)	0.9500
C(4)-C(5)	1.539(4)
C(4)-H(4A)	0.9900
C(4)-H(4B)	0.9900
C(5)-C(12)	1.524(4)
C(5)-C(13)	1.525(4)
C(6)-C(11)	1.394(4)
C(6)-C(7)	1.400(4)
C(7)-C(8)	1.385(4)
C(7)-H(7A)	0.9500
C(8)-C(9)	1.385(5)
C(8)-H(8A)	0.9500
C(9)-C(10)	1.378(5)
C(9)-H(9A)	0.9500
C(10)-C(11)	1.392(4)
C(10)-H(10A)	0.9500
C(11)-H(11A)	0.9500
C(12)-H(12A)	0.9800

C(12)-H(12B)	0.9800
C(12)-H(12C)	0.9800
C(13)-H(13A)	0.9800
C(13)-H(13B)	0.9800
C(13)-H(13C)	0.9800
C(14)-C(21)	1.401(4)
C(14)-C(15)	1.516(4)
C(14)-H(14A)	1.0000
C(15)-C(16)	1.539(4)
C(15)-H(15A)	0.9900
C(15)-H(15B)	0.9900
C(16)-C(17)	1.516(4)
C(16)-H(16A)	0.9900
C(16)-H(16B)	0.9900
C(17)-C(18)	1.401(5)
C(17)-H(17A)	1.0000
C(18)-C(19)	1.503(4)
C(18)-H(18A)	1.0000
C(19)-C(20)	1.530(4)
C(19)-H(19A)	0.9900
C(19)-H(19B)	0.9900
C(20)-C(21)	1.527(4)
C(20)-H(20A)	0.9900
C(20)-H(20B)	0.9900
C(21)-H(21A)	1.0000
C(22)-H(22A)	0.9800
C(22)-H(22B)	0.9800
C(22)-H(22C)	0.9800
O(1)-Ir(1)-N(1)	91.57(8)
O(1)-Ir(1)-C(22)	82.21(10)
N(1)-Ir(1)-C(22)	79.87(10)
O(1)-Ir(1)-C(17)	83.47(10)
N(1)-Ir(1)-C(17)	158.96(11)
C(22)-Ir(1)-C(17)	79.20(12)
O(1)-Ir(1)-C(14)	156.93(10)

N(1)-Ir(1)-C(14)	97.87(10)
C(22)-Ir(1)-C(14)	78.83(11)
C(17)-Ir(1)-C(14)	80.19(11)
O(1)-Ir(1)-C(18)	81.78(10)
N(1)-Ir(1)-C(18)	161.82(11)
C(22)-Ir(1)-C(18)	115.62(11)
C(17)-Ir(1)-C(18)	37.21(12)
C(14)-Ir(1)-C(18)	94.65(12)
O(1)-Ir(1)-C(21)	157.73(9)
N(1)-Ir(1)-C(21)	103.83(10)
C(22)-Ir(1)-C(21)	115.93(11)
C(17)-Ir(1)-C(21)	87.35(11)
C(14)-Ir(1)-C(21)	37.10(11)
C(18)-Ir(1)-C(21)	78.67(11)
O(1)-Ir(1)-I(1)	85.50(6)
N(1)-Ir(1)-I(1)	81.65(6)
C(22)-Ir(1)-I(1)	157.45(8)
C(17)-Ir(1)-I(1)	118.12(9)
C(14)-Ir(1)-I(1)	116.60(8)
C(18)-Ir(1)-I(1)	80.98(8)
C(21)-Ir(1)-I(1)	81.03(8)
C(1)-O(1)-Ir(1)	126.99(18)
C(3)-O(2)-C(4)	106.7(2)
C(3)-N(1)-C(5)	108.0(2)
C(3)-N(1)-Ir(1)	119.03(19)
C(5)-N(1)-Ir(1)	132.90(18)
O(1)-C(1)-C(2)	125.6(3)
O(1)-C(1)-C(6)	112.9(2)
C(2)-C(1)-C(6)	121.5(3)
C(1)-C(2)-C(3)	125.2(3)
C(1)-C(2)-H(2A)	117.4
C(3)-C(2)-H(2A)	117.4
N(1)-C(3)-O(2)	115.3(2)
N(1)-C(3)-C(2)	131.5(3)
O(2)-C(3)-C(2)	113.1(3)
O(2)-C(4)-C(5)	105.2(2)

O(2)-C(4)-H(4A)	110.7
C(5)-C(4)-H(4A)	110.7
O(2)-C(4)-H(4B)	110.7
C(5)-C(4)-H(4B)	110.7
H(4A)-C(4)-H(4B)	108.8
N(1)-C(5)-C(12)	112.7(2)
N(1)-C(5)-C(13)	111.3(2)
C(12)-C(5)-C(13)	111.8(3)
N(1)-C(5)-C(4)	100.9(2)
C(12)-C(5)-C(4)	110.1(3)
C(13)-C(5)-C(4)	109.4(2)
C(11)-C(6)-C(7)	119.0(3)
C(11)-C(6)-C(1)	122.4(3)
C(7)-C(6)-C(1)	118.6(3)
C(8)-C(7)-C(6)	120.3(3)
C(8)-C(7)-H(7A)	119.8
C(6)-C(7)-H(7A)	119.8
C(9)-C(8)-C(7)	120.4(3)
C(9)-C(8)-H(8A)	119.8
C(7)-C(8)-H(8A)	119.8
C(10)-C(9)-C(8)	119.5(3)
C(10)-C(9)-H(9A)	120.2
C(8)-C(9)-H(9A)	120.2
C(9)-C(10)-C(11)	120.8(3)
C(9)-C(10)-H(10A)	119.6
C(11)-C(10)-H(10A)	119.6
C(10)-C(11)-C(6)	119.9(3)
C(10)-C(11)-H(11A)	120.1
C(6)-C(11)-H(11A)	120.1
C(5)-C(12)-H(12A)	109.5
C(5)-C(12)-H(12B)	109.5
H(12A)-C(12)-H(12B)	109.5
C(5)-C(12)-H(12C)	109.5
H(12A)-C(12)-H(12C)	109.5
H(12B)-C(12)-H(12C)	109.5
C(5)-C(13)-H(13A)	109.5

C(5)-C(13)-H(13B)	109.5
H(13A)-C(13)-H(13B)	109.5
C(5)-C(13)-H(13C)	109.5
H(13A)-C(13)-H(13C)	109.5
H(13B)-C(13)-H(13C)	109.5
C(21)-C(14)-C(15)	123.3(3)
C(21)-C(14)-Ir(1)	72.25(17)
C(15)-C(14)-Ir(1)	111.1(2)
C(21)-C(14)-H(14A)	114.3
C(15)-C(14)-H(14A)	114.3
Ir(1)-C(14)-H(14A)	114.3
C(14)-C(15)-C(16)	113.7(3)
C(14)-C(15)-H(15A)	108.8
C(16)-C(15)-H(15A)	108.8
C(14)-C(15)-H(15B)	108.8
C(16)-C(15)-H(15B)	108.8
H(15A)-C(15)-H(15B)	107.7
C(17)-C(16)-C(15)	113.7(2)
C(17)-C(16)-H(16A)	108.8
C(15)-C(16)-H(16A)	108.8
C(17)-C(16)-H(16B)	108.8
C(15)-C(16)-H(16B)	108.8
H(16A)-C(16)-H(16B)	107.7
C(18)-C(17)-C(16)	123.7(3)
C(18)-C(17)-Ir(1)	72.06(17)
C(16)-C(17)-Ir(1)	113.8(2)
C(18)-C(17)-H(17A)	113.5
C(16)-C(17)-H(17A)	113.5
Ir(1)-C(17)-H(17A)	113.5
C(17)-C(18)-C(19)	123.6(3)
C(17)-C(18)-Ir(1)	70.73(17)
C(19)-C(18)-Ir(1)	113.7(2)
C(17)-C(18)-H(18A)	113.8
C(19)-C(18)-H(18A)	113.8
Ir(1)-C(18)-H(18A)	113.8
C(18)-C(19)-C(20)	112.3(3)

C(18)-C(19)-H(19A)	109.1
C(20)-C(19)-H(19A)	109.1
C(18)-C(19)-H(19B)	109.1
C(20)-C(19)-H(19B)	109.1
H(19A)-C(19)-H(19B)	107.9
C(21)-C(20)-C(19)	113.8(3)
C(21)-C(20)-H(20A)	108.8
C(19)-C(20)-H(20A)	108.8
C(21)-C(20)-H(20B)	108.8
C(19)-C(20)-H(20B)	108.8
H(20A)-C(20)-H(20B)	107.7
C(14)-C(21)-C(20)	121.7(3)
C(14)-C(21)-Ir(1)	70.65(17)
C(20)-C(21)-Ir(1)	113.97(19)
C(14)-C(21)-H(21A)	114.4
C(20)-C(21)-H(21A)	114.4
Ir(1)-C(21)-H(21A)	114.4
Ir(1)-C(22)-H(22A)	109.5
Ir(1)-C(22)-H(22B)	109.5
H(22A)-C(22)-H(22B)	109.5
Ir(1)-C(22)-H(22C)	109.5
H(22A)-C(22)-H(22C)	109.5
H(22B)-C(22)-H(22C)	109.5

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\approx 2 \times 10^3$) for d1872_a. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Ir(1)	9(1)	12(1)	13(1)	-2(1)	3(1)	1(1)
I(1)	16(1)	18(1)	17(1)	2(1)	0(1)	0(1)
O(1)	10(1)	14(1)	24(1)	-2(1)	7(1)	2(1)
O(2)	15(1)	14(1)	23(1)	0(1)	6(1)	1(1)
N(1)	10(1)	13(1)	13(1)	-1(1)	3(1)	4(1)
C(1)	9(1)	20(1)	10(2)	-1(1)	1(1)	2(1)
C(2)	12(2)	20(2)	16(2)	1(1)	6(1)	1(1)
C(3)	16(2)	14(1)	11(2)	0(1)	1(1)	2(1)
C(4)	19(2)	14(1)	18(2)	-1(1)	3(1)	3(1)
C(5)	16(2)	11(1)	19(2)	0(1)	3(1)	2(1)
C(6)	10(2)	19(1)	10(2)	-3(1)	-1(1)	1(1)
C(7)	16(2)	20(2)	20(2)	-2(1)	2(1)	4(1)
C(8)	24(2)	22(2)	21(2)	-5(1)	-2(2)	6(1)
C(9)	17(2)	37(2)	17(2)	-10(2)	-1(1)	13(2)
C(10)	16(2)	39(2)	14(2)	-1(1)	4(1)	5(1)
C(11)	13(2)	25(2)	13(2)	1(1)	2(1)	3(1)
C(12)	21(2)	16(2)	22(2)	-2(1)	8(1)	6(1)
C(13)	16(2)	20(2)	20(2)	6(1)	0(1)	2(1)
C(14)	7(1)	18(2)	22(2)	-1(1)	4(1)	4(1)
C(15)	12(2)	28(2)	19(2)	-2(1)	0(1)	-3(1)
C(16)	19(2)	24(2)	21(2)	-10(1)	3(1)	-10(1)
C(17)	17(2)	12(1)	28(2)	-8(1)	8(1)	-2(1)
C(18)	18(2)	10(1)	28(2)	-3(1)	6(1)	-1(1)
C(19)	17(2)	16(2)	27(2)	4(1)	5(1)	-3(1)
C(20)	16(2)	20(2)	20(2)	2(1)	5(1)	1(1)
C(21)	11(2)	18(2)	18(2)	-2(1)	6(1)	4(1)
C(22)	4(1)	17(2)	24(2)	-4(1)	4(1)	-2(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\approx^2 \times 10^{-3}$) for d1872_a.

	x	y	z	U(eq)
H(2A)	4750	4987	3885	18
H(4A)	2883	8461	4123	20
H(4B)	3247	7881	5161	20
H(7A)	4423	553	4230	23
H(8A)	5522	-893	3783	28
H(9A)	6599	-27	2951	29
H(10A)	6627	2311	2642	27
H(11A)	5538	3787	3092	21
H(12A)	1127	6279	5089	29
H(12B)	1623	7729	5350	29
H(12C)	2134	6345	5763	29
H(13A)	983	6362	3380	29
H(13B)	1830	6935	2950	29
H(13C)	1276	7942	3510	29
H(14A)	509	4652	3941	19
H(15A)	75	2992	2785	24
H(15B)	-500	2548	3557	24
H(16A)	293	616	3889	25
H(16B)	477	800	2857	25
H(17A)	2009	773	3457	22
H(18A)	2637	596	4952	22
H(19A)	1902	1040	6188	24
H(19B)	1173	70	5523	24
H(20A)	133	1734	5138	22
H(20B)	629	2301	6139	22
H(21A)	981	4257	5458	18
H(22A)	1430	4325	2704	23
H(22B)	1623	2736	2545	23
H(22C)	2461	3818	2660	23

Table 6. Torsion angles [$^{\circ}$] for d1872_a.

Ir(1)-O(1)-C(1)-C(2)	3.9(4)
Ir(1)-O(1)-C(1)-C(6)	-174.50(18)
O(1)-C(1)-C(2)-C(3)	-2.2(5)
C(6)-C(1)-C(2)-C(3)	176.1(3)
C(5)-N(1)-C(3)-O(2)	3.9(3)
Ir(1)-N(1)-C(3)-O(2)	-177.90(18)
C(5)-N(1)-C(3)-C(2)	-174.3(3)
Ir(1)-N(1)-C(3)-C(2)	3.8(4)
C(4)-O(2)-C(3)-N(1)	9.2(3)
C(4)-O(2)-C(3)-C(2)	-172.2(2)
C(1)-C(2)-C(3)-N(1)	-2.3(5)
C(1)-C(2)-C(3)-O(2)	179.4(3)
C(3)-O(2)-C(4)-C(5)	-17.7(3)
C(3)-N(1)-C(5)-C(12)	-131.6(3)
Ir(1)-N(1)-C(5)-C(12)	50.6(4)
C(3)-N(1)-C(5)-C(13)	101.8(3)
Ir(1)-N(1)-C(5)-C(13)	-76.0(3)
C(3)-N(1)-C(5)-C(4)	-14.2(3)
Ir(1)-N(1)-C(5)-C(4)	168.0(2)
O(2)-C(4)-C(5)-N(1)	19.0(3)
O(2)-C(4)-C(5)-C(12)	138.2(2)
O(2)-C(4)-C(5)-C(13)	-98.5(3)
O(1)-C(1)-C(6)-C(11)	159.6(3)
C(2)-C(1)-C(6)-C(11)	-18.9(4)
O(1)-C(1)-C(6)-C(7)	-18.3(4)
C(2)-C(1)-C(6)-C(7)	163.2(3)
C(11)-C(6)-C(7)-C(8)	-1.2(5)
C(1)-C(6)-C(7)-C(8)	176.8(3)
C(6)-C(7)-C(8)-C(9)	-0.5(5)
C(7)-C(8)-C(9)-C(10)	1.9(5)
C(8)-C(9)-C(10)-C(11)	-1.7(5)
C(9)-C(10)-C(11)-C(6)	0.0(5)
C(7)-C(6)-C(11)-C(10)	1.4(4)
C(1)-C(6)-C(11)-C(10)	-176.5(3)

C(21)-C(14)-C(15)-C(16)	-51.5(4)
Ir(1)-C(14)-C(15)-C(16)	30.6(3)
C(14)-C(15)-C(16)-C(17)	-24.2(4)
C(15)-C(16)-C(17)-C(18)	89.6(4)
C(15)-C(16)-C(17)-Ir(1)	5.9(4)
C(16)-C(17)-C(18)-C(19)	-1.0(5)
Ir(1)-C(17)-C(18)-C(19)	106.1(3)
C(16)-C(17)-C(18)-Ir(1)	-107.2(3)
C(17)-C(18)-C(19)-C(20)	-51.4(4)
Ir(1)-C(18)-C(19)-C(20)	30.5(3)
C(18)-C(19)-C(20)-C(21)	-27.6(4)
C(15)-C(14)-C(21)-C(20)	-2.8(4)
Ir(1)-C(14)-C(21)-C(20)	-106.8(3)
C(15)-C(14)-C(21)-Ir(1)	104.0(3)
C(19)-C(20)-C(21)-C(14)	93.4(3)
C(19)-C(20)-C(21)-Ir(1)	12.1(3)

Symmetry transformations used to generate equivalent atoms:

F11: Crystal data and structure refinement for **18**.

Identification code	d18124_a (18)	
Empirical formula	C ₃₉ H ₄₂ Ir N O ₂ Sn	
Formula weight	867.62	
Temperature	150(2) K	
Wavelength	0.71073 Å	
Crystal system	Monoclinic	
Space group	P2 ₁ /n	
Unit cell dimensions	a = 13.2953(8) Å	α = 90°.
	b = 17.8294(12) Å	β = 92.525(2)°.
	c = 14.0144(9) Å	γ = 90°.
Volume	3318.8(4) Å ³	
Z	4	
Density (calculated)	1.736 Mg/m ³	
Absorption coefficient	4.793 mm ⁻¹	
F(000)	1704	
Crystal size	0.130 x 0.120 x 0.090 mm ³	
Theta range for data collection	1.849 to 27.533°.	
Index ranges	-17 ≤ h ≤ 16, -23 ≤ k ≤ 23, -18 ≤ l ≤ 16	
Reflections collected	70635	
Independent reflections	7637 [R(int) = 0.0404]	
Completeness to theta = 25.242°	100.0 %	
Absorption correction	Semi-empirical from equivalents	
Max. and min. transmission	0.7456 and 0.6796	

Refinement method	Full-matrix least-squares on F ²
Data / restraints / parameters	7637 / 0 / 403
Goodness-of-fit on F ²	1.048
Final R indices [I>2sigma(I)]	R1 = 0.0199, wR2 = 0.0359
R indices (all data)	R1 = 0.0320, wR2 = 0.0399
Extinction coefficient	n/a
Largest diff. peak and hole	1.669 and -0.604 e.Å ⁻³

Table 2. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d18124_a. $U(\text{eq})$ is defined as one third of the trace of the orthogonalized U^{ij} tensor.

	x	y	z	$U(\text{eq})$
Ir(1)	5417(1)	1734(1)	6988(1)	15(1)
Sn(1)	5982(1)	3097(1)	7440(1)	16(1)
O(1)	5848(1)	1998(1)	5558(1)	21(1)
O(2)	8341(2)	803(1)	6850(1)	32(1)
N(1)	6861(2)	1286(1)	7244(2)	17(1)
C(1)	6591(2)	1702(2)	5136(2)	20(1)
C(2)	7346(2)	1276(2)	5567(2)	26(1)
C(3)	7457(2)	1138(2)	6557(2)	22(1)
C(4)	8241(2)	604(2)	7840(2)	30(1)
C(5)	7410(2)	1119(1)	8182(2)	20(1)
C(6)	6595(2)	1874(2)	4085(2)	22(1)
C(7)	6018(2)	2472(2)	3731(2)	26(1)
C(8)	5994(2)	2652(2)	2766(2)	31(1)
C(9)	6537(2)	2229(2)	2142(2)	30(1)
C(10)	7101(2)	1628(2)	2480(2)	30(1)
C(11)	7133(2)	1452(2)	3441(2)	28(1)
C(12)	6775(2)	721(2)	8897(2)	34(1)
C(13)	7852(2)	1838(2)	8609(2)	33(1)
C(14)	4817(2)	538(2)	7169(2)	28(1)
C(15)	4842(2)	668(2)	6202(2)	30(1)
C(16)	3912(3)	848(2)	5573(3)	39(1)
C(17)	3672(2)	1680(2)	5495(2)	29(1)
C(18)	4007(2)	2143(2)	6352(2)	21(1)
C(19)	3856(2)	1966(2)	7304(2)	22(1)
C(20)	3269(2)	1270(2)	7598(3)	36(1)
C(21)	3898(2)	567(2)	7752(3)	38(1)
C(22)	7436(2)	3423(1)	6936(2)	20(1)
C(23)	7598(2)	3431(2)	5957(2)	27(1)
C(24)	8537(2)	3596(2)	5612(2)	35(1)
C(25)	9340(2)	3748(2)	6238(3)	36(1)
C(26)	9204(2)	3743(2)	7206(3)	32(1)

C(27)	8264(2)	3586(2)	7555(2)	25(1)
C(28)	5920(2)	3407(1)	8926(2)	19(1)
C(29)	6307(2)	4088(2)	9271(2)	21(1)
C(30)	6066(2)	4356(2)	10165(2)	25(1)
C(31)	5430(2)	3957(2)	10723(2)	28(1)
C(32)	5051(2)	3272(2)	10405(2)	28(1)
C(33)	5300(2)	3004(2)	9519(2)	25(1)
C(34)	4962(2)	3957(1)	6854(2)	17(1)
C(35)	4757(2)	4043(2)	5878(2)	24(1)
C(36)	4102(2)	4591(2)	5521(2)	28(1)
C(37)	3633(2)	5068(2)	6135(2)	31(1)
C(38)	3822(2)	4998(2)	7105(2)	32(1)
C(39)	4478(2)	4444(2)	7458(2)	23(1)

Table 3. Bond lengths [$\text{\AA}$] and angles [$^\circ$] for d18124_a.

Ir(1)-N(1)	2.095(2)
Ir(1)-O(1)	2.1592(19)
Ir(1)-C(18)	2.166(2)
Ir(1)-C(19)	2.180(3)
Ir(1)-C(14)	2.294(3)
Ir(1)-C(15)	2.310(3)
Ir(1)-Sn(1)	2.6137(2)
Ir(1)-H(1IR)	1.37(3)
Sn(1)-C(28)	2.160(3)
Sn(1)-C(22)	2.166(3)
Sn(1)-C(34)	2.182(2)
O(1)-C(1)	1.286(3)
O(2)-C(3)	1.365(3)
O(2)-C(4)	1.445(3)
N(1)-C(3)	1.302(3)
N(1)-C(5)	1.505(3)
C(1)-C(2)	1.377(4)
C(1)-C(6)	1.505(4)
C(2)-C(3)	1.411(4)
C(2)-H(2A)	0.9500
C(4)-C(5)	1.529(4)
C(4)-H(4A)	0.9900
C(4)-H(4B)	0.9900
C(5)-C(12)	1.514(4)
C(5)-C(13)	1.521(4)
C(6)-C(7)	1.392(4)
C(6)-C(11)	1.397(4)
C(7)-C(8)	1.388(4)
C(7)-H(7A)	0.9500
C(8)-C(9)	1.383(4)
C(8)-H(8A)	0.9500
C(9)-C(10)	1.380(4)
C(9)-H(9A)	0.9500
C(10)-C(11)	1.382(4)

C(10)-H(10A)	0.9500
C(11)-H(11A)	0.9500
C(12)-H(12A)	0.9800
C(12)-H(12B)	0.9800
C(12)-H(12C)	0.9800
C(13)-H(13A)	0.9800
C(13)-H(13B)	0.9800
C(13)-H(13C)	0.9800
C(14)-C(15)	1.377(4)
C(14)-C(21)	1.501(4)
C(14)-H(14A)	1.0000
C(15)-C(16)	1.521(4)
C(15)-H(15A)	1.0000
C(16)-C(17)	1.521(4)
C(16)-H(16A)	0.9900
C(16)-H(16B)	0.9900
C(17)-C(18)	1.508(4)
C(17)-H(17A)	0.9900
C(17)-H(17B)	0.9900
C(18)-C(19)	1.393(4)
C(18)-H(18A)	1.0000
C(19)-C(20)	1.531(4)
C(19)-H(19A)	1.0000
C(20)-C(21)	1.517(4)
C(20)-H(20A)	0.9900
C(20)-H(20B)	0.9900
C(21)-H(21A)	0.9900
C(21)-H(21B)	0.9900
C(22)-C(23)	1.398(4)
C(22)-C(27)	1.402(4)
C(23)-C(24)	1.390(4)
C(23)-H(23A)	0.9500
C(24)-C(25)	1.378(5)
C(24)-H(24A)	0.9500
C(25)-C(26)	1.377(5)
C(25)-H(25A)	0.9500

C(26)-C(27)	1.391(4)
C(26)-H(26A)	0.9500
C(27)-H(27A)	0.9500
C(28)-C(33)	1.396(4)
C(28)-C(29)	1.397(4)
C(29)-C(30)	1.392(4)
C(29)-H(29A)	0.9500
C(30)-C(31)	1.376(4)
C(30)-H(30A)	0.9500
C(31)-C(32)	1.387(4)
C(31)-H(31A)	0.9500
C(32)-C(33)	1.385(4)
C(32)-H(32A)	0.9500
C(33)-H(33A)	0.9500
C(34)-C(39)	1.391(4)
C(34)-C(35)	1.391(4)
C(35)-C(36)	1.389(4)
C(35)-H(35A)	0.9500
C(36)-C(37)	1.379(4)
C(36)-H(36A)	0.9500
C(37)-C(38)	1.376(4)
C(37)-H(37A)	0.9500
C(38)-C(39)	1.394(4)
C(38)-H(38A)	0.9500
C(39)-H(39A)	0.9500

N(1)-Ir(1)-O(1)	87.95(8)
N(1)-Ir(1)-C(18)	165.61(10)
O(1)-Ir(1)-C(18)	78.79(9)
N(1)-Ir(1)-C(19)	155.50(10)
O(1)-Ir(1)-C(19)	116.16(9)
C(18)-Ir(1)-C(19)	37.39(10)
N(1)-Ir(1)-C(14)	86.95(9)
O(1)-Ir(1)-C(14)	114.38(9)
C(18)-Ir(1)-C(14)	93.41(10)
C(19)-Ir(1)-C(14)	79.33(10)

N(1)-Ir(1)-C(15)	92.83(9)
O(1)-Ir(1)-C(15)	80.36(10)
C(18)-Ir(1)-C(15)	79.56(10)
C(19)-Ir(1)-C(15)	87.41(11)
C(14)-Ir(1)-C(15)	34.79(11)
N(1)-Ir(1)-Sn(1)	93.55(6)
O(1)-Ir(1)-Sn(1)	86.42(5)
C(18)-Ir(1)-Sn(1)	91.19(7)
C(19)-Ir(1)-Sn(1)	92.31(7)
C(14)-Ir(1)-Sn(1)	159.20(8)
C(15)-Ir(1)-Sn(1)	165.09(8)
N(1)-Ir(1)-H(1IR)	84.6(13)
O(1)-Ir(1)-H(1IR)	165.3(13)
C(18)-Ir(1)-H(1IR)	109.6(13)
C(19)-Ir(1)-H(1IR)	72.8(13)
C(14)-Ir(1)-H(1IR)	78.0(13)
C(15)-Ir(1)-H(1IR)	112.6(13)
Sn(1)-Ir(1)-H(1IR)	81.4(13)
C(28)-Sn(1)-C(22)	108.56(10)
C(28)-Sn(1)-C(34)	97.70(10)
C(22)-Sn(1)-C(34)	103.76(10)
C(28)-Sn(1)-Ir(1)	116.66(7)
C(22)-Sn(1)-Ir(1)	114.89(7)
C(34)-Sn(1)-Ir(1)	113.26(6)
C(1)-O(1)-Ir(1)	125.28(17)
C(3)-O(2)-C(4)	106.4(2)
C(3)-N(1)-C(5)	108.5(2)
C(3)-N(1)-Ir(1)	122.24(18)
C(5)-N(1)-Ir(1)	129.15(17)
O(1)-C(1)-C(2)	125.8(3)
O(1)-C(1)-C(6)	113.7(2)
C(2)-C(1)-C(6)	120.5(2)
C(1)-C(2)-C(3)	124.7(3)
C(1)-C(2)-H(2A)	117.6
C(3)-C(2)-H(2A)	117.6
N(1)-C(3)-O(2)	114.2(2)

N(1)-C(3)-C(2)	130.7(2)
O(2)-C(3)-C(2)	115.1(2)
O(2)-C(4)-C(5)	104.6(2)
O(2)-C(4)-H(4A)	110.8
C(5)-C(4)-H(4A)	110.8
O(2)-C(4)-H(4B)	110.8
C(5)-C(4)-H(4B)	110.8
H(4A)-C(4)-H(4B)	108.9
N(1)-C(5)-C(12)	114.0(2)
N(1)-C(5)-C(13)	109.9(2)
C(12)-C(5)-C(13)	110.6(3)
N(1)-C(5)-C(4)	100.1(2)
C(12)-C(5)-C(4)	111.0(2)
C(13)-C(5)-C(4)	110.9(2)
C(7)-C(6)-C(11)	118.2(3)
C(7)-C(6)-C(1)	118.7(3)
C(11)-C(6)-C(1)	123.1(3)
C(8)-C(7)-C(6)	120.9(3)
C(8)-C(7)-H(7A)	119.6
C(6)-C(7)-H(7A)	119.6
C(9)-C(8)-C(7)	120.0(3)
C(9)-C(8)-H(8A)	120.0
C(7)-C(8)-H(8A)	120.0
C(10)-C(9)-C(8)	119.8(3)
C(10)-C(9)-H(9A)	120.1
C(8)-C(9)-H(9A)	120.1
C(9)-C(10)-C(11)	120.3(3)
C(9)-C(10)-H(10A)	119.9
C(11)-C(10)-H(10A)	119.9
C(10)-C(11)-C(6)	120.9(3)
C(10)-C(11)-H(11A)	119.6
C(6)-C(11)-H(11A)	119.6
C(5)-C(12)-H(12A)	109.5
C(5)-C(12)-H(12B)	109.5
H(12A)-C(12)-H(12B)	109.5
C(5)-C(12)-H(12C)	109.5

H(12A)-C(12)-H(12C)	109.5
H(12B)-C(12)-H(12C)	109.5
C(5)-C(13)-H(13A)	109.5
C(5)-C(13)-H(13B)	109.5
H(13A)-C(13)-H(13B)	109.5
C(5)-C(13)-H(13C)	109.5
H(13A)-C(13)-H(13C)	109.5
H(13B)-C(13)-H(13C)	109.5
C(15)-C(14)-C(21)	125.8(3)
C(15)-C(14)-Ir(1)	73.23(16)
C(21)-C(14)-Ir(1)	108.88(19)
C(15)-C(14)-H(14A)	113.8
C(21)-C(14)-H(14A)	113.8
Ir(1)-C(14)-H(14A)	113.8
C(14)-C(15)-C(16)	123.5(3)
C(14)-C(15)-Ir(1)	71.98(16)
C(16)-C(15)-Ir(1)	110.25(18)
C(14)-C(15)-H(15A)	114.5
C(16)-C(15)-H(15A)	114.5
Ir(1)-C(15)-H(15A)	114.5
C(17)-C(16)-C(15)	114.2(2)
C(17)-C(16)-H(16A)	108.7
C(15)-C(16)-H(16A)	108.7
C(17)-C(16)-H(16B)	108.7
C(15)-C(16)-H(16B)	108.7
H(16A)-C(16)-H(16B)	107.6
C(18)-C(17)-C(16)	115.0(2)
C(18)-C(17)-H(17A)	108.5
C(16)-C(17)-H(17A)	108.5
C(18)-C(17)-H(17B)	108.5
C(16)-C(17)-H(17B)	108.5
H(17A)-C(17)-H(17B)	107.5
C(19)-C(18)-C(17)	126.0(3)
C(19)-C(18)-Ir(1)	71.86(15)
C(17)-C(18)-Ir(1)	111.31(18)
C(19)-C(18)-H(18A)	113.4

C(17)-C(18)-H(18A)	113.4
Ir(1)-C(18)-H(18A)	113.4
C(18)-C(19)-C(20)	122.5(3)
C(18)-C(19)-Ir(1)	70.75(15)
C(20)-C(19)-Ir(1)	113.75(18)
C(18)-C(19)-H(19A)	114.2
C(20)-C(19)-H(19A)	114.2
Ir(1)-C(19)-H(19A)	114.2
C(21)-C(20)-C(19)	115.0(2)
C(21)-C(20)-H(20A)	108.5
C(19)-C(20)-H(20A)	108.5
C(21)-C(20)-H(20B)	108.5
C(19)-C(20)-H(20B)	108.5
H(20A)-C(20)-H(20B)	107.5
C(14)-C(21)-C(20)	114.1(3)
C(14)-C(21)-H(21A)	108.7
C(20)-C(21)-H(21A)	108.7
C(14)-C(21)-H(21B)	108.7
C(20)-C(21)-H(21B)	108.7
H(21A)-C(21)-H(21B)	107.6
C(23)-C(22)-C(27)	117.0(3)
C(23)-C(22)-Sn(1)	120.1(2)
C(27)-C(22)-Sn(1)	122.8(2)
C(24)-C(23)-C(22)	121.5(3)
C(24)-C(23)-H(23A)	119.2
C(22)-C(23)-H(23A)	119.2
C(25)-C(24)-C(23)	120.2(3)
C(25)-C(24)-H(24A)	119.9
C(23)-C(24)-H(24A)	119.9
C(26)-C(25)-C(24)	119.7(3)
C(26)-C(25)-H(25A)	120.2
C(24)-C(25)-H(25A)	120.2
C(25)-C(26)-C(27)	120.4(3)
C(25)-C(26)-H(26A)	119.8
C(27)-C(26)-H(26A)	119.8
C(26)-C(27)-C(22)	121.2(3)

C(26)-C(27)-H(27A)	119.4
C(22)-C(27)-H(27A)	119.4
C(33)-C(28)-C(29)	117.4(3)
C(33)-C(28)-Sn(1)	119.44(19)
C(29)-C(28)-Sn(1)	121.8(2)
C(30)-C(29)-C(28)	121.0(3)
C(30)-C(29)-H(29A)	119.5
C(28)-C(29)-H(29A)	119.5
C(31)-C(30)-C(29)	120.4(3)
C(31)-C(30)-H(30A)	119.8
C(29)-C(30)-H(30A)	119.8
C(30)-C(31)-C(32)	119.8(3)
C(30)-C(31)-H(31A)	120.1
C(32)-C(31)-H(31A)	120.1
C(33)-C(32)-C(31)	119.7(3)
C(33)-C(32)-H(32A)	120.2
C(31)-C(32)-H(32A)	120.2
C(32)-C(33)-C(28)	121.8(3)
C(32)-C(33)-H(33A)	119.1
C(28)-C(33)-H(33A)	119.1
C(39)-C(34)-C(35)	117.0(2)
C(39)-C(34)-Sn(1)	120.42(19)
C(35)-C(34)-Sn(1)	122.6(2)
C(36)-C(35)-C(34)	121.6(3)
C(36)-C(35)-H(35A)	119.2
C(34)-C(35)-H(35A)	119.2
C(37)-C(36)-C(35)	120.2(3)
C(37)-C(36)-H(36A)	119.9
C(35)-C(36)-H(36A)	119.9
C(38)-C(37)-C(36)	119.5(3)
C(38)-C(37)-H(37A)	120.2
C(36)-C(37)-H(37A)	120.2
C(37)-C(38)-C(39)	120.0(3)
C(37)-C(38)-H(38A)	120.0
C(39)-C(38)-H(38A)	120.0
C(34)-C(39)-C(38)	121.7(3)

C(34)-C(39)-H(39A)	119.2
C(38)-C(39)-H(39A)	119.2

Symmetry transformations used to generate equivalent atoms:

Table 4. Anisotropic displacement parameters ($\text{\AA}^2 \times 10^3$) for d18124_a. The anisotropic displacement factor exponent takes the form: $-2\pi^2 [h^2 a^{*2} U^{11} + \dots + 2 h k a^* b^* U^{12}]$

	U^{11}	U^{22}	U^{33}	U^{23}	U^{13}	U^{12}
Ir(1)	15(1)	12(1)	20(1)	-2(1)	-1(1)	0(1)
Sn(1)	16(1)	13(1)	17(1)	-1(1)	-1(1)	0(1)
O(1)	21(1)	22(1)	20(1)	-1(1)	2(1)	4(1)
O(2)	26(1)	41(1)	27(1)	3(1)	2(1)	18(1)
N(1)	16(1)	16(1)	20(1)	1(1)	-2(1)	3(1)
C(1)	21(1)	19(1)	22(1)	-2(1)	2(1)	-2(1)
C(2)	28(2)	32(2)	20(2)	-3(1)	3(1)	10(1)
C(3)	19(1)	20(1)	27(2)	-1(1)	0(1)	6(1)
C(4)	32(2)	30(2)	28(2)	4(1)	-2(1)	14(1)
C(5)	19(1)	19(1)	21(1)	0(1)	-2(1)	3(1)
C(6)	19(1)	25(1)	22(2)	0(1)	1(1)	-4(1)
C(7)	27(2)	27(2)	23(2)	0(1)	-1(1)	-1(1)
C(8)	32(2)	30(2)	29(2)	5(1)	-3(1)	-1(1)
C(9)	28(2)	39(2)	24(2)	5(1)	-1(1)	-10(1)
C(10)	27(2)	39(2)	24(2)	-4(1)	7(1)	-4(1)
C(11)	26(2)	32(2)	26(2)	-1(1)	2(1)	1(1)
C(12)	29(2)	45(2)	29(2)	15(1)	-2(1)	-2(1)
C(13)	36(2)	22(2)	39(2)	-1(1)	-18(2)	4(1)
C(14)	26(2)	12(1)	47(2)	1(1)	-4(1)	-2(1)
C(15)	29(2)	17(1)	44(2)	-15(1)	-5(1)	0(1)
C(16)	46(2)	26(2)	45(2)	-10(1)	-18(2)	0(1)
C(17)	24(2)	29(2)	34(2)	-7(1)	-9(1)	4(1)
C(18)	13(1)	18(1)	30(2)	-3(1)	-4(1)	1(1)
C(19)	12(1)	21(1)	33(2)	-2(1)	0(1)	4(1)
C(20)	24(2)	35(2)	50(2)	13(2)	7(2)	-2(1)
C(21)	34(2)	26(2)	53(2)	6(2)	5(2)	-8(1)
C(22)	18(1)	13(1)	29(2)	0(1)	1(1)	-1(1)
C(23)	27(2)	22(1)	31(2)	0(1)	2(1)	1(1)
C(24)	36(2)	29(2)	41(2)	5(1)	15(2)	2(1)
C(25)	22(2)	25(2)	63(2)	9(2)	12(2)	1(1)
C(26)	20(2)	19(1)	58(2)	3(1)	-6(2)	-1(1)

C(27)	25(2)	15(1)	36(2)	1(1)	-3(1)	1(1)
C(28)	21(1)	17(1)	18(1)	-1(1)	-4(1)	4(1)
C(29)	21(1)	21(1)	22(2)	0(1)	-2(1)	2(1)
C(30)	29(2)	20(1)	26(2)	-8(1)	-4(1)	2(1)
C(31)	32(2)	31(2)	20(2)	-5(1)	1(1)	7(1)
C(32)	31(2)	31(2)	22(2)	4(1)	3(1)	0(1)
C(33)	32(2)	20(1)	22(2)	0(1)	-2(1)	-2(1)
C(34)	16(1)	14(1)	22(1)	1(1)	-3(1)	-3(1)
C(35)	26(2)	21(1)	24(2)	-2(1)	0(1)	1(1)
C(36)	28(2)	32(2)	24(2)	6(1)	-6(1)	-1(1)
C(37)	23(2)	26(2)	43(2)	2(1)	-10(1)	6(1)
C(38)	27(2)	34(2)	33(2)	-11(1)	-5(1)	11(1)
C(39)	18(1)	27(2)	24(2)	-5(1)	-5(1)	4(1)

Table 5. Hydrogen coordinates ($\times 10^4$) and isotropic displacement parameters ($\text{\AA}^2 \times 10^{-3}$) for d18124_a.

	x	y	z	U(eq)
H(1IR)	5350(20)	1684(17)	7960(20)	39(9)
H(2A)	7828	1060	5168	32
H(4A)	8880	691	8213	36
H(4B)	8049	71	7902	36
H(7A)	5636	2761	4154	31
H(8A)	5605	3066	2536	37
H(9A)	6522	2352	1481	36
H(10A)	7468	1333	2050	36
H(11A)	7527	1039	3667	34
H(12A)	6533	244	8626	52
H(12B)	6198	1037	9042	52
H(12C)	7182	626	9484	52
H(13A)	7305	2173	8780	49
H(13B)	8265	2085	8140	49
H(13C)	8270	1718	9182	49
H(14A)	5338	174	7416	34
H(15A)	5373	381	5874	36
H(16A)	4008	648	4924	47
H(16B)	3326	586	5832	47
H(17A)	3994	1883	4927	35
H(17B)	2935	1739	5389	35
H(18A)	4008	2694	6224	25
H(19A)	3767	2411	7724	26
H(20A)	2931	1385	8196	43
H(20B)	2738	1167	7098	43
H(21A)	4110	533	8436	45
H(21B)	3473	125	7593	45
H(23A)	7054	3320	5518	32
H(24A)	8625	3605	4943	42
H(25A)	9983	3855	6002	44

H(26A)	9757	3847	7639	39
H(27A)	8181	3589	8225	30
H(29A)	6741	4372	8889	26
H(30A)	6342	4818	10391	30
H(31A)	5251	4149	11325	33
H(32A)	4622	2989	10793	34
H(33A)	5041	2532	9310	29
H(35A)	5073	3718	5445	29
H(36A)	3976	4638	4851	33
H(37A)	3183	5443	5891	38
H(38A)	3505	5327	7533	38
H(39A)	4598	4399	8129	28

Table 6. Torsion angles [°] for d18124_a.

Ir(1)-O(1)-C(1)-C(2)	-12.7(4)
Ir(1)-O(1)-C(1)-C(6)	168.62(16)
O(1)-C(1)-C(2)-C(3)	-4.3(5)
C(6)-C(1)-C(2)-C(3)	174.3(3)
C(5)-N(1)-C(3)-O(2)	6.9(3)
Ir(1)-N(1)-C(3)-O(2)	-175.93(17)
C(5)-N(1)-C(3)-C(2)	-172.0(3)
Ir(1)-N(1)-C(3)-C(2)	5.2(4)
C(4)-O(2)-C(3)-N(1)	9.8(3)
C(4)-O(2)-C(3)-C(2)	-171.1(2)
C(1)-C(2)-C(3)-N(1)	8.8(5)
C(1)-C(2)-C(3)-O(2)	-170.0(3)
C(3)-O(2)-C(4)-C(5)	-21.6(3)
C(3)-N(1)-C(5)-C(12)	-137.7(3)
Ir(1)-N(1)-C(5)-C(12)	45.3(3)
C(3)-N(1)-C(5)-C(13)	97.5(3)
Ir(1)-N(1)-C(5)-C(13)	-79.4(3)
C(3)-N(1)-C(5)-C(4)	-19.2(3)
Ir(1)-N(1)-C(5)-C(4)	163.85(18)
O(2)-C(4)-C(5)-N(1)	24.0(3)
O(2)-C(4)-C(5)-C(12)	144.7(2)
O(2)-C(4)-C(5)-C(13)	-91.9(3)
O(1)-C(1)-C(6)-C(7)	17.8(4)
C(2)-C(1)-C(6)-C(7)	-161.0(3)
O(1)-C(1)-C(6)-C(11)	-160.8(3)
C(2)-C(1)-C(6)-C(11)	20.4(4)
C(11)-C(6)-C(7)-C(8)	-1.2(4)
C(1)-C(6)-C(7)-C(8)	-179.8(3)
C(6)-C(7)-C(8)-C(9)	0.9(4)
C(7)-C(8)-C(9)-C(10)	0.1(4)
C(8)-C(9)-C(10)-C(11)	-0.7(4)
C(9)-C(10)-C(11)-C(6)	0.4(4)
C(7)-C(6)-C(11)-C(10)	0.5(4)
C(1)-C(6)-C(11)-C(10)	179.1(3)

C(21)-C(14)-C(15)-C(16)	-1.3(4)
Ir(1)-C(14)-C(15)-C(16)	-102.8(3)
C(21)-C(14)-C(15)-Ir(1)	101.5(3)
C(14)-C(15)-C(16)-C(17)	89.5(4)
Ir(1)-C(15)-C(16)-C(17)	8.3(4)
C(15)-C(16)-C(17)-C(18)	-28.9(4)
C(16)-C(17)-C(18)-C(19)	-47.0(4)
C(16)-C(17)-C(18)-Ir(1)	35.6(3)
C(17)-C(18)-C(19)-C(20)	-2.9(4)
Ir(1)-C(18)-C(19)-C(20)	-106.5(2)
C(17)-C(18)-C(19)-Ir(1)	103.6(3)
C(18)-C(19)-C(20)-C(21)	89.6(4)
Ir(1)-C(19)-C(20)-C(21)	8.0(4)
C(15)-C(14)-C(21)-C(20)	-49.5(4)
Ir(1)-C(14)-C(21)-C(20)	33.1(3)
C(19)-C(20)-C(21)-C(14)	-28.3(4)
C(27)-C(22)-C(23)-C(24)	-0.3(4)
Sn(1)-C(22)-C(23)-C(24)	-176.4(2)
C(22)-C(23)-C(24)-C(25)	0.9(4)
C(23)-C(24)-C(25)-C(26)	-0.7(5)
C(24)-C(25)-C(26)-C(27)	0.0(4)
C(25)-C(26)-C(27)-C(22)	0.6(4)
C(23)-C(22)-C(27)-C(26)	-0.4(4)
Sn(1)-C(22)-C(27)-C(26)	175.5(2)
C(33)-C(28)-C(29)-C(30)	-1.0(4)
Sn(1)-C(28)-C(29)-C(30)	165.3(2)
C(28)-C(29)-C(30)-C(31)	-0.7(4)
C(29)-C(30)-C(31)-C(32)	1.9(4)
C(30)-C(31)-C(32)-C(33)	-1.3(4)
C(31)-C(32)-C(33)-C(28)	-0.6(4)
C(29)-C(28)-C(33)-C(32)	1.7(4)
Sn(1)-C(28)-C(33)-C(32)	-165.0(2)
C(39)-C(34)-C(35)-C(36)	0.2(4)
Sn(1)-C(34)-C(35)-C(36)	179.7(2)
C(34)-C(35)-C(36)-C(37)	-0.1(4)
C(35)-C(36)-C(37)-C(38)	0.2(5)

C(36)-C(37)-C(38)-C(39)	-0.4(5)
C(35)-C(34)-C(39)-C(38)	-0.4(4)
Sn(1)-C(34)-C(39)-C(38)	-179.9(2)
C(37)-C(38)-C(39)-C(34)	0.5(5)

Symmetry transformations used to generate equivalent atoms:

References

- (1) Witte, H.; Seeliger, W. *Angew. Chem. Int. Ed. Engl.* **1972**, *11*, 287–288.
- (2) Evans, D. A.; Peterson, G. S.; Johnson, J. S.; Barnes, D. M.; Campos, K. R.; Woerpel, K. A. *J. Org. Chem.* **1998**, *63*, 4541–4544.
- (3) Schwekendiek, K.; Glorius, F. *Synthesis* **2006**, 2996–3002.
- (4) Wiley, R. H.; Bennett, L. L. *Chem. Rev.* **1949**, *44*, 447–476.
- (5) Hoogenboom, R. *Angew. Chem. Int. Ed. Engl.* **2009**, *48*, 7978–7994.
- (6) Meyers, A. I.; Mihelich, D. M. *Angew. Chem. Int. Ed. Engl.* **1976**, *15*, 270–281.
- (7) Gant, T. G.; Meyers, A. I.; Collins, F.; Oxazolines, A.; Metalation, C.; Oxazolines, A.; Metalations, A. *Tetrahedron* **1994**, *50*, 2297–2360.
- (8) Tohda, Y.; Yanagidani, T.; Asaka, N. *Bull. Chem. Soc. Jpn.* **2016**, *89*, 810–822.
- (9) Song, Y.; De Silva, H. I.; Henry, W. P.; Ye, G.; Chatterjee, S.; Pittman, C. U. *Tetrahedron Lett.* **2011**, *52*, 4507–4511.
- (10) Van Kirk, C. C.; Béreau, V.; Abu-Omar, M. M.; Evans, D. H. *J. Electroanal. Chem.* **2003**, *541*, 31–38.
- (11) Gomez-Simon, M.; Jansat, S.; Muller, G.; Panyella, D.; Font-Bardia, M.; Solans, X. *J. Chem. Soc., Dalton. Trans.* **1997**, 3755–3764.
- (12) Gold, V. *IUPAC Compend. Chem. Terminol.* **2014**, 1670.
- (13) Gossage, R. A.; Jenkins, H. A.; Yadav, P. N. *Tetrahedron Lett.* **2004**, *45*, 7689–7691.
- (14) Decken, A.; Eisnor, C. R.; Gossage, R. A.; Jackson, S. M. *Inorg. Chim. Acta* **2006**, *359*, 1743–1753.
- (15) Barclay, T. M.; Río, I. del; Gossage, R. A.; Jackson, S. M. *Can. J. Chem.* **2003**, *81*,

1482–1491.

- (16) Eisnor, C. R.; Gossage, R. A.; Yadav, P. N. *Tetrahedron* **2006**, *62*, 3395–3401.
- (17) Takayama, T.; Nakazawa, J.; Hikichi, S. *Acta Crystallogr. Sect. C Struct. Chem.* **2016**, *72*, 842–845.
- (18) Zhang, J.; Pattacini, R.; Braunstein, P. *Inorg. Chem.* **2009**, *48*, 11954–11962.
- (19) Herasymchuk, K.; Huynh, J.; Lough, A.; Roces Fernández, L.; Gossage, R. *Synthesis* **2016**, *48*, 2121–2129.
- (20) Jones, R. C.; Herasymchuk, K.; Mahdi, T.; Petrov, A.; Resanovic, S.; Vaughan, D. G.; Lough, A. J.; Quail, J. W.; Koivisto, B. D.; Wylie, R. S.; Gossage, R. A. *Org. Biomol. Chem.* **2013**, *11*, 3484–3493.
- (21) Drover, M. W.; Love, J. A.; Schafer, L. L.; Goze, C.; Brandès, S.; Guillard, R.; Zrineh, A.; Denat, F.; Ray, K.; Company, A. *Chem. Soc. Rev.* **2017**, *373*, 150–158.
- (22) Sun, Y.; Chan, H. S.; Dixneuf, P. H.; Xie, Z. *J. Organomet. Chem.* **2006**, *691*, 3071–3082.
- (23) Karpin, G. W.; Merola, J. S.; Falkinham, J. O. *Antimicrob. Agents Chemother.* **2013**, *57*, 3434–3436.
- (24) Molodkin, A. K.; Esina, N. Y.; Kurasova, M. N. *Russ. J. Inorg. Chem.* **2010**, *55*, 370–376.
- (25) Varshavsky, Y. S.; Galding, M. R.; Cherkasova, T. G.; Smirnov, S. N.; Khrustalev, V. N. *J. Organomet. Chem.* **2007**, *692*, 5788–5794.
- (26) Thai, T. T.; Therrien, B.; Süß-Fink, G. *Inorg. Chem. Commun.* **2009**, *12*, 806–807.
- (27) Vasam, C. S.; Modem, S.; Kankala, S.; Kanne, S.; Budige, G.; Vadde, R. *Cent. Eur. J. Chem.* **2010**, *8*, 77–86.

- (28) Gorol, M.; Roesky, H. W.; Noltemeyer, M.; Schmidt, H. G. *Eur. J. Inorg. Chem.* **2005**, 4840–4844.
- (29) Heras, J. V.; Pinilla, E.; Martinez, M. *Polyhedron* **1983**, 2, 1009–1012.
- (30) Menendez Rodriguez, G.; Bucci, A.; Hutchinson, R.; Bellachioma, G.; Zuccaccia, C.; Giovagnoli, S.; Idriss, H.; Macchioni, A. *ACS Energy Lett.* **2017**, 2, 105–110.
- (31) Zhang, T.; Wang, W.; Gu, X.; Shi, M. *Organometallics* **2008**, 27, 753–757.
- (32) Drover, M. W.; Bowes, E. G.; Love, J. A.; Schafer, L. L. *Organometallics* **2017**, 36, 331–341.
- (33) Drover, M. W.; Johnson, H. C.; Schafer, L. L.; Love, J. A.; Weller, A. S. *Organometallics* **2015**, 34, 3849–3856.
- (34) Meyers, A. I.; Temple, D. L.; Nolen, R. L.; Mihelich, E. D. *J. Org. Chem.* **1974**, 39, 2778–2783.
- (35) Tohda, Y.; Kawashima, T.; Ariga, M.; Akiyama, R.; Shudoh, H.; Mori, Y. *Bull. Chem. Soc. Jpn.* **1984**, 57, 2329–2330.
- (36) Tohda, Y.; Morikawa, M.; Kawashima, T.; Ariga, M.; Mori, Y. *Chem. Lett.* **1986**, 272–274.
- (37) Tohda, Y.; Yanagidani, T.; Hiramatsu, S. I.; Nishiwaki, N.; Tani, K.; Imagawa, K.; Ariga, M. *Bull. Chem. Soc. Jpn.* 1997, pp 2781–2790.
- (38) Chanu, L. G.; Singh, O. M.; Jang, S.-H.; Lee, S.-G. *Bull. Korean Chem. Soc.* **2010**, 31, 859–862.
- (39) Kumar, G. M. V. N. A. R. R.; Thangamani, A. *Orient. J. Chem.* **2017**, 33, 1555–1562.
- (40) Huang, Z.; Zhang, P. *Chem. Ber.* **1989**, 122, 2011–2016.

- (41) Bergs, R.; Krämer, R.; Maurus, M.; Schreiner, B.; Urban, R.; Christopher, M.; Polborn, K.; Siinkel, K.; Beck, W. *Naturforsch.* **1996**, 1–14.
- (42) Kramer, R.; Maurus, M.; Bergs, R.; Polborn, K.; Siinkel, K.; Wagner, B.; Beck, W. *Chem. Ber.* **1993**, 126, 1969–1980.
- (43) Lampeka, R.; Bergs, R.; Kramer, R.; Polborn, K.; Beck, W. *J. Nat. Res.* **1994**, 49, 225–232.
- (44) Yuwen, J.; Chakraborty, S.; Brennessel, W. W.; Jones, W. D. *ACS Catal.* **2017**, 7, 3735–3740.
- (45) Tokmic, K.; Jackson, B. J.; Salazar, A.; Woods, T. J.; Fout, A. R. *J. Am. Chem. Soc.* **2017**, 139, 13554–13561.
- (46) Bertilsson, S. K.; Tendenborg, L.; Alonso, D. A.; Andersson, P. G. *Organometallics* **1999**, 18, 1281–1286.
- (47) García, F.; Perles, J.; Zamora, F.; Amo-Ochoa, P. *Inorg. Chim. Acta* **2016**, 453, 574–582.
- (48) El-Saied, F. A.; Salem, T. A.; Shakhofa, M. M. E.; Al-Hakimi, A. N.; Radwan, A. S. *Beni-Suef Univ. J. Basic Appl. Sci.* **2017**.
- (49) Van den Berg, M.; Minnaard, A. J.; Schudde, E. P.; Van Esch, J.; De Vries, A. H. M.; De Vries, J. G.; Feringa, B. L. *J. Am. Chem. Soc.* **2000**, 122, 11539–11540.
- (50) Burk, M. J.; Feaster, J. E. *J. Am. Chem. Soc.* **1992**, 114, 6266–6267.
- (51) Kang, G.; Cullen, W. R.; Fryzuk, M. D.; James, B. R.; Kutney, J. P. *J. Chem. Soc. Chem. Commun.* **1988**, 0, 1466–1467.
- (52) Morimoto, T.; Chiba, M.; Achiwa, K. *Chem. Pharm. Bull.* **1992**, 40, 2894–2896.
- (53) Perea-Buceta, J. E.; Fernández, I.; Heikkinen, S.; Axenov, K.; King, A. W. T.;

- Niemi, T.; Nieger, M.; Leskelä, M.; Repo, T. *Angew. Chem. Int. Ed. Engl.* **2015**, *54*, 14321–14325.
- (54) Carlton, L.; Molapisi, J. J. *J. Organomet. Chem.* **2000**, *609*, 60–65.
- (55) Drover, M. W.; Schafer, L. L.; Love, J. A. *Angew. Chem. Int. Ed. Engl.* **2016**, *55*, 3181–3186.
- (56) Herde, J. L.; Lambert, J. C.; Senoff, C. V. *Inorg. Synth.* **1974**, *15*, 18–20.
- (57) Rane, S. S.; Choi, P. *Chem. Mater.* **2005**, *17*, 926.
- (58) Odian, G. *Principles of Polymerization*, 4th ed.; JOHN WILEY & SONS, INC.,: New York, 2004.
- (59) Kato, M.; Kamigaito, M.; Sawamoto, M.; Higashimuras, T. *Macromolecules* **1996**, *28*, 1721–1723.
- (60) Wang, J. S.; Matyjaszewski, K. *J. Am. Chem. Soc.* **1995**, *117*, 5614–5615.
- (61) Matyjaszewski, K.; Xia, J. *Chem. Rev.* **2001**, *101*, 2921–2990.
- (62) Ribelli, T. G.; Fantin, M.; Daran, J. C.; Augustine, K. F.; Poli, R.; Matyjaszewski, K. *J. Am. Chem. Soc.* **2018**, *140*, 1525–1534.
- (63) Kaneyoshi, H.; Matyjaszewski, K. *J. Polym. Sci. Part A Polym. Chem.* **2006**, *45*, 477–459.
- (64) Zhang, Z.; Zhang, W.; Zhu, X.; Cheng, Z.; Zhu, J. *J. Polym. Sci. Part A Polym. Chem.* **2007**, *45*, 5722–5730.
- (65) Yan, Y.; Zhang, J.; Wilbon, P.; Qiao, Y.; Tang, C. *Macromol. Rapid Commun.* **2014**, *35*, 1840–1845.
- (66) Resanović, S.; Gossage, R. A. Unpublished Results, 2009.
- (67) Adjei, J. A. Nickel and Palladium (N,O)-Donor Ligand Complexes: Synthesis,

Structure & Catalysis, Ryerson University, 2018.

- (68) Kruck, M.; Sauer, D. C.; Enders, M.; Wadepohl, H.; Gade, L. H. *Dalton Trans.* **2011**, 40, 10406–10415.
- (69) Okuniewski, A.; Rosiak, D.; Chojnacki, J.; Becker, B. *Polyhedron* **2015**, 90, 47–57.
- (70) Yang, L.; Powell, D. R.; Houser, R. P. *Dalton Trans.* **2007**, 955–964.
- (71) Ghandi, M.; Feizi, S.; Ziaie, F.; Fazaeli, Y.; Notash, B. *Ann. Nucl. Med.* **2014**, 28, 880–890.
- (72) Baerlocher, F. J.; Bucur, R.; Decken, A.; Eisnor, C. R.; Gossage, R. A.; Jackson, S. M.; Jolly, L.; Wheaton, S. L.; Wylie, R. S. *Aust. J. Chem.* **2010**, 63, 47–55.
- (73) Gurr, P. A.; Mills, M. F.; Qiao, G. G.; Solomon, D. H. *Polymer (Guildf)*. **2005**, 46, 2097–2104.
- (74) Matsubara, K.; Matsumoto, M. *J. Polym. Sci. Part A Polym. Chem.* **2006**, 44, 4222–4228.
- (75) Carlton, L.; Fernandes, M. a; Sitabule, E. *Proc. Natl. Acad. Sci. U. S. A.* **2007**, 104, 6969–6973.
- (76) Herrmann, W. A.; Frey, G. D.; Herdtweck, E.; Steinbeck, M. *Adv. Synth. Catal.* **2007**, 349, 1677–1691.
- (77) Srour, M.; Hadjiali, S.; Sauer, G.; Brunnengräber, K.; Breitzke, H.; Xu, Y.; Weidler, H.; Limbach, H. H.; Gutmann, T.; Buntkowsky, G. *ChemCatChem* **2016**, 8, 3409–3416.
- (78) Nishihara, Y.; Takemura, M.; Osakada, K. *Inorg. Chim. Acta* **2009**, 362, 2951–2956.
- (79) Kirchmann, M.; Fleischhauer, S.; Wesemann, L. *Organometallics* **2008**, 27, 2803–2808.

- (80) Rampersadh, T.; Fernandes, M. A.; Carlton, L. *J. Chem. Crystallogr.* **2014**, 151–160.
- (81) Taubmann, S.; Alt, H. G. *J. Organomet. Chem.* **2008**, 693, 1808–1814.
- (82) Usón, R.; Oro, L. A.; Ciriano, M. A.; Gonzalez, R. *J. Organomet. Chem.* **1981**, 205, 259–271.
- (83) Drover, M. W.; Schafer, L. L.; Love, J. A. *Organometallics* **2015**, 34, 1783–1786.
- (84) Ekengard, E.; Kumar, K.; Fogeron, T.; Kock, C. De; Smith, P. J.; Haukka, M.; Monari, M.; Nordlander, E. *Dalton Trans.* **2016**, 45, 3905–3917.
- (85) Van Der Zeijden, A. A. H.; Van Koten, G.; Wouters, J. M. A.; Wijsmuller, W. F. A.; Grove, D. M.; Spek, A. L.; Smeets, W. J. J. *J. Am. Chem. Soc.* **1988**, 110, 5354–5361.
- (86) Balch, A. L.; Olmstead, M. M.; Oram, D. E.; Reedy, P. E.; Reimer, S. H. *J. Am. Chem. Soc.* **1989**, 111, 4021–4028.
- (87) Petrov, A.; Jones, R. C.; Vaughan, D. G.; Lough, A. J.; Gossage, R. A. *Crystals* **2011**, 1, 229–235.
- (88) Huang, Z.-T.; Zhang, P.-C. *Synth. Commun.* **1990**, 20, 1399–1407.
- (89) Collman, J. P.; Sears Jr, C. T.; Kubota, M. *Inorg. Synth.* **1990**, 92–94.